

# **Analytic tomography of the mantle in a spherically Earth. The technique MZY**

Yoël Lana-Renault

*Geophysics. Department of Theoretical Physics.  
University of Zaragoza. 50009 Zaragoza. Spain  
e-mail: [yoel@kepler.unizar.es](mailto:yoel@kepler.unizar.es)*

**Abstract.** An explicit expression for P-wave velocity is proposed to develop a novel tomographic technique in a spherically symmetric model of the Earth (MZY). The distribution of the P velocity structure in the mantle is determined using only 34 P- and 2 PcP- observed traveltimes. By applying a non-linear inversion, the P-residuals in the range between  $0^\circ$  and  $100^\circ$  are minimised up to a maximum value of 0.015 s. Furthermore, from the high quality computation of PcP traveltimes, with residuals much better than 0.13 s., it is possible to infer the existence of a brief low velocity layer in the D" region. This is then followed by a gradual increasing in the velocity profile towards the core, which begins at a depth of 2893.9 km.

**Key words:** D" shell, Earth's mantle, P-wave velocity, tomography, traveltimes.

## **Introduction**

To date, numerous studies use the arrival times of seismic waves to explore the Earth structure. Seismic arrival times have provided a fundamental constraint on the radial and lateral velocity structure of our planet. Sengupta and Toksoz (1976), Clayton and Comer (1983), Dziewonski (1984) among others, studied the variation of the P-wave velocity in the lower mantle. These works have been extended rapidly to the whole mantle (Pulliam et al., 1993) from many different viewpoints and perspectives, but concluding in almost all cases in interesting correlations with the structure predicted by the plate tectonics. On the other hand, reference models constitute the common basis for all the different studies concerning the Earth. Some of them are fairly relevant and well known in the seismological literature, as PREM (Dziewonski and Anderson, 1981), IASP91 (Kennet and Engdahl, 1991) and SP6 (Morelli and Dziewonski, 1993). They constitute the starting point for a number of applications, including seismic tomography and synthetic seismogram calculations. The strategy of finding an agreement between physical meaningful and

achieving observations is of crucial importance. A decreasing of the relative error between the reproduced and measured data becomes in an increasing knowledge of the main features concerning the Earth structure. In this sense, any effort made to improve the available reference models, will benefit on the current seismological knowledge, especially those concerning local deviations in boundary interfaces in the Earth's interior.

From this viewpoint, in this preliminary work we pretend to improve the fitting of reference traveltimes tables (JB: Jeffreys and Bullen, 1958; BSSA: Herrin et al., 1968) to observed traveltimes and, as consequence of that, to infer the slight deviations of the whole structure with respect to the average models. We have focused our attention on the tomography of the mantle, using and developing a non-linear inversion technique based on the analytical solution of the elliptical integrals involved in the theory of wave propagation. In this sense the approach described in this paper cannot be viewed like an empirical model. We demonstrate that the range of the achievement is large enough and, therefore, the real interpretation is to be an improvement for the reference model derived from the Herrin et al. traveltimes tables, used in this work. Eventually, this sort of agreement to respect the *observed* data (errors not larger than a particular threshold) has been imposed as a first objective of this study, but it is not unique. The use of an analytical function avoids the common strategy of deriving spherical averages from seismological observations via an inversion procedure (i.e., the least-square approach). An interesting comment of this performing can be found in Morelli and Dziewonski (1993). In our scheme, the inherent biased data distribution is largely overcome since only traveltimes tables are taking into account. This absence of real data is a major lack in the model we present in this paper, and we agree. However, we keep the opinion that the results should be interpreted in a different way as those derived from a reference model, because they maintain internal consistency and do not pretend to be an alternative to PREM, ISAP91 or SP6 models.

The use of analytical functions to derive a model that globally reproduces the observed traveltimes by acting locally on a multilayer and spherical mantle does not prescribe the meaning, from a physical viewpoint, of the new model. Indeed, the analytical tomography results in an improved understanding of some particular areas, for example the D'' layer at the base of the mantle. These features are the most relevant conclusions of our work as they provide some slight differences to the current knowledge of the mantle.

## Methodology

The trial P-wave velocity function used in this work to analyse the structure of the mantle can be summarised by the expression

$$v(r) = r \cdot (B - A \cdot \ln(r)) \quad , \quad (1)$$

where  $r$  is the radius, and  $(A,B)$  two independent parameters to be determined. This formula can be simplified by defining the function

$$w(r) = (B - A \ln(r)) \quad , \quad (2)$$

and then:

$$v(r) = r \cdot w(r) \quad . \quad (3)$$

The P-wave velocity function expressed in Eq. (1) has been used (Lana-Renault and Cid, 1991; Lana-Renault, 1998) to obtain different Earth models by varying the different parameters. The smoothness of this function makes it adequate to tomographic studies of the Earth's mantle, once a proper parameterisation is applied, i.e. a division in many spherical layers, which is the one followed in this work. Another useful property of the function described in Eq. (1) is that converts the elliptical integrals arisen during the hamiltonian formulation of ray propagation, into analytical functions. For example, for a ray crossing the first layer ( $i = 1$ ) of the mantle, who radius of the top surface is  $R_1$  (Earth's radius), the epicentral distance  $D$  can be expressed as

$$D = \frac{2 \cdot w_1}{A_1} \cdot \sinh\left(\frac{A_1 \cdot T}{2}\right) \quad , \quad (4)$$

where:

$$w_1 = w_1(R_1) = B_1 - A_1 \cdot \ln(R_1) \quad .$$

The general analytical expressions for  $D$  and  $T$  can be obtained using the classical integral expressions (Bullen and Bolt, 1985)

$$D = p \cdot \int_{r_p}^{r_o} r^{-1} \cdot \sqrt{h^2 - p^2} \, dr \quad , \quad (5)$$


$$T = \int_{r_p}^{r_o} h^2 \cdot r^{-1} \cdot \sqrt{h^2 - p^2} \, dr \quad .$$

Denoting the angle of incidence at the top surface of the  $i^{th}$  layer by  $I_i$  and its radius by  $R_i$ , and similarly for the variables at the bottom ( $I_i'$  and  $R_i'$ ), see figure 1, it is always possible to write

$$w_i = w_i(R_i) = B_i - A_i \ln(R_i) = \frac{\sin(I_i)}{p} \quad , \quad (6)$$

and

$$w'_i = w'_i(R'_i) = B_i - A_i \ln(R'_i) = \frac{\sin(I'_i)}{p} \quad . \quad (7)$$



**Fig. 1.** P-trajectory traveling through a layer  $i$

In general, for a point  $P(r)$  we have

$$w_i(r) = B_i - A_i \ln(r) = \frac{\sin(I)}{p} \quad . \quad (8)$$

Hence, through its derivative,

$$\frac{dr}{r \cos(I)} = -\frac{dI}{p \cdot A_i} \quad (9)$$

we can calculate the expressions (5) for a P-trajectory which travels from  $R_i$  to  $R'_i$

$$D_i = \frac{\cos(I_i) - \cos(I'_i)}{p \cdot A_i} \quad , \quad (10)$$

$$T_i = \frac{1}{A_i} \cdot \ln \left[ \frac{\operatorname{tg}\left(\frac{I'_i}{2}\right)}{\operatorname{tg}\left(\frac{I_i}{2}\right)} \right] \quad . \quad (11)$$

Therefore, the observables at the Earth surface can be computed as a result of several additions of these computed values at each layer. That is, if one ray travels along  $k$  layers, the final epicentral distance and traveltime are calculated through of the following  $2k+1$  equations:

$$D = \frac{2}{p} \cdot \left( \sum_i \frac{\cos(I_i) - \cos(I'_i)}{A_i} + \frac{\cos(I_k)}{A_k} \right) , \quad (12)$$

$$T = 2 \cdot \left[ \sum_i \frac{1}{A_i} \cdot \ln \left[ \frac{\operatorname{tg}\left(\frac{I'_i}{2}\right)}{\operatorname{tg}\left(\frac{I_i}{2}\right)} \right] - \frac{1}{A_k} \cdot \ln \left( \operatorname{tg}\left(\frac{I_k}{2}\right) \right) \right] \quad (13)$$

and these  $2k-1$  auxiliary equations

$$p = \frac{\operatorname{sen}(I_i)}{w_i(R_i)} = \frac{\operatorname{sen}(I'_i)}{w_i(R'_i)} = \frac{\operatorname{sen}(I_k)}{w_k(R_k)} , \quad (14)$$

where:  $i = 1, 2, \dots, k-1$ .

On the other hand, the observables for a PcP-trajectory are calculated by the following  $2(k+1)$  equations:

$$D = \frac{2}{p} \cdot \left( \sum_i \frac{\cos(I_i) - \cos(I'_i)}{A_i} \right) , \quad (15)$$

$$T = 2 \cdot \left[ \sum_i \frac{1}{A_i} \cdot \ln \left[ \frac{\operatorname{tg}\left(\frac{I'_i}{2}\right)}{\operatorname{tg}\left(\frac{I_i}{2}\right)} \right] \right] \quad (16)$$

and these  $2k$  auxiliary equations

$$p = \frac{\operatorname{sen}(I_i)}{w_i(R_i)} = \frac{\operatorname{sen}(I'_i)}{w_i(R'_i)} . \quad (17)$$

where:  $i = 1, 2, \dots, k$ .

Finally, by integrating Eq. (9) between  $P(R_i)$  and  $P(r)$ , it is easy to calculate the radius of any single point  $P(r)$  along the trajectory:

$$r = R_i \cdot \exp \left( \frac{\operatorname{sen}(I_i) - \operatorname{sen}(I)}{p \cdot A_i} \right) = R_i \cdot \exp \left( \frac{w_i - w_i(r)}{A_i} \right) . \quad (18)$$

## Results

With a single collection of observed traveltimes, it is possible to reproduce the observations on the Earth's surface for any event. For the sake of simplicity, as an example of the versatility and functionality of the proposed methodology, we have selected the datasets reproduced in *Herrin et al.* (1968). The sequence of calculations consists of determining the specific constants  $A_i$ ,  $w_i$  and  $w'_i$  ( $i = 1, \dots, N$ ), for each layer,  $N$  being the number of layers.

(Note that  $w'_i$  is a measure of the thickness of the  $i^{th}$  layer and that we don't use  $B_i$ . The parameter  $B_i$  is calculated after using the Eq. (6))

Let suppose these quantities are already known for the first  $k-1$  layers, except  $w'_{k-1}$ , the starting point for the  $k^{th}$  layer. The inverse problem can be posed as a system of non-linear equations (12-14) that will provide the parameters  $A_k$  and  $w_k$  of the layer  $k$ . We must use three P-observed trajectories reproducing three fixed points  $(D_l, T_{ol}; l = 1, 2, 3)$  as boundary conditions for the system of  $3(2k+1)$  non linear equations with  $3(2k+1)$  unknowns ( $p_l, I_{ll}, I'_{ll}, I_{kl}, w'_{k-1}, w_k, A_k$ ). The solution is then iterated till assure a convergence criterion, in our case, a threshold for the computed residuals less than a certain value ( $10^{-15}$ ).

Known the values  $w'_{k-1}, w_k, y A_k$ , we prove that the residual times  $T_o - T_c$  (observed minus computed time) of the all the others P-trajectories which also return to the surface-focus from the  $k^{th}$  layer are smaller than a determined  $\epsilon$ . If it is not so, we begin again taking others three observables  $(D_l, T_{ol})$  nearer among them.


It is to be noted that the algebra applied in our methodology permits a discontinuity of the 1<sup>st</sup> kind ( $w'_{k-1} \neq w_k$ ) in the velocity function.

This last property can be analysed through the study of the derivative ( $dT/dD$ ) (*Herrin et al.*, 1968), in order to detect jumps in the selected velocity pattern. If we have the security that only a discontinuity of the 2<sup>nd</sup> kind ( $w'_{k-1} = w_k$ ) is present, then it is possible to work with only 2 observables  $(D, T_o)$  or fixed boundary conditions, thus eliminating  $2k+1$  redundant equations from the global system.

In this case, the experience tells us that is much better to work with one observable  $(D, T_o)$  and, thus, fixing the final of the  $k-1^{th}$  layer by a value for  $w'_{k-1}$ . and insuring that the residual times of all the P-trajectories which return from the  $k-1^{th}$  layer to the surface-focus are less than our  $\epsilon$ . Thereby, we resolve a non linear system with only  $2k+1$  equations.

We have performed a complete description of the Mantle using a maximum residual time  $\epsilon = 0.015$  s. and only 34 P-observed traveltimes. The total number of layers used in this description is 28. The last one finishes at a depth of 2810.1 km., maximum for the last P-observed trajectory at  $D = 100^\circ$  according to *Herrin et al.* (1968), with  $T_o = 826.7303$  s. Once known the problematic of the lack of information for  $D > 100^\circ$  and the special case of the D" shell, we have worked with

data available from  $D > 88^\circ$  and maximum residuals of 0.002 s. See Table 1 and Residuals of P-travel times (Figure 2) for a graphic representation and further details.




**Fig. 2.** P-residual times  $T_o - T_c$ .  $D$ :  $0 - 100^\circ$  every  $0.5^\circ$ . Maximum residual 0.015 sec. at  $D = 18.5^\circ$ .


**The 29<sup>th</sup> layer.** Since the derivative associated with the surface-focus travel time ( $dT/dD$ ) is effectively constant (Herrin et al., 1968) beyond  $99.0^\circ$ , and that our residuals are practically zero for those points, we consider the boundary condition  $w'_{28} = 1/p(100^\circ) = w_{29}$  produces the best results. Thus, our problem is reduced to calculate the parameters  $w'_{29}$  (final layer) and  $A_{29}$ . For this purpose, we pose a non linear system with two PcP observables. One of these fixed points should always be the axial trajectory ( $D = 0^\circ$ ;  $T_o = 511.3$  sec.) that allows us to use only one equation:

$$T = 2 \cdot \sum_i \frac{1}{A_i} \cdot \ln \left( \frac{w'_i}{w_i} \right) \quad (19)$$

We have considered that the other observable should be very separated from the first, and thus, selected  $D = 93^\circ$ ;  $T_o = 795.2$  s as second observable. Once obtained the values  $w'_{29}$  and  $A_{29}$ , all the PcP residuals were balanced with an error less than 0.13 s. (Figure 3, Table 2 of PcP-travel times). By applying Eq. (18) we found the outer core at 2893.9 km. Further details can be seen in Figures 4, 5, 6 and Table 3 of P-wave velocity.




**Fig. 3.** PcP-residual times  $T_o - T_c$ .  $\mathbf{D}$ :  $0 - 96^\circ$  every  $1^\circ$ . Maximum residual 0.13 sec. at  $\mathbf{D} = 75^\circ$ . Depth Outer Core: 2893.9 km.



**Fig. 4.** P-wave velocity in the mantle. *Herrin et al.* versus MZY.


Fig. 4 shows a comparison between our velocity distribution and the one provided by *Herrin et al.* (1968). The most conspicuous difference is observed at 2749.8 km., final of the 25<sup>th</sup> layer, where the last trajectory returns to the surface at  $\mathbf{D} = 92^\circ$  and the corresponding residual is null. Figure 5 exhibits residuals of computed velocity *Herrin et al.* minus MZY. Let us note how the maximum residual is 0.0044 km/s at a depth of 755 km.





**Fig. 5.** Residual velocity *Herrin et al.* minus *MZY* in the mantle until a depth of 2745 km. Maximum residual: 0.0044 km/sec at a depth of 755 km.

From 2749.8 km. until the core-mantle boundary (*region D''*), our velocity distribution begins to be completely different to *Herrin et al.* (1968), as can be seen in Fig. 6. This is due that *Herrin et al.* adopted a special smooth velocity distribution to the region *D''* to explain the last results from *Taggart and Engdahl*, (1968), which indicated a slow increase of velocity towards the core. Morelli and Dziewonski (1993), in their SP6 model, obtained a continuous decrease from 2741 km. In our *MZY* model, we propose that *D''* region begins at 2780.7 km. with a brief (29.4 km.) negative gradient (layers number 27 and 28) followed by a slow increase until the core. With this profile we insure the residuals of all the observables P and PcP from *Herrin et al.* (1968) are minima, and reproduce accurately the observed times.



**Fig. 6.** P-wave velocity in *D''* region. *Herrin et al.* versus *MZY*

## **Concluding remarks**

We have presented a new technique to tomography the interior of the Earth for which one is able to obtain residual times less than a determined value  $\epsilon$  for all observed trajectories  $P$ . The more minor is the value of  $\epsilon$ , more genuine and real is the tomography.

Also, we have seen that the technique MZY developed is very easy to apply. Its potentiality is based in the function velocity found that it provides us analytical solutions for  $\Delta$  and  $T$ .

## **Acknowledgements**

The author is grateful to Dr. Javier Sabadell by the comments and suggestions made during the writing of the manuscript.

Table 1						P travel times (sec.)					p. 1/2
La- yers	$\Delta_{\text{initial}}$ $\Delta_{\text{final}}$	$\Delta$	Observed (Herrin & al. )	Computed MZY	Residual $T_o - T_c$	La- yers	$\Delta_{\text{initial}}$ $\Delta_{\text{final}}$	$\Delta$	Observed (Herrin & al. )	Computed MZY	Residual $T_o - T_c$
1	0	0.00	0.0000	0.0000	0.0000	15	27.14	27.50	347.2025	347.1930	0.0095
		0.50	9.2663	9.2662	0.0001			28.00	351.6796	351.6743	0.0053
	7.85	1.00	18.5323	18.5321	0.0002			28.50	356.1456	356.1490	-0.0034
2	0.52	1.50	26.9525	26.9525	0.0000			29.00	360.6048	360.6144	-0.0096
	10.66							29.50	365.0596	365.0682	-0.0086
3	0.989	2.00	34.8630	34.8630	0.0000	16	30	30.00	369.5086	369.5086	0.0000
		2.50	41.7231	41.7213	0.0018		30	30.50	373.9477	373.9401	0.0076
		3.00	48.5813	48.5782	0.0031			31.00	378.3751	378.3650	0.0101
		3.50	55.4373	55.4334	0.0039			31.50	382.7900	382.7810	0.0090
		4.00	62.2906	62.2864	0.0042			32.00	387.1923	387.1867	0.0056
		4.50	69.1410	69.1367	0.0043			32.50	391.5831	391.5808	0.0023
		5.00	75.9880	75.9840	0.0040			33.00	395.9621	395.9627	-0.0006
		5.50	82.8312	82.8278	0.0034			33.50	400.3281	400.3315	-0.0034
		6.00	89.6703	89.6676	0.0027			34.00	404.6807	404.6864	-0.0057
		6.50	96.5049	96.5030	0.0019			34.50	409.0193	409.0270	-0.0077
		7.00	103.3346	103.3337	0.0009			35.00	413.3435	413.3527	-0.0092
		7.50	110.1591	110.1591	0.0000			35.50	417.6532	417.6629	-0.0097
		8.00	116.9779	116.9789	-0.0010			36.00	421.9479	421.9573	-0.0094
		8.50	123.7908	123.7926	-0.0018			36.50	426.2269	426.2353	-0.0084
		9.00	130.5973	130.5998	-0.0025			37.00	430.4894	430.4967	-0.0073
		9.50	137.3970	137.4000	-0.0030			37.50	434.7347	434.7411	-0.0064
		10.00	144.1896	144.1930	-0.0034			38.00	438.9626	438.9681	-0.0055
		10.50	150.9747	150.9783	-0.0036			38.50	443.1730	443.1775	-0.0045
		11.00	157.7519	157.7554	-0.0035			39.00	447.3662	447.3690	-0.0028
		11.50	164.5209	164.5240	-0.0031	17	39.5	39.50	451.5425	451.5425	0.0000
		12.00	171.2813	171.2836	-0.0023		39.5	40.00	455.7020	455.7008	0.0012
		12.50	178.0326	178.0340	-0.0014			40.50	459.8449	459.8440	0.0009
	13.15	13.00	184.7746	184.7746	0.0000			41.00	463.9710	463.9709	0.0001
4	13.15	13.50	191.4964	191.4929	0.0035			41.50	468.0802	468.0808	-0.0006
	14.14	14.14	200.0582 *	200.0655	-0.0073			42.00	472.1723	472.1736	-0.0013
5	14.14	14.50	204.8555	204.8480	0.0075			42.50	476.2473	476.2488	-0.0015
	15.13	15.13	213.1831 *	213.1946	-0.0115			43.00	480.3051	480.3064	-0.0013
6	15.13	15.50	218.0429	218.0320	0.0109			43.50	484.3454	484.3462	-0.0008
	16.11	16.11	225.9644 *	225.9771	-0.0127			44.00	488.3680	488.3680	0.0000
7	16.11	16.50	230.9845	230.9726	0.0119			44.50	492.3728	492.3716	0.0012
	17.09	17.09	238.4697 *	238.4824	-0.0127			45.00	496.3596	496.3571	0.0025
8	17.09	17.50	243.6096	243.5956	0.0140			45.50	500.3285	500.3244	0.0041
	18.08	18.08	250.7491 *	250.7627	-0.0136			46.00	504.2791	504.2733	0.0058
9	18.08	18.50	255.8408	255.8261	0.0147			46.50	508.2111	508.2038	0.0073
	19.06	19.06	262.4864 *	262.4973	-0.0109			47.00	512.1242	512.1159	0.0083
10	19.06	19.50	267.6136	267.6016	0.0120			47.50	516.0178	516.0096	0.0082
	20.04	20.04	273.7653 *	273.7721	-0.0068			48.00	519.8920	519.8847	0.0073
11	20.04	20.50	278.9036	278.8956	0.0080			48.50	523.7469	523.7415	0.0054
	21.02	21.02	284.5832 *	284.5863	-0.0031			49.00	527.5828	527.5797	0.0031
12	21.02	21.50	289.7160	289.7114	0.0046			49.50	531.4001	531.3995	0.0006
	22	22.00	294.9501	294.9501	0.0000			50.00	535.1992	535.2008	-0.0016
13	22	22.50	300.0806	300.0759	0.0047			50.50	538.9802	538.9837	-0.0035
		23.00	305.1134	305.1050	0.0084			51.00	542.7433	542.7482	-0.0049
		23.50	310.0533	310.0470	0.0063			51.50	546.4887	546.4944	-0.0057
		24.00	314.9070	314.9070	0.0000			52.00	550.2164	550.2222	-0.0058
		24.50	319.6818	319.6890	-0.0072			52.50	553.9266	553.9317	-0.0051
14	25.46	25.00	324.3869	324.3961	-0.0092			53.00	557.6192	557.6230	-0.0038
	25.46	25.46	328.6614 *	328.6630	-0.0016			53.50	561.2941	561.2962	-0.0021
		26.00	333.6295	333.6262	0.0033			54.00	564.9510	564.9512	-0.0002
		26.50	338.1848	338.1873	-0.0025			54.50	568.5899	568.5882	0.0017
	27.14	27.14	343.9656 *	343.9643	0.0013			55.00	572.2107	572.2072	0.0035

Table 1						P travel times (sec.)						p. 2/2
La- yers	$\Delta_{\text{initial}}$ $\Delta_{\text{final}}$	$\Delta$	Observed (Herrin & al. )	Computed MZY	Residual $T_o - T_c$	La- yers	$\Delta_{\text{initial}}$ $\Delta_{\text{final}}$	$\Delta$	Observed (Herrin & al. )	Computed MZY	Residual $T_o - T_c$	
17		55.50	575.8137	575.8082	0.0055	22	81.8	81.80	740.2336 *	740.2409	-0.0073	
		56.00	579.3986	579.3915	0.0071			82.50	743.9007	743.8974	0.0033	
		56.50	582.9653	582.9569	0.0084			83.00	746.4926	746.4852	0.0074	
		57.00	586.5135	586.5047	0.0088			83.50	749.0611	749.0544	0.0067	
		57.50	590.0430	590.0348	0.0082		84.04	84.04	751.8075 *	751.8092	-0.0017	
		58.00	593.5538	593.5475	0.0063		84.04	84.50	754.1271	754.1271	0.0000	
		58.50	597.0462	597.0427	0.0035			85.00	756.6260	756.6263	-0.0003	
59	59.00	600.5205	600.5205	0.0000			85.50	759.1042	759.1061	-0.0019		
59	59.50	603.9770	603.9803	-0.0033	23		86.00	761.5636	761.5676	-0.0040		
	60.00	607.4162	607.4225	-0.0063			86.50	764.0064	764.0117	-0.0053		
	60.50	610.8385	610.8471	-0.0086			87.00	766.4338	766.4390	-0.0052		
	61.00	614.2444	614.2545	-0.0101			87.50	768.8465	768.8502	-0.0037		
	61.50	617.6343	617.6447	-0.0104		88	88.00	771.2455	771.2455	0.0000		
	62.00	621.0084	621.0179	-0.0095		88	88.50	773.6315	773.6301	0.0014		
	62.50	624.3668	624.3743	-0.0075		24		89.00	776.0056	776.0053	0.0003	
	63.00	627.7094	627.7138	-0.0044			89.50	778.3687	778.3695	-0.0008		
	63.50	631.0356	631.0367	-0.0011	90		90.00	780.7222	780.7222	0.0000		
18		64.00	634.3452	634.3430	0.0022	25	90	90.50	783.0673	783.0673	0.0000	
		64.50	637.6379	637.6329	0.0050			91.00	785.4049	785.4061	-0.0012	
		65.00	640.9137	640.9064	0.0073			91.50	787.7356	787.7371	-0.0015	
		65.50	644.1724	644.1637	0.0087	26	92	92.00	790.0597	790.0597	0.0000	
		66.00	647.4142	647.4049	0.0093		92	92.50	792.3774	792.3767	0.0007	
		66.50	650.6392	650.6301	0.0091			93.00	794.6891	794.6892	-0.0001	
		67.00	653.8477	653.8395	0.0082		93.50	796.9953	796.9961	-0.0008		
		67.50	657.0398	657.0330	0.0068	94.03	94.03	799.4344 *	799.4344	0.0000		
		68.00	660.2151	660.2108	0.0043	27	94.03	94.50	801.5937	801.5930	0.0007	
		68.50	663.3731	663.3731	0.0000			95.00	803.8872	803.8873	-0.0001	
		69.08	667.0129 *	667.0220	-0.0091			95.50	806.1777	806.1784	-0.0007	
	69.08	69.50	669.6355	669.6387	-0.0032	96.123	96.123	809.0282 *	809.0279	0.0003		
	19	70.00	672.7383	672.7354	0.0029	28	96.123	96.50	810.7518	810.7503	0.0015	
		70.50	675.8202	675.8127	0.0075			97.00	813.0361	813.0345	0.0016	
		71.00	678.8805	678.8716	0.0089			97.50	815.3192	815.3185	0.0007	
		71.50	681.9193	681.9128	0.0065			98.00	817.6016	817.6021	-0.0005	
72.35		72.00	684.9366	684.9366	0.0000			98.50	819.8838	819.8851	-0.0013	
72.35	72.35	687.0340 *	687.0432	-0.0092			99.00	822.1660	822.1676	-0.0016		
20		73.00	690.9092	690.9170	-0.0078		99.50	824.4481	824.4494	-0.0013		
		73.50	693.8665	693.8729	-0.0064	100	100.00	826.7303	826.7303	0.0000		
		74.00	696.8054	696.8096	-0.0042	29	100	99.5	824.4481	824.4498	-0.0017	
		74.50	699.7264	699.7279	-0.0015			99.0	822.1660	822.1694	-0.0034	
		75.00	702.6299	702.6283	0.0016			98.5	819.8838	819.8892	-0.0054	
		75.50	705.5159	705.5115	0.0044			98.0	817.6016	817.6094	-0.0078	
		76.00	708.3843	708.3776	0.0067			97.5	815.3192	815.3302	-0.0110	
		76.50	711.2346	711.2272	0.0074			97.0	813.0361	813.0520	-0.0159	
		77.00	714.0661	714.0604	0.0057		96.5	810.7518	810.7757	-0.0239		
	77.9	77.50	716.8776	716.8776	0.0000		96.0	808.4658	808.5028	-0.0370		
77.9	77.90	719.1107 *	719.1200	-0.0093		95.618	806.7177 *	806.7715	-0.0538			
21		78.50	722.4405	722.4426	-0.0021		96.0	808.4658	808.4805	-0.0147		
		79.00	725.1920	725.1893	0.0027	96.289	96.289	809.7871 *	809.7642	0.0229		
		79.50	727.9234	727.9171	0.0063	interpolated *						
		80.00	730.6349	730.6267	0.0082							
		80.50	733.3270	733.3190	0.0080							
		81.00	735.9998	735.9944	0.0054							
	81.8	81.50	738.6533	738.6533	0.0000							

<b>Table 2</b> <b>PcP travel times (sec.)</b>							
$\Delta$	Observed (Herrin & al.)	Computed MZY	Residual $T_o - T_c$	$\Delta$	Observed (Herrin & al.)	Computed MZY	Residual $T_o - T_c$
0	511.3	511.300	0.00	49	611.9	611.870	0.03
1	511.4	511.348	0.05	50	615.5	615.511	-0.01
2	511.5	511.492	0.01	51	619.2	619.191	0.01
3	511.7	511.731	-0.03	52	622.9	622.910	-0.01
4	512.1	512.066	0.03	53	626.7	626.666	0.03
5	512.5	512.497	0.00	54	630.4	630.458	-0.06
6	513.0	513.022	-0.02	55	634.3	634.284	0.02
7	513.6	513.641	-0.04	56	638.1	638.144	-0.04
8	514.4	514.355	0.04	57	642.0	642.037	-0.04
9	515.2	515.163	0.04	58	646.0	645.961	0.04
10	516.1	516.063	0.04	59	649.9	649.914	-0.01
11	517.1	517.055	0.04	60	653.9	653.896	0.00
12	518.1	518.139	-0.04	61	657.9	657.906	-0.01
13	519.3	519.314	-0.01	62	661.9	661.943	-0.04
14	520.6	520.578	0.02	63	666.0	666.005	0.00
15	521.9	521.932	-0.03	64	670.1	670.091	0.01
16	523.4	523.373	0.03	65	674.2	674.201	0.00
17	524.9	524.901	0.00	66	678.3	678.333	-0.03
18	526.5	526.515	-0.01	67	682.5	682.486	0.01
19	528.2	528.214	-0.01	68	686.6	686.660	-0.06
20	530.0	529.996	0.00	69	690.8	690.853	-0.05
21	531.9	531.861	0.04	70	695.1	695.065	0.04
22	533.8	533.806	-0.01	71	699.3	699.294	0.01
23	535.8	535.832	-0.03	72	703.5	703.540	-0.04
24	537.9	537.936	-0.04	73	707.9	707.802	0.10
25	540.1	540.117	-0.02	74	712.1	712.079	0.02
26	542.4	542.374	0.03	75	716.5	716.370	0.13
27	544.7	544.705	-0.01	76	720.7	720.675	0.03
28	547.1	547.110	-0.01	77	725.0	724.992	0.01
29	549.6	549.586	0.01	78	729.3	729.320	-0.02
30	552.1	552.132	-0.03	79	733.6	733.660	-0.06
31	554.7	554.747	-0.05	80	738.0	738.010	-0.01
32	557.4	557.428	-0.03	81	742.4	742.370	0.03
33	560.2	560.176	0.02	82	746.7	746.739	-0.04
34	563.0	562.988	0.01	83	751.1	751.116	-0.02
35	565.9	565.863	0.04	84	755.5	755.500	0.00
36	568.8	568.798	0.00	85	759.9	759.892	0.01
37	571.8	571.794	0.01	86	764.3	764.290	0.01
38	574.8	574.848	-0.05	87	768.7	768.693	0.01
39	578.0	577.959	0.04	88	773.1	773.102	0.00
40	581.1	581.125	-0.02	89	777.5	777.515	-0.01
41	584.3	584.345	-0.04	90	781.9	781.932	-0.03
42	587.6	587.617	-0.02	91	786.3	786.352	-0.05
43	590.9	590.940	-0.04	92	790.8	790.775	0.03
44	594.3	594.313	-0.01	93	795.2	795.200	0.00
45	597.7	597.734	-0.03	94	799.6	799.627	-0.03
46	601.2	601.202	0.00	95	804.0	804.055	-0.05
47	604.7	604.714	-0.01	96	808.5	808.484	0.02
48	608.3	608.271	0.03	96.289		809.764	

Table 3

Data	MZY		P-wave velocity (km/s)			Data	MZY	p. 1/4	P-wave velocity (km/s)		
Radius (km)	$v_i$ (km/s)	Depth (km)	Radius of surface-focus = 6371.028			Radius (km)	$v_i$ (km/s)	Depth (km)			
Layers	$B_i$ ( $\times 10^{-2}$ )		Herrin & al.	MZY	Residual H. & al. - MZY	Layers	$B_i$ ( $\times 10^{-2}$ )		Herrin & al.	MZY	Residual H. & al. - MZY
Depth (km)	$v'_i$ (km/s)					Depth (km)	$A_i$ ( $\times 10^{-3}$ )				
6371.028	6.0000	0	6.0000	6.0001	-0.0001	6021.290062	8.8862	350	8.8905	8.8875	0.0030
1	0.92341045	5	6.0000	6.0001	-0.0001			355	8.9131	8.9114	0.0017
	0.94666548	10	6.0000	6.0001	-0.0001			360	8.9360	8.9352	0.0008
15.001533	6.0001	15	6.0000	6.0001	-0.0001			365	8.9590	8.9591	-0.0001
6356.026467	6.7500	20	6.7500	6.7501	-0.0001			370	8.9823	8.9829	-0.0006
2	1.04491768	25	6.7500	6.7501	-0.0001	10	5.59329351	375	9.0058	9.0067	-0.0009
	1.07194400	30	6.7500	6.7502	-0.0002		6.25724227	380	9.0294	9.0304	-0.0010
		35	6.7500	6.7502	-0.0002			385	9.0532	9.0542	-0.0010
40.053935	6.7502	40	6.7500	6.7502	-0.0002			390	9.0773	9.0779	-0.0006
6330.974065	8.0540	45	8.0582	8.0593	-0.0011			395	9.1015	9.1016	-0.0001
		50	8.0642	8.0645	-0.0003			400	9.1258	9.1252	0.0006
		55	8.0698	8.0698	0.0000			405	9.1503	9.1489	0.0014
		60	8.0753	8.0751	0.0002	411.322422	9.1787	410	9.1750	9.1725	0.0025
		65	8.0806	8.0803	0.0003	5959.705578	9.1787	415	9.1999	9.1977	0.0022
3	2.16670136	70	8.0859	8.0856	0.0003			420	9.2248	9.2236	0.0012
	2.32998530	75	8.0911	8.0908	0.0003			425	9.2499	9.2494	0.0005
		80	8.0962	8.0960	0.0002			430	9.2752	9.2752	0.0000
		85	8.1013	8.1012	0.0001			435	9.3007	9.3010	-0.0003
		90	8.1064	8.1064	0.0000	11	5.99325871	440	9.3262	9.3267	-0.0005
		95	8.1115	8.1116	-0.0001		6.71735442	445	9.3519	9.3524	-0.0005
104.957687	8.1219	100	8.1165	8.1168	-0.0003			450	9.3778	9.3781	-0.0003
6266.070313	8.1219	105	8.1219	8.1220	-0.0001			455	9.4038	9.4038	0.0000
4	2.51841026	110	8.1285	8.1292	-0.0007			460	9.4299	9.4294	0.0005
	2.73226453	115	8.1356	8.1363	-0.0007			465	9.4562	9.4550	0.0012
		120	8.1432	8.1435	-0.0003	472.071820	9.4911	470	9.4826	9.4805	0.0021
125.320	8.1511	125	8.1513	8.1506	0.0007	5898.956180	9.4911	475	9.5091	9.5072	0.0019
6245.707939	8.1511	130	8.1599	8.1602	-0.0003			480	9.5358	9.5345	0.0013
		135	8.1690	8.1699	-0.0009			485	9.5626	9.5618	0.0008
5	2.97405843	140	8.1786	8.1796	-0.0010			490	9.5895	9.5891	0.0004
	3.25362200	145	8.1886	8.1893	-0.0007			495	9.6165	9.6164	0.0001
		150	8.1991	8.1990	0.0001	12	6.31396167	500	9.6437	9.6436	0.0001
155.027311	8.2087	155	8.2101	8.2087	0.0014		7.08672019	505	9.6709	9.6709	0.0000
6216.000689	8.2087	160	8.2214	8.2213	0.0001			510	9.6981	9.6980	0.0001
		165	8.2332	8.2340	-0.0008			515	9.7255	9.7252	0.0003
6	3.50244082	170	8.2454	8.2467	-0.0013			520	9.7530	9.7523	0.0007
	3.85853274	175	8.2580	8.2593	-0.0013			525	9.7805	9.7794	0.0011
		180	8.2710	8.2719	-0.0009	531.559687	9.8149	530	9.8080	9.8064	0.0016
		185	8.2843	8.2845	-0.0002	5839.468313	9.8149	535	9.8356	9.8342	0.0014
193.795314	8.3067	190	8.2980	8.2971	0.0009			540	9.8632	9.8623	0.0009
6177.232686	8.3067	195	8.3120	8.3104	0.0016			545	9.8908	9.8903	0.0005
		200	8.3264	8.3260	0.0004			550	9.9185	9.9184	0.0001
		205	8.3410	8.3415	-0.0005			555	9.9462	9.9464	-0.0002
		210	8.3560	8.3571	-0.0011			560	9.9740	9.9743	-0.0003
		215	8.3713	8.3726	-0.0013			565	10.0018	10.0023	-0.0005
		220	8.3870	8.3881	-0.0011			570	10.0296	10.0302	-0.0006
		225	8.4029	8.4036	-0.0007			575	10.0574	10.0580	-0.0006
		230	8.4191	8.4191	0.0000			580	10.0853	10.0859	-0.0006
239.227039	8.4476	235	8.4357	8.4345	0.0012			585	10.1132	10.1137	-0.0005
6131.800961	8.4476	240	8.4525	8.4504	0.0021			590	10.1411	10.1414	-0.0003
		245	8.4696	8.4689	0.0007			595	10.1690	10.1692	-0.0002
		250	8.4870	8.4873	-0.0003			600	10.1970	10.1969	0.0001
		255	8.5047	8.5057	-0.0010			605	10.2249	10.2246	0.0003
		260	8.5227	8.5241	-0.0014	13	6.50401417	610	10.2528	10.2522	0.0006
		265	8.5410	8.5424	-0.0014		7.30586668	615	10.2807	10.2798	0.0009
		270	8.5595	8.5607	-0.0012			620	10.3086	10.3074	0.0012
		275	8.5783	8.5791	-0.0008			625	10.3364	10.3350	0.0014
		280	8.5973	8.5974	-0.0001			630	10.3642	10.3625	0.0017
		285	8.6167	8.6156	0.0011			635	10.3920	10.3900	0.0020
291.392218	8.6390	290	8.6362	8.6339	0.0023			640	10.4197	10.4174	0.0023
6079.635782	8.6390	295	8.6561	8.6543	0.0018			645	10.4474	10.4449	0.0025
		300	8.6762	8.6756	0.0006			650	10.4750	10.4723	0.0027
		305	8.6966	8.6969	-0.0003			655	10.5024	10.4996	0.0028
		310	8.7172	8.7182	-0.0010			660	10.5297	10.5270	0.0027
		315	8.7380	8.7394	-0.0014			665	10.5570	10.5542	0.0028
		320	8.7591	8.7606	-0.0015			670	10.5840	10.5815	0.0025
		325	8.7804	8.7818	-0.0014			675	10.6109	10.6087	0.0022
		330	8.8020	8.8029	-0.0009			680	10.6375	10.6359	0.0016
		335	8.8238	8.8241	-0.0003			685	10.6638	10.6631	0.0007
		340	8.8458	8.8452	0.0006			690	10.6899	10.6903	-0.0004
349.737938	8.8862	345	8.8680	8.8663	0.0017			695	10.7157	10.7174	-0.0017
						700.177664	10.7454	700	10.7412	10.7444	-0.0032

Table 3

Data MZY		Depth (km)	P-wave velocity (km/s)			Data MZY		p. 2/4	P-wave velocity (km/s)		
Radius (km)	$V_i$ (km/s)		Radius of surface-focus = 6371.028			Radius (km)	$V_i$ (km/s)				
Layers	$B_i$ ( $\times 10^{-2}$ )		Herrin & al.	MZY	Residual	Layers	$B_i$ ( $\times 10^{-2}$ )		Herrin & al.	MZY	Residual
Depth (km)	$V'_i$ (km/s)		H. & al. - MZY			Depth (km)	$A_i$ ( $\times 10^{-3}$ )		H. & al. - MZY		
5670.850336	10.7454	705	10.7664	10.7679	-0.0015			1115	11.6288	11.6296	-0.0008
		710	10.7911	10.7912	-0.0001			1120	11.6367	11.6375	-0.0008
		715	10.8154	10.8144	0.0010			1125	11.6446	11.6455	-0.0009
14	5.85891401	720	10.8392	10.8376	0.0016			1130	11.6525	11.6534	-0.0009
	6.55949029	725	10.8624	10.8608	0.0016			1135	11.6604	11.6613	-0.0009
		730	10.8850	10.8840	0.0010			1140	11.6684	11.6692	-0.0008
		735	10.9068	10.9071	-0.0003			1145	11.6763	11.6771	-0.0008
744.109784	10.9492	740	10.9279	10.9302	-0.0023			1150	11.6842	11.6849	-0.0007
5626.918216	10.9492	745	10.9479	10.9515	-0.0036			1155	11.6921	11.6928	-0.0007
		750	10.9663	10.9645	0.0018			1160	11.7000	11.7006	-0.0006
15	4.12047826	755	10.9819	10.9775	0.0044			1165	11.7080	11.7084	-0.0004
	4.54632080	760	10.9933	10.9904	0.0029			1170	11.7159	11.7162	-0.0003
		765	11.0029	11.0033	-0.0004			1175	11.7238	11.7239	-0.0001
772.052042	11.0215	770	11.0134	11.0163	-0.0029			1180	11.7316	11.7317	-0.0001
5598.975958	11.0215	775	11.0240	11.0273	-0.0033			1185	11.7395	11.7394	0.0001
		780	11.0348	11.0370	-0.0022			1190	11.7473	11.7471	0.0002
		785	11.0455	11.0467	-0.0012			1195	11.7551	11.7548	0.0003
		790	11.0561	11.0564	-0.0003			1200	11.7629	11.7624	0.0005
		795	11.0666	11.0660	0.0006			1205	11.7707	11.7701	0.0006
		800	11.0766	11.0757	0.0009			1210	11.7785	11.7777	0.0008
		805	11.0865	11.0853	0.0012			1215	11.7862	11.7853	0.0009
		810	11.0962	11.0949	0.0013			1220	11.7938	11.7929	0.0009
		815	11.1060	11.1045	0.0015			1225	11.8015	11.8005	0.0010
		820	11.1156	11.1141	0.0015			1230	11.8091	11.8081	0.0010
		825	11.1252	11.1236	0.0016			1235	11.8167	11.8156	0.0011
		830	11.1348	11.1332	0.0016			1240	11.8242	11.8231	0.0011
		835	11.1443	11.1427	0.0016			1245	11.8318	11.8306	0.0012
		840	11.1538	11.1522	0.0016			1250	11.8393	11.8381	0.0012
		845	11.1632	11.1617	0.0015			1255	11.8467	11.8456	0.0011
		850	11.1726	11.1711	0.0015			1260	11.8542	11.8530	0.0012
16	3.57557893	855	11.1819	11.1806	0.0013			1265	11.8616	11.8605	0.0011
	3.91494425	860	11.1912	11.1900	0.0012			1270	11.8689	11.8679	0.0010
		865	11.2004	11.1994	0.0010			1275	11.8763	11.8753	0.0010
		870	11.2096	11.2088	0.0008			1280	11.8836	11.8826	0.0010
		875	11.2189	11.2182	0.0007			1285	11.8909	11.8900	0.0009
		880	11.2281	11.2276	0.0005			1290	11.8982	11.8973	0.0009
		885	11.2373	11.2369	0.0004			1295	11.9054	11.9046	0.0008
		890	11.2466	11.2462	0.0004	17	3.48245516	1300	11.9126	11.9119	0.0007
		895	11.2558	11.2555	0.0003		3.80663410	1305	11.9198	11.9192	0.0006
		900	11.2651	11.2648	0.0003			1310	11.9269	11.9265	0.0004
		905	11.2743	11.2741	0.0002			1315	11.9341	11.9337	0.0004
		910	11.2835	11.2833	0.0002			1320	11.9412	11.9409	0.0003
		915	11.2927	11.2926	0.0001			1325	11.9483	11.9481	0.0002
		920	11.3019	11.3018	0.0001			1330	11.9554	11.9553	0.0001
		925	11.3110	11.3110	0.0000			1335	11.9624	11.9625	-0.0001
		930	11.3201	11.3202	-0.0001			1340	11.9695	11.9696	-0.0001
		935	11.3291	11.3293	-0.0002			1345	11.9765	11.9768	-0.0003
		940	11.3381	11.3385	-0.0004			1350	11.9836	11.9839	-0.0003
		945	11.3470	11.3476	-0.0006			1355	11.9906	11.9910	-0.0004
950.864911	11.3583	950	11.3558	11.3567	-0.0009			1360	11.9976	11.9980	-0.0004
5420.163089	11.3583	955	11.3646	11.3654	-0.0008			1365	12.0046	12.0051	-0.0005
		960	11.3734	11.3739	-0.0005			1370	12.0116	12.0121	-0.0005
		965	11.3820	11.3824	-0.0004			1375	12.0185	12.0191	-0.0006
		970	11.3907	11.3909	-0.0002			1380	12.0255	12.0261	-0.0006
		975	11.3993	11.3994	-0.0001			1385	12.0324	12.0331	-0.0007
		980	11.4079	11.4079	0.0000			1390	12.0393	12.0401	-0.0008
		985	11.4164	11.4163	0.0001			1395	12.0462	12.0470	-0.0008
		990	11.4249	11.4247	0.0002			1400	12.0531	12.0539	-0.0008
		995	11.4333	11.4331	0.0002			1405	12.0599	12.0608	-0.0009
		1000	11.4418	11.4415	0.0003			1410	12.0668	12.0677	-0.0009
		1005	11.4502	11.4499	0.0003			1415	12.0736	12.0746	-0.0010
		1010	11.4585	11.4582	0.0003			1420	12.0805	12.0814	-0.0009
		1015	11.4668	11.4666	0.0002			1425	12.0873	12.0882	-0.0009
		1020	11.4751	11.4749	0.0002			1430	12.0941	12.0950	-0.0009
		1025	11.4834	11.4832	0.0002			1435	12.1009	12.1018	-0.0009
		1030	11.4917	11.4915	0.0002			1440	12.1077	12.1086	-0.0009
		1035	11.4999	11.4998	0.0001			1445	12.1145	12.1153	-0.0008
17	3.48245516	1040	11.5081	11.5080	0.0001			1450	12.1213	12.1221	-0.0008
	3.80663410	1045	11.5163	11.5162	0.0001			1455	12.1281	12.1288	-0.0007
		1050	11.5244	11.5245	-0.0001			1460	12.1349	12.1354	-0.0005
		1055	11.5326	11.5326	0.0000			1465	12.1417	12.1421	-0.0004
		1060	11.5407	11.5408	-0.0001			1470	12.1485	12.1488	-0.0003
		1065	11.5488	11.5490	-0.0002			1475	12.1553	12.1554	-0.0001
		1070	11.5569	11.5571	-0.0002			1480	12.1620	12.1620	0.0000
		1075	11.5649	11.5652	-0.0003			1485	12.1688	12.1686	0.0002
		1080	11.5730	11.5734	-0.0004			1490	12.1755	12.1752	0.0003
		1085	11.5810	11.5814	-0.0004			1495	12.1822	12.1817	0.0005
		1090	11.5890	11.5895	-0.0005			1500	12.1889	12.1882	0.0007
		1095	11.5970	11.5976	-0.0006			1505	12.1956	12.1948	0.0008
		1100	11.6049	11.6056	-0.0007			1510	12.2023	12.2013	0.0010
		1105	11.6129	11.6136	-0.0007	1516.957451	12.2103	1515	12.2089	12.2077	0.0012
		1110	11.6208	11.6216	-0.0008						

[illegible]



Data MZY		Depth (km)	P-wave velocity (km/s)			Data MZY		p. 4/4	P-wave velocity (km/s)		
Radius (km)	$V_f$ (km/s)		Radius of surface-focus = 6371.028			Radius (km)	$V_f$ (km/s)		Radius of surface-focus = 6371.028		
Layers	$B_f$ ( $\times 10^{-2}$ ) $A_f$ ( $\times 10^{-3}$ )		Herrin & al.	MZY	Residual	Layers	$B_f$ ( $\times 10^{-2}$ ) $A_f$ ( $\times 10^{-3}$ )		Herrin & al.	MZY	Residual
Depth (km)	$V'_f$ (km/s)			H. & al. - MZY	Depth (km)	$V'_f$ (km/s)			H. & al. - MZY		
4169.610716	13.0106	2205	13.0175	13.0151	0.0024	3887.018610	13.3469	2485	13.3499	13.3482	0.0017
		2210	13.0234	13.0214	0.0020			2490	13.3562	13.3549	0.0013
		2215	13.0292	13.0277	0.0015			2495	13.3626	13.3615	0.0011
		2220	13.0350	13.0339	0.0011			2500	13.3690	13.3680	0.0010
		2225	13.0408	13.0401	0.0007			2505	13.3753	13.3746	0.0007
		2230	13.0466	13.0462	0.0004			2510	13.3817	13.3811	0.0006
		2235	13.0525	13.0524	0.0001			2515	13.3881	13.3876	0.0005
		2240	13.0583	13.0585	-0.0002			2520	13.3945	13.3940	0.0005
		2245	13.0641	13.0646	-0.0005			2525	13.4009	13.4004	0.0005
		2250	13.0700	13.0707	-0.0007			2530	13.4073	13.4068	0.0005
		2255	13.0758	13.0767	-0.0009			2535	13.4137	13.4132	0.0005
		2260	13.0817	13.0827	-0.0010			2540	13.4200	13.4195	0.0005
		2265	13.0876	13.0887	-0.0011			2545	13.4264	13.4258	0.0006
		2270	13.0934	13.0947	-0.0013			2550	13.4327	13.4321	0.0006
		2275	13.0993	13.1006	-0.0013			2555	13.4391	13.4383	0.0008
		2280	13.1052	13.1065	-0.0013			2560	13.4454	13.4445	0.0009
		2285	13.1110	13.1124	-0.0014			2565	13.4516	13.4507	0.0009
		2290	13.1169	13.1182	-0.0013			2570	13.4578	13.4568	0.0010
		2295	13.1228	13.1241	-0.0013			2575	13.4640	13.4629	0.0011
		2300	13.1287	13.1298	-0.0011			2580	13.4702	13.4690	0.0012
		2305	13.1345	13.1356	-0.0011			2585	13.4763	13.4750	0.0013
2310	13.1404	13.1414	-0.0010	2590	13.4823	13.4810	0.0013				
2315	13.1462	13.1471	-0.0009	2595	13.4883	13.4870	0.0013				
2320	13.1521	13.1528	-0.0007	2600	13.4942	13.4930	0.0012				
2325	13.1579	13.1584	-0.0005	2605	13.5000	13.4989	0.0011				
2330	13.1637	13.1641	-0.0004	2610	13.5058	13.5048	0.0010				
2335	13.1696	13.1697	-0.0001	2615	13.5116	13.5106	0.0010				
2340	13.1754	13.1753	0.0001	2620	13.5172	13.5164	0.0008				
2345	13.1812	13.1808	0.0004	2625	13.5229	13.5222	0.0007				
2350	13.1871	13.1863	0.0008	2630	13.5285	13.5280	0.0005				
2355	13.1929	13.1918	0.0011	2635	13.5340	13.5337	0.0003				
2360	13.1987	13.1973	0.0014	2640	13.5394	13.5394	0.0000				
2365	13.2046	13.2028	0.0018	2645	13.5448	13.5451	-0.0003				
2370	13.2104	13.2082	0.0022	2650	13.5501	13.5507	-0.0006				
2372.501241	13.2109				2655	13.5554	13.5563	-0.0009			
3998.526759	13.2109	2375	13.2163	13.2141	0.0022	2660	13.5606	13.5619	-0.0013		
22	4.13594750 4.58849794	2380	13.2221	13.2205	0.0016	2660.414706	13.5623	2665	13.5657	13.5666	-0.0009
		2385	13.2280	13.2268	0.0012	3710.613294	13.5623	2670	13.5707	13.5712</	

## REFERENCES

- Bullen K. E. & Bolt Bruce A., 1985, *An introduction to the theory of seismology*, Cambridge University Press, Cambridge, 499 pp.
- Clayton, R. W. & Comer, R. P., 1983, A tomographic analysis of mantle heterogeneities from body wave travel times, *Eos Trans. AGU*, 776.
- Dziewonski, A. M. & Anderson, D. L., 1981, Preliminary reference Earth model, *Phys. Earth. planet. Interiors*, **25**, 297-356.
- Dziewonski, A. M., 1984, Mapping the lower mantle: Determination of lateral heterogeneity in P velocity up to degree and order 6, *J. Geophys. Res.*, **89**, 5929-5952.
- Herrin et al., 1968, 1968 seismological tables for P phases, *Bull. Seism. Soc. Am.*, Vol. 58, No 4, pp. 1193-1235.
- Jeffreys, H. & Bullen, K. E., 1958, *Seismological Tables*, British Association for the Advancement of Science, London.
- Kennett, B. L. N. & Engdahl, E. R., 1991, Traveltimes for global earthquake location and phase identification, *Geophys. J. Int.*, **105**, 429-465.
- Lana-Renault, Yoël & Cid Palacios, Rafael, 1991, *On the problem of the internal constitution of the Earth*, Academia de Ciencias de Zaragoza, Univ. Zaragoza, Vol. 4, 158 pp.
- Lana-Renault, Yoël, 1998, *Modelo de constitución interna de la Tierra*, Doctoral Dissertation, Departamento de Física Teórica, Univ. Zaragoza, 146 pp.
- Morelli, A. & Dziewonski, A. M., 1993, Body wave traveltimes and a spherically symmetric P- and S-wave velocity model, *Geophys. J. Int.* **112**, 178-194.
- Pulliam, R. J., Vasco, D. W. & Johnson L. R., 1993, Tomographic inversions for mantle P wave velocity structure based on the minimisation of  $l^2$  and  $l^1$  norms of International Seismological Centre travel time residuals, *J. Geophys. Res.*, **98**, 699-734.
- Sengupta, M.K. & Toksoz, M. N., 1976, Three-dimensional model of seismic velocity variation in the Earth's mantle, *Geophys. Res. Lett.*, **3**, 84-86.
- Taggart, J. N. & Engdahl, E. R., 1968, Estimation of PcP travel times and the depth to the core, *Bull. Seism. Soc. Am.*, **58**, 1243-1260.