

## Fermat's Last Theorem, Solution Sets v3

Ernest R. Lucier

### Abstract

A solution set is derived for  $C^2 = A^2 + B^2$  where  $n = 2$ . The non-zero integer solution set for  $n = 2$  is derived. The solution set methodology for  $n = 2$  is applied to  $n = 3$ ,  $n = 4$ , and generalized to  $n$ . I will show that there are no non-zero integer solutions for all  $n > 2$ .

### Derive a solution for $C^n = A^n + B^n$ with $n = 2$ .

$C^n = A^n + B^n$  with  $n = 2$  is:

$C^2 = A^2 + B^2$	(1)
-------------------	-----

Rewriting A and B in terms of C, a, and b:

$A = C - a$	(2)
$B = C - b$	(3)

Substitute for A and B in Equation 1:

$C^2 = (C - a)^2 + (C - b)^2$	(4)
-------------------------------	-----

Expand terms

$C^2 = C^2 - 2Ca + a^2 + C^2 - 2Cb + b^2$	(5)
---	-----

Combine like terms

$C^2 - 2C(a + b) + a^2 + b^2 = 0$	(6)
-----------------------------------	-----

Complete the square

$C^2 - 2C(a + b) + (a + b)^2 = \{(a + b)^2 - (a^2 + b^2)\}$	(7)
and	
$C^2 - 2C(a + b) + (a + b)^2 = 2ab$	(8)
and	
$[C - (a + b)]^2 = 2ab$	(9)

Take the square root of both sides of Equation 9:

$C - (a + b) = \pm (2ab)^{1/2}$	(10)
---------------------------------	------

Solve for C

$C = a + b \pm (2ab)^{1/2}$	(11)
-----------------------------	------

Equation 11 is the solution for  $C^n = A^n + B^n$  with  $n = 2$ .

**Define the non-zero integer solution set for  $n = 2$ .**

Factor both sides of Equation 9:

$[C - (a + b)][C - (a + b)] = 2*a*b$	(12)
--------------------------------------	------




Figure 1. Physical representation of Equations 9 and 11 (Not to scale)

Notes:

1.  $2*ab$  completes the square, Equation 9. The area completing the square equals the common core area,  $[C - (a + b)]^2$ .
2.  $[C - (a + b)]^2$  is area common to both  $A^2$  and  $B^2$ , i.e., overlapping areas. The area inside  $C^2$  and outside  $A^2$  and  $B^2$  (i.e.,  $2*ab$ ) must equal the common area  $[C - (a + b)]^2$  (i.e., common core). A common core can exist only if  $C > (a + b)$ .

Define M in terms of a and b to ensure that  $(2*a*b)$  of Equation 11 will be a whole number.

Let  $2*M^2 = a*b$ .

$2M^2 = ab$	(13)
$[C - (a + b)]*[C - (a + b)] = 2*2*M^2$	(14)
Then	
$C = a + b \pm 2*M$	(15)

Equation 15 is the non-zero integer solution set for Equation 1,  $C^2 = A^2 + B^2$ , when M is a non-zero integer and a and b are non-zero integer factors of  $2M^2$ . There are an infinite number of non-zero integer solutions (i.e., Pythagorean triples) for  $C^2 = A^2 + B^2$  where  $M = 1, 2, 3 \dots$

**Example:** let  $M = 1$  so that  $2M^2 = 2$

Using Equations 13, 15, 2, and 3:

The factors of  $2M^2$  are  $a = 1$  and  $b = 2$  since  $2M^2 = 2$ .

$C = 1 + 2 + 2, C = 5, \text{ and } C^2 = 25$	(16)
$A = C - a, A = 5 - 1, A = 4, \text{ and } A^2 = 16$	(17)
$B = C - b, B = 5 - 2, B = 3, \text{ and } B^2 = 9$	(18)

Therefore, when  $n = 2$  and  $M = 1$ , this Equation results in a 3, 4, 5 triangle, i.e.,  $9 + 16 = 25$ .

**Derive a solution for  $C^n = A^n + B^n$  with  $n = 3$ .**

$C^n = A^n + B^n$  with  $n = 3$  is:

$C^3 = A^3 + B^3$	(19)
-------------------	------

Rewriting A and B in terms of C, a, and b as in  $n = 2$  (Equations 2 and 3):

$A = C - a$	(20)
$B = C - b$	(21)

Substitute for A and B in Equation 19:

$C^3 = (C - a)^3 + (C - b)^3$	(22)
-------------------------------	------

Expand terms

$C^3 = C^3 - 3C^2a + 3Ca^2 - a^3 + C^3 - 3C^2b + 3Cb^2 - b^3$	(23)
---	------

Combine like terms

$C^3 - 3C^2(a + b) + 3C(a^2 + b^2) - (a^3 + b^3) = 0$	(24)
---	------

Complete the cube

$C^3 - 3C^2(a + b) + 3C(a + b)^2 - (a + b)^3 =$ $3C\{(a + b)^2 - (a^2 + b^2)\} - \{(a + b)^3 - (a^3 + b^3)\}$ and $C^3 - 3C^2(a + b) + 3C(a + b)^2 - (a + b)^3 = 3C(2ab) - (3a^2b + 3ab^2)$ and $[C - (a + b)]^3 = 3ab[2C - (a + b)]$	(25)
$C^3 - 3C^2(a + b) + 3C(a + b)^2 - (a + b)^3 = 3C(2ab) - (3a^2b + 3ab^2)$	(26)
$[C - (a + b)]^3 = 3ab[2C - (a + b)]$	(27)

Take the cube root of both sides of Equation 27.

$C - (a + b) = \{3ab[2C - (a + b)]\}^{1/3}$	(28)
---	------

Solve for C

$C = a + b + \{3ab[2C - (a + b)]\}^{1/3}$	(29)
---	------

Equation 29 is the solution for  $C^n = A^n + B^n$  with  $n = 3$ .

**Define the non-zero integer solution set for  $n = 3$ .**

Factor both sides of Equation 27:

$[C - (a + b)] * [C - (a + b)] * [C - (a + b)] = 3 * a * b * [2C - (a + b)]$	$(30)$
--	--------




Figure 2. Physical representation of Equation 30 (Not to scale)

Notes:

1.  $3 * Cab + 3 * ab(C - (a + b)) = 3ab[2C - (a + b)]$  completes the cube, Equation 27.

The volume completing the cube equals the common core volume,  $[C - (a + b)]^3$ .

2.  $[C - (a + b)]^3$  is volume common to both  $A^3$  and  $B^3$ . The volume inside  $C^3$  and outside  $A^3$  and  $B^3$  (i.e.,  $3ab[2C - (a + b)]$ ) must equal the common volume  $[C - (a + b)]^3$  (i.e., common core). A common core exist only if  $C > (a + b)$ .

Divide both sides of Equation 27 by  $[C - (a + b)]$ .

$[C - (a + b)]*[C - (a + b)] = 3*a*b*[2C - (a + b)] \div [C - (a + b)]$	(31)
$[C - (a + b)]*[C - (a + b)] = 3*a*b*\{C + [C - (a + b)] \div [C - (a + b)]\}$	(32)
$[C - (a + b)]*[C - (a + b)] = 3*a*b*\{C \div [C - (a + b)] + [C - (a + b)] \div [C - (a + b)]\}$	(33)
$[C - (a + b)]*[C - (a + b)] = 3*a*b*\{C \div [C - (a + b)] + 1\}$	(34)

Add and subtract  $(a + b)$  to  $C$  in the numerator on the right side of the Equation 34 and rewrite:

$[C - (a + b)]*[C - (a + b)] = 3*a*b*\{[C - (a + b) + (a + b)] \div [C - (a + b)] + 1\}$	(35)
$[C - (a + b)]*[C - (a + b)] = 3*a*b*\{1 + \{(a + b) \div [C - (a + b)]\} + 1\}$	(36)
$[C - (a + b)]*[C - (a + b)] = 3*a*b*\{2 + \{(a + b) \div [C - (a + b)]\}\}$	(37)

The non-zero integer solution set for  $[C - (a + b)]^2$  from Equation 9 is  $[2ab]$  when  $M$  is a non-zero integer and  $a$  and  $b$  are non-zero integer factors of  $2M^2$  with  $2M^2 = ab$  (see Equation 15). Find the value of  $C$  in Equation 37 when  $[C - (a + b)]^2$  equals  $[2ab]$ .

Let  $[C - (a + b)]*[C - (a + b)]$  equal  $[2ab]$  in Equation 37 and solve for  $C$  and  $(a + b)$ .

$[2ab] = 3*a*b*\{2 + \{(a + b) \div [C - (a + b)]\}\}$	(38)
--	------



Divide Equation 38 by  $3ab$  and combine terms:

$\frac{2}{3} = 2 + \{(a + b) \div [C - (a + b)]\}$	(39)
$\frac{2}{3} - 2 = (a + b) \div [C - (a + b)]$	(40)
$-\frac{4}{3} = (a + b) \div [C - (a + b)]$	(41)
$-\frac{4}{3} * [C - (a + b)] = (a + b)$	(42)
$C - (a + b) = -\frac{3}{4} * (a + b)$	(43)
$C = -\frac{3}{4} * (a + b) + (a + b)$	(44)
$C = (1 - \frac{3}{4}) * (a + b)$	(45)
$C = \frac{1}{4} * (a + b)$	(46)

$C$  is less than  $(a + b)$ . A common core exists only when  $C$  is greater than  $(a + b)$ . Therefore, there is no common core and there are no non-zero integer solutions for  $n = 3$ .

**Derive a solution for  $C^n = A^n + B^n$  with  $n = 4$ .**

$C^n = A^n + B^n$  with  $n = 4$  is:

$C^4 = A^4 + B^4$	(47)
-------------------	------

Rewriting A and B in terms of C, a, and b as in  $n = 2$  (Equations 2 and 3):

$A = C - a$	(48)
$B = C - b$	(49)

Substitute for A and B in Equation 47:

$C^4 = (C - a)^4 + (C - b)^4$	(50)
-------------------------------	------

Expand terms

$C^4 = C^4 - 4C^3a + 6C^2a^2 - 4Ca^3 + a^4 + C^4 - 4C^3b + 6C^2b^2 - 4Cb^3 + b^4$	(51)
---	------

Combine like terms

$C^4 - 4C^3(a + b) + 6C^2(a^2 + b^2) - 4C(a^3 + b^3) + (a^4 + b^4) = 0$	(52)
---	------

Complete the fourth power

$C^4 - 4C^3(a + b) + 6C^2(a + b)^2 - 4C(a + b)^3 + (a^4 + b^4) =$ $6C^2\{(a + b)^2 - (a^2 + b^2)\} - 4C\{(a + b)^3 - (a^3 + b^3)\} + \{(a + b)^4 - (a^4 + b^4)\}$ <p style="text-align: center;">and</p> $C^4 - 4C^3(a + b) + 6C^2(a + b)^2 - 4C(a + b)^3 + (a^4 + b^4) =$ $6C^2(2ab) - 4C(3a^2b + 3ab^2) + (4a^3b + 6a^2b^2 + 4ab^3)$ <p style="text-align: center;">and</p> $[C - (a + b)]^4 = 2ab[6C^2 - 6C(a + b) + (2a^2 + 3ab + 2b^2)]$	<p>(53)</p> <p>(54)</p> <p>(55)</p>
---	-------------------------------------

Take the fourth root of both sides of Equation 55.

$C - (a + b) = \{2ab[6C^2 - 6C(a + b) + (2a^2 + 3ab + 2b^2)]\}^{1/4}$	<p>(56)</p>
---	-------------

Solve for C

$C = a + b + \{2ab[6C^2 - 6C(a + b) + (2a^2 + 3ab + 2b^2)]\}^{1/4}$	<p>(57)</p>
---	-------------

Equation 57 is the solution for  $C^n = A^n + B^n$  with  $n = 4$ .

**Define the non-zero integer solution set for  $n = 4$ .**

Factor both sides of Equation 55.

$[C - (a + b)]*[C - (a + b)]*[C - (a + b)]*[C - (a + b)]$ $= 2ab[6C^2 - 6C(a + b) + (2a^2 + 3ab + 2b^2)]$	(58)
---	------

The fourth root of the left side of Equation 58 is  $[C - (a + b)]$ . The fourth root of the right side must also Equal  $[C - (a + b)]$  and both sides can be divided by  $[C - (a + b)]^2$ .

$[C - (a + b)]^2 = 2ab[6C^2 - 6C(a + b) + (2a^2 + 3ab + 2b^2)] \div [C - (a + b)]^2$	(59)
--	------

C must be greater than  $(a + b)$  for a common core to exist. Let  $C = (a + b + x)$  and  $x = 1$  in the right side of Equation 59 and solve for  $[C - (a + b)]^2$ .

$[C - (a + b)]^2 = 2ab[6C^2 - 6C(a + b) + (2a^2 + 3ab + 2b^2)]$	(60)
$[C - (a + b)]^2 = 2ab[6(a + b + 1)^2 - 6(a + b + 1)(a + b) + (2a^2 + 3ab + 2b^2)]$	(61)
$[C - (a + b)]^2 = 2ab*[6(a + b + 1) + (2a^2 + 3ab + 2b^2)]$	(62)

$[C - (a + b)]^2$  must equal  $[2ab]$  since the non-zero integer solution set for  $[C - (a + b)]*[C - (a + b)]$  from Equation 9 is  $[2ab]$  when M is a non-zero integer and a and b are non-zero integer factors of  $2M^2$  with  $2M^2 = ab$  (see Equation 15).

From Equation 62  $[C - (a + b)]^2$  does not equal  $[2ab]$  when C is greater than  $(a + b)$  and  $n = 4$  since:

$[2ab] < 2ab*[6(a + b + 1) + (2a^2 + 3ab + 2b^2)]$ <p style="text-align: center;">for all C greater than <math>(a + b)</math></p>	(63)
---	------

Therefore, there is no non-zero integer solution set for  $n = 4$ .

**Derive a solution for  $C^n = A^n + B^n$  for all  $n$ .**

$C^n = A^n + B^n$  for all  $n$ :

$C^n = A^n + B^n$	(64)
-------------------	------

Rewriting A and B in terms of C, a, and b as in  $n = 2$  (Equations 2 and 3):

$A = C - a$	(65)
$B = C - b$	(66)

Substitute for A and B in Equation 64:

$C^n = (C - a)^n + (C - b)^n$	(67)
-------------------------------	------

Expand terms

$C^n =$ $C^n - nC^{n-1}a + [{}^n_2]C^{n-2}a^2 - [{}^n_3]C^{n-3}a^3 + \dots \pm [{}^n_n]a^n +$ $C^n - nC^{n-1}b + [{}^n_2]C^{n-2}b^2 - [{}^n_3]C^{n-3}b^3 + \dots \pm [{}^n_n]b^n$	(68)
<p>Where the binomial coefficient <math>[{}^n_r] = n! / [(n - r)!r!]</math> is the coefficient of the <math>x^r</math> term in the polynomial expansion of the binomial power <math>(1 + x)^n</math> and <math>[{}^n_r] = 0</math> if <math>r &gt; n</math>.</p>	

Combine like terms

$C^n - nC^{n-1}(a + b) + [{}^n_2]C^{n-2}(a^2 + b^2) - [{}^n_3]C^{n-3}(a^3 + b^3) + \dots \pm [{}^n_n](a^n + b^n) = 0$	(69)
---	------

Complete the  $n^{\text{th}}$  power

$C^n - nC^{n-1}(a+b) + [{}^n_2]C^{n-2}(a+b)^2 - [{}^n_3]C^{n-3}(a+b)^3 + \dots \pm (a+b)^n =$ $[{}^n_2]C^{n-2}\{(a+b)^2 - (a^2+b^2)\} - [{}^n_3]C^{n-3}\{(a+b)^3 - (a^3+b^3)\} + \dots$ $\pm [{}^n_n]C^{n-n}\{(a+b)^n - (a^n+b^n)\}$ <p style="text-align: center;">and</p> $C^n - nC^{n-1}(a+b) + [{}^n_2]C^{n-2}(a+b)^2 - [{}^n_3]C^{n-3}(a+b)^3 + \dots \pm (a+b)^n =$ $[{}^n_2]C^{n-2}(2ab) - [{}^n_3]C^{n-3}(3a^2b + 3ab^2) + \dots$ $\pm [{}^n_n]C^{n-n}\{[{}^n_1]a^{n-1}b + [{}^n_2]a^{n-2}b^2 + [{}^n_3]a^{n-3}b^3 + \dots + [{}^n_{n-1}]ab^{n-1}\}$ <p style="text-align: center;">and</p> $[C - (a+b)]^n =$ $[{}^n_2]C^{n-2}(2ab) - [{}^n_3]C^{n-3}(3a^2b + 3ab^2) + \dots$ $\pm [{}^n_n]\{[{}^n_1]a^{n-1}b + [{}^n_2]a^{n-2}b^2 + [{}^n_3]a^{n-3}b^3 + \dots + [{}^n_{n-1}]ab^{n-1}\}$ <p style="text-align: center;">or</p> $[C - (a+b)]^n = ab([{}^n_2]C^{n-2}(2) - [{}^n_3]C^{n-3}(3a + 3b) + \dots$ $\pm [{}^n_n]\{[{}^n_1]a^{n-2} + [{}^n_2]a^{n-3}b^1 + [{}^n_3]a^{n-4}b^2 + \dots + [{}^n_{n-1}]b^{n-2}\})$	<p style="text-align: right;">(70)</p> <p style="text-align: right;">(71)</p> <p style="text-align: right;">(72)</p> <p style="text-align: right;">(73)</p>
---	---

Note: The  $[{}^n_n]$  term is the last term in the series. That is, the first term is the last term when  $n = 2$ , the second term when  $n = 3$ , etc.

Take the  $n^{\text{th}}$  root of both sides of Equation 73.

$C - (a+b) = a^{1/n}b^{1/n}([{}^n_2]C^{n-2}(2) - [{}^n_3]C^{n-3}(3a + 3b) + \dots$ $\pm [{}^n_n]\{[{}^n_1]a^{n-2} + [{}^n_2]a^{n-3}b^1 + [{}^n_3]a^{n-4}b^2 + \dots + [{}^n_{n-1}]b^{n-2}\})^{1/n}$	(74)
---	------

Solve for C

$C = a + b + a^{1/n}b^{1/n}([{}^n_2]C^{n-2}(2) - [{}^n_3]C^{n-3}(3a + 3b) + \dots$ $\pm [{}^n_n]\{[{}^n_1]a^{n-2} + [{}^n_2]a^{n-3}b^1 + [{}^n_3]a^{n-4}b^2 + \dots + [{}^n_{n-1}]b^{n-2}\})^{1/n}$	(75)
---	------

Equation 75 is the solution for  $C^n = A^n + B^n$  for any  $n$ .

**Define the non-zero integer solution set for any n.**

Factor both sides of Equation 73.

$[C - (a + b)]_1 * [C - (a + b)]_2 * \dots * [C - (a + b)]_{n-1} * [C - (a + b)]_n =$ $a * b * ([^n_2] C^{n-2}(2) - [^n_3] C^{n-3}(3a + 3b) + \dots$ $\pm [^n_n] \{ [^n_1] a^{n-2} + [^n_2] a^{n-3} b^1 + [^n_3] a^{n-4} b^2 + \dots + [^n_{n-1}] b^{n-2} \})$	(76)
--	------

The  $n^{\text{th}}$  root of the left side of Equation 76 is  $[C - (a + b)]$ .  $[C - (a + b)]^n$  can be divided by  $[C - (a + b)]^{n-2}$  to obtain  $[C - (a + b)]^2$ . The non-zero integer solution set for  $[C - (a + b)]^2$  is  $2ab$  when  $a$  and  $b$  are non-zero integer factors of  $2M^2$  and  $M$  is a non-zero integer. (See Equation 15).

Divide both sides of Equation 73 by  $[C - (a + b)]^{n-2}$ .

$[C - (a + b)]^2 =$ $a * b * ([^n_2] C^{n-2}(2) - [^n_3] C^{n-3}(3a + 3b) + \dots$ $\pm [^n_n] \{ [^n_1] a^{n-2} + [^n_2] a^{n-3} b^1 + [^n_3] a^{n-4} b^2 + \dots + [^n_{n-1}] b^{n-2} \}) \div [C - (a + b)]^{n-2}$	(77)
---	------

Solve for  $[C - (a + b)]^2$  for  $n = 2, 3, 4$ , and all  $n$ .

For n = 2 rewrite Equation 77 with n = 2.

(Note: The  $[\cdot]_n$  term in Equation 77 is the first term when n = 2):

$[C - (a + b)]^2 = a*b*([2]_2 C^{2-2}(2)) \div [C - (a + b)]^{2-2}$	(78)
$[C - (a + b)]^2 = 2ab$	(79)

Equation 79 is the same as Equation 9 (n = 2). The non-zero integer solution set for  $[C - (a + b)]^2$  from Equation 9 is  $[2ab]$  when M is a non-zero integer and a and b are non-zero integer factors of  $2M^2$  with  $2M^2 = ab$  (see Equation 15). There are an infinite number of non-zero integer solutions (i.e., Pythagorean triples) for  $C^2 = A^2 + B^2$  where M = 1, 2, 3 ...

For n = 3 rewrite Equation 77 with n = 3.

(Note: The  $[\cdot]_n$  term in Equation 77 is the second term when n = 3):

$[C - (a + b)]^2 = a*b*([3]_3 C^{3-2}(2) - [3]_3 C^{3-3}(3a + 3b)) \div [C - (a + b)]^{3-2}$	(80)
$[C - (a + b)]^2 = 3ab(2C - (a + b)) \div [C - (a + b)]$	(81)

C must be greater than (a + b) for a common core to exist (see Figures 1 and 2). The non-zero integer solution set for  $[C - (a + b)]^2$  from Equation 9 is  $[2ab]$  when M is a non-zero integer and a and b are non-zero integer factors of  $2M^2$  with  $2M^2 = ab$  (see Equation 15).

From Equation 46,  $C = 1/4 * (a + b)$  when n = 3, C is not greater than (a + b), and  $[C - (a + b)]^2$  can not equal  $[2ab]$  when n = 3.

Therefore, there is no non-zero integer solution set for n = 3.



For n = 4 rewrite Equation 77 with n = 4.

(Note: The  $[^n_n]$  term in Equation 77 is the third term when n = 4):

$[C - (a + b)]^2 = a*b*([^4_2]C^{4-2}(2) - [^4_3]C^{4-3}(3a + 3b) + [^4_4]C^{4-4}(4a^2 + 6ab + 4b^2))$ $\div [C - (a + b)]^{4-2}$	(82)
$[C - (a + b)]^2 = a*b*(6C^2(2) - 4C(3a + 3b) + (4a^2 + 6ab + 4b^2)) \div [C - (a + b)]^2$	(83)
$[C - (a + b)]^2 = 2ab[6C^2 - 6C(a + b) + (2a^2 + 3ab + 2b^2)] \div [C - (a + b)]^2$	(84)

Equation 84 is the same as Equation 59.

Let  $C = (a + b + x)$  and  $x = 1$  in the right side of Equation 81 and solve for  $[C - (a + b)]^2$ .

$[C - (a + b)]^2 = 2*a*b*(6C^2 - 6C(a + b) + (2a^2 + 3ab + 2b^2))$	(85)
$[C - (a + b)]^2 = 2*a*b*(6(a + b + 1)^2 - 6(a + b + 1)(a + b) + (2a^2 + 3ab + 2b^2))$	(86)
$[C - (a + b)]^2 = 2ab*(6(a + b + 1) + (2a^2 + 3ab + 2b^2))$	(87)

C must be greater than (a + b) for a common core to exist (see Figures 1 and 2). The non-zero integer solution set for  $[C - (a + b)]^2$  from Equation 9 is  $[2ab]$  when M is a non-zero integer and a and b are non-zero integer factors of  $2M^2$  with  $2M^2 = ab$  (see Equation 15). The quantity equal to  $[C - (a + b)]^2$  is always greater than  $[2ab]$  when C is greater than (a + b) when n = 4 from Equation 87.

Therefore, there is no non-zero integer solution set for n = 4.

For  $n = n$  repeat Equation 73:

$[C - (a + b)]^n = ab\left(\binom{n}{2}C^{n-2}(2) - \binom{n}{3}C^{n-3}(3a + 3b) + \dots\right. \\ \left. \pm \binom{n}{n} \{ \binom{n}{1}a^{n-2} + \binom{n}{2}a^{n-3}b^1 + \binom{n}{3}a^{n-4}b^2 + \dots + \binom{n}{n-1}b^{n-2} \} \right)$	(88)
---	------

The root of both sides of Equation 88 (and 73) is  $[C - (a + b)]$ .

Divide Equation 88 by  $[C - (a + b)]^{n-2}$ .

$[C - (a + b)]^2 = a*b*\left(\binom{n}{2}C^{n-2}(2) - \binom{n}{3}C^{n-3}(3a + 3b) + \dots\right. \\ \left. \pm \binom{n}{n} \{ \binom{n}{1}a^{n-2} + \binom{n}{2}a^{n-3}b^1 + \binom{n}{3}a^{n-4}b^2 + \dots + \binom{n}{n-1}b^{n-2} \} \right) \div [C - (a + b)]^{n-2}$	(89)
--	------

The non-zero integer solution set for  $[C - (a + b)]^2$  is  $2ab$  when the variables  $a$  and  $b$  are non-zero integer factors of  $2M^2$  and  $M$  is a non-zero integer. Therefore, the quantity equal to  $[C - (a + b)]^2$  in Equation 89 must equal  $2ab$  for whole number solutions to exist. Also,  $C$  must be greater than  $(a + b)$  for a common core to exist (see Figures 1 and 2).

Note: The  $\binom{n}{n}$  in Equations 88 and 89 are binomial coefficients. The number of applicable terms in Equation 89 increase as  $n$  increases (i.e., number of applicable terms in Equation 89 equals  $n - 1$ ). Completion of the  $n$ th power always begins with the  $\binom{n}{2}$  term in the series and ends with  $(n - 1)$  terms (see A in Figure 3). The quantity equal to  $[C - (a + b)]^2$  is always greater than  $2ab$  when  $n$  is greater than 2.

		Applicable terms - A	
1			
1 1			
1 -2	1	n = 2	A = 1
1 -3	3 -1	n = 3	A = 2
1 -4	6 -4 1	n = 4	A = 3
1 -5	10 -10 5 -1	n = 5	A = 4
1 -6	15 -20 15 -6 1	n = 6	A = 5
1 -7	21 -35 35 -21 7 -1	n = 7	A = 6
1 -8	28 -56 70 -56 28 -8 1	n = 8	A = 7
1 -9	36 -84 126 -126 84 -36 9 -1	n = 9	A = 8
1 -10	45 -120 210 -252 210 -120 45 -10 1	n = 10	A = 9

Figure 3. Binomial Coefficients for n = 2 to n = 10 for Equations 88 and 89. The highlighted numbers are the binomial coefficients for completing the n<sup>th</sup> power beginning with  $\binom{n}{2}$ .

For example:

Completion of the square for n = 2 is  $ab(\binom{2}{2}C^{2-2}(2)) = 2ab$

Completion of the cube for n = 3 is  $ab(\binom{3}{2}C^{3-2}(2) - \binom{3}{3}C^{3-3}(3a + 3b))$

$$= ab(\binom{3}{2}C^1(2) - \binom{3}{3}(3a + 3b)) = 3ab(2C - (a + b)), \text{ the same as Equation 27.}$$

Completion of the fourth power for n = 4 is  $ab(\binom{4}{2}C^{4-2}(2) - \binom{4}{3}C^{4-3}(3a + 3b) + \binom{4}{4}C^{4-4}(2a^2 + 3ab + 2b^2))$

$$= 2ab[6C^2 - 6C(a + b) + (2a^2 + 3ab + 2b^2)]$$

The quantity  $[C - (a + b)]^2$  in Equation 89 equals 2ab only when n = 2. Therefore, there is no non-zero integer solution set for any n greater than 2 since the quantity equal to  $[C - (a + b)]^2$  is always greater than 2ab when n is greater than 2 and C is greater than (a + b).

#### Bibliography:

My cursory review of approaches to a proof of Fermat's Last Theorem did not find any similar to the one in this paper. Two references I relied on most are:

Fermat's Last Theorem (Vandiver 1914; Vandiver publisher; L'Intermédiaire des Math. 15, 217-218.;

Swinnerton-Dwyer, P. Nature 364, 13--14, 1993) found at

<http://mathworld.wolfram.com/FermatsLastTheorem.html>.

Fermat's Enigma, by Simon Singh, 1997, Anchor Books, New York