

On the Homotopy Type of the Fredholm Lagrangian-Grassmannian

J. C. C. Eidam, P. Piccione*

February 24, 2005

Abstract

In this article, using the results of [1], we show that the Fredholm Lagrangian-Grassmannian $\mathcal{F}_{L_0}(\Lambda)$ has the homotopy type of the quotient $U(\infty)/O(\infty)$. As a corollary, we obtain that the Maslov index with respect to a lagrangian is a isomorphism between the fundamental group of $\mathcal{F}_{L_0}(\Lambda)$ and the integers. We remark that the fundamental group $\pi_1(\mathcal{F}_{L_0}(\Lambda))$ was computed by B. Booss-Bavnbek and K. Furutani in [2], [4], with considerable effort, but the method works only for the π_1 case.

1 Introduction

Let H be a separable infinite dimensional Hilbert space endowed with a symplectic structure, that is, there exists a bounded operator J in H such that $J^2 = -I$ and $J^* = -J$ and a symplectic form ω given by $\omega(\cdot, \cdot) = \langle J\cdot, \cdot \rangle$. The space Λ of all lagrangian subspaces in H admits a structure of real-analytic contractible Banach manifold, so uninteresting from the topological viewpoint. To obtain a space with nontrivial topology, consider $L_0 \in \Lambda$ and the space $\mathcal{F}_{L_0}(\Lambda)$ of all $L \in \Lambda$ which forms a Fredholm pair with L_0 . It is a open set in Λ , so it is also a analytic Banach manifold.

In this article, we prove that $\mathcal{F}_{L_0}(\Lambda)$ is homotopy equivalent with the space of compact perturbations of a fixed lagrangian; this last space is homotopy equivalent with $U(\infty)/O(\infty)$.

2 Proof of the result

Let us fix some notation. Denote by P_M the orthogonal projection over a closed subspace $M \subset H$. Denote by $GL(H)$ the group of all invertible operators in H and by $O(H)$ the group of unitary operators in H . If H is a symplectic space with complex structure J , the group $U(H; J)$ is formed by elements of $O(H)$ which commute with J . We define $GL_c(H) = GL(H) \cap (I + \mathcal{L}_c(H))$, $O_c(H) = O(H) \cap (I + \mathcal{L}_c(H))$ and $U_c(H; J) =$

*Research supported by Fapesp, São Paulo, Brasil, grants 01/00046-3 and 02/02528-8. The second author is partially sponsored by Cnpq, Brasil.

$U(\mathbf{H}; J) \cap (I + \mathcal{L}_c(\mathbf{H}))$, where $\mathcal{L}_c(\mathbf{H})$ denotes the space of compact operators in \mathbf{H} . Elements of $I + \mathcal{L}_c(\mathbf{H})$ are called *compact perturbations of identity*.

We begin by studying some properties of compact perturbations of lagrangians. Given $L, L' \in \Lambda$, L' is a compact perturbation of L if $P_{L'} - P_L$ is compact. This fact is denoted by $L' \sim L$. Obviously, \sim is a equivalence relation in Λ . The quotient space of this relation is called *essential grassmannian-lagrangian* and is denoted by Λ_e . It is endowed with the quotient topology induced by the canonical map $\pi : \Lambda \rightarrow \Lambda_e$. For $L_0 \in \Lambda$, we define $\mathcal{G}_c(L_0)$ as the equivalence class of L_0 . The next results determine the topology of $\mathcal{G}_c(L_0)$.

LEMMA 2.1 *The group action*

$$\begin{aligned} U_c(\mathbf{H}; J) \times \mathcal{G}_c(L_0) &\rightarrow \mathcal{G}_c(L_0) \\ (U, L) &\mapsto U(L) \end{aligned} \quad (2.1)$$

is transitive.

PROOF. Let $L \in \mathcal{G}_c(L_0)$; we must find $U \in U_c(\mathbf{H}; J)$ such that $U(L_0) = L$. Define $T_1 = I + (P_L - P_{L_0})|_{L_0} = P_L|_{L_0} : L_0 \rightarrow L$; since T_1 is a Fredholm operator with $\text{ind } T_1 = 0$, there exists $K : L_0 \rightarrow L$ of finite rank such that $T_2 = T_1 + K : L_0 \rightarrow L$ is a linear isomorphism. Putting

$$T = T_2 P_{L_0} - J T_2 J P_{L_0}^\perp = I + ((P_L - P_{L_0}) + K) P_{L_0} - J((P_L - P_{L_0}) + K) J P_{L_0}^\perp,$$

we have that $T \in \text{GL}_c(\mathbf{H})$, $TJ = JT$ and $T(L_0) = L$. This last equality implies that $T^*(L^\perp) = L_0^\perp$, so, since $T^*J = JT^*$, we have

$$T^*(L) = T^*J(L^\perp) = JT^*(L^\perp) = J(L_0^\perp) = L_0.$$

In particular, $(T^*T)(L_0) = L_0$. Let S be the unique positive square root for T^*T . It is easy (see for example [10, Chapter VII]) that S can be given as the limit of the sequence $\{S_n\}$ given by $S_0 = 0$ and

$$S_{n+1} = \frac{1}{2}(T^*T + S_n^2), \quad (2.2)$$

for $n \geq 0$. Since T is a compact perturbation of the identity, we see, by formula (2.2) and induction that S is a compact perturbation of the identity, $SJ = JS$ and $S(L_0) = L_0$. Taking $T = US$ the polar decomposition of T , with $U = TS^{-1}$, we see that $U(L_0) = L$ and $U \in U_c(\mathbf{H}; J)$, as desired. \blacksquare

PROPOSITION 2.2 *The space $\mathcal{G}_c(L_0)$ is homotopy equivalent to $U(\infty)/O(\infty)$.*

PROOF. Consider the differentiable action (2.1). Since the isotropy group at L_0 is $\{U \in U_c(\mathbf{H}; J) : U(L_0) = L_0\} \approx O_c(L_0)$, we conclude that there is a diffeomorphism $\mathcal{G}_c(L_0) \approx U_c(\mathbf{H}; J)/O_c(L_0)$. Since the spaces $U_c(\mathbf{H}; J)$ and $O_c(L_0)$ are homotopy equivalent with $U(\infty)$ and $O(\infty)$, respectively, (see [7]), the result follows. \blacksquare

If $U \in U(\mathbf{H}; J)$ and $L \in \Lambda$ then $P_{U(L)} = U P_L U^*$. This fact is important to prove the next theorem, which describes the topology of Λ_e .

THEOREM 2.3 *The canonical map $\pi : \Lambda \rightarrow \Lambda_e$ defines a fibre bundle with typical fiber $\mathcal{G}_c(L_0)$.*

PROOF. Let $L_0 \in \Lambda$ and consider the Calkin algebra $\mathcal{C}(\mathbf{H}) = \mathcal{L}(\mathbf{H})/\mathcal{L}_c(\mathbf{H})$, where $\mathcal{L}(\mathbf{H})$ ($\mathcal{L}_c(\mathbf{H})$) denotes the space of bounded (compact) operators in \mathbf{H} . As proved in [1, Proposition 1.2], if $U(\mathcal{C}(\mathbf{H}))$ denotes the unitary elements of $\mathcal{C}(\mathbf{H})$ (that is, $u \in U(\mathcal{C}(\mathbf{H}))$ if and only if $u^*u = uu^* = 1$), then the map

$$\begin{aligned} U(\mathcal{C}(\mathbf{H})) &\rightarrow U(\mathcal{C}(\mathbf{H})) \cdot [P_{L_0}] \\ u &\mapsto u[P_{L_0}]u^* \end{aligned}$$

is a analytic principal bundle with structural group $\{u \in U(\mathcal{C}(\mathbf{H})) : u[P_{L_0}]u^* = [P_{L_0}]\}$, where $[P_{L_0}]$ denotes the image of P_{L_0} by the canonical map $p : \mathcal{L}(\mathbf{H}) \rightarrow \mathcal{C}(\mathbf{H})$. By restricting the structural group to the closed subgroup

$$G = p(\{U \in U(\mathbf{H}; J) : U(L_0) = L_0\}),$$

we obtain a principal bundle

$$\begin{aligned} \Phi : p(U(\mathbf{H}; J)) &\rightarrow p(U(\mathbf{H}; J)) \cdot [P_{L_0}] \\ u &\mapsto u[P_{L_0}]u^* \end{aligned} \tag{2.3}$$

with G as structural group. The map (2.3) can be rewritten as

$$\begin{aligned} \Phi : p(U(\mathbf{H}; J)) &\rightarrow p(U(\mathbf{H}; J)) \cdot [P_{L_0}] \approx \Lambda_e \\ u &\mapsto [P_{U(L_0)}], \end{aligned}$$

where $U \in U(\mathbf{H}; J)$ is such that $p(U) = u$. Since we can obtain local sections for the map Φ and the orbit $p(U(\mathbf{H}; J)) \cdot [P_{L_0}]$ can be identified with the space Λ_e , there exists a neighbourhood $\mathcal{U} \subset \Lambda_e$ of $\pi(L_0)$ and a map $s' : \mathcal{U} \rightarrow p(U(\mathbf{H}; J))$ such that $\Phi \circ s' = I$. Now, again by the results of [1, Proposition 1.7], the map $p|_{U(\mathbf{H}; J)} : U(\mathbf{H}; J) \rightarrow p(U(\mathbf{H}; J))$ admit local sections around all points of $p(U(\mathbf{H}; J))$, so, we obtain a neighbourhood $\mathcal{V} \subset p(U(\mathbf{H}; J))$ of I and a map $s'' : \mathcal{V} \rightarrow U(\mathbf{H}; J)$ such that $p \circ s'' = I$. Putting $s = s'' \circ s' : \mathcal{U} \rightarrow U(\mathbf{H}; J)$, the map

$$\begin{aligned} \mathcal{U} \times \mathcal{G}_c(L_0) &\rightarrow \pi^{-1}(\mathcal{U}) \\ ([L], L') &\mapsto s([L])(L') \end{aligned}$$

is a local trivialization, proving the result. ■

By theorem 2.3, the long exact sequence in homotopy and the contractibility of Λ , we have the next result.

COROLLARY 2.4 *There exists isomorphisms $\pi_j(\Lambda_e) \approx \pi_{j-1}(\mathcal{G}_c(L_0)) \approx \pi_{j-1}(U(\infty)/O(\infty))$ for all $j \in \mathbb{N}$.*

A pair (L_0, L_1) of lagrangians of \mathbf{H} is said *complementary* if $L_0 \cap L_1 = 0$ and $L_0 + L_1 = \mathbf{H}$. The set of all lagrangians complementary to L_1 is denoted by $\Lambda_0(L_1)$. Define

$$\begin{aligned} \varphi_{L_0, L_1} : \Lambda_0(L_1) &\rightarrow \mathcal{L}^{\text{sa}}(L_0) \\ L &\mapsto P_{L_0}JS, \end{aligned}$$

where $S : L_0 \rightarrow L_1$ has graph $\text{Gr}(S) = L$ and $\mathcal{L}^{\text{sa}}(L_0)$ denotes the Banach space of bounded self-adjoint operators in L_0 . This map is a bijection and the collection $\{\varphi_{L_0, L_1}\}$, where (L_0, L_1) runs over the set of all complementary lagrangians in \mathbf{H} , is a real-analytic atlas in Λ which makes it a Banach manifold modeled on the space of bounded self-adjoint operators on a Hilbert space. More details can be found in [3], [6].

LEMMA 2.5 *Given a pair (L_0, L_1) of complementary lagrangians in H and $L \in \Lambda_0(L_1)$, we have that (L, L_0) is a Fredholm pair if and only if $\varphi_{L_0, L_1}(L)$ is a Fredholm operator.*

PROOF. Since $\ker \varphi_{L_0, L_1}(L) = L \cap L_0$, we have that $\dim \ker \varphi_{L_0, L_1}(L) = \dim(L \cap L_0)$. The inequalities

$$\gamma(L_0, L_1) \operatorname{dist}(u, L \cap L_0) \leq \operatorname{dist}(u + Su, L \cap L_0) \leq \|I + S\| \operatorname{dist}(u, L \cap L_0),$$

where $\gamma(L_0, L_1)$ is the usual minimum gap between L_0 and L_1 (see [5, Chapter IV]) and $S : L_0 \rightarrow L_1$ has graph L , imply that

$$\|I + S\|^{-1} \frac{\|P_{L_0}JSu\|}{\operatorname{dist}(u, L \cap L_0)} \leq \frac{\operatorname{dist}(u + Su, L_0)}{\operatorname{dist}(u + Su, L \cap L_0)} \leq \gamma(L_0, L_1)^{-1} \frac{\|P_{L_0}JSu\|}{\operatorname{dist}(u, L \cap L_0)},$$

so,

$$\|I + S\|^{-1} \gamma(P_{L_0}JS) \leq \gamma(L, L_0) \leq \gamma(L_0, L_1)^{-1} \gamma(P_{L_0}JS),$$

where $\gamma(P_{L_0}JS)$ denotes the minimum modulus of $P_{L_0}JS$, as defined in [5, Chapter IV]. In particular, $L + L_0$ is closed if and only if $P_{L_0}JS$ has closed range. Since $P_{L_0}JS$ is self-adjoint and $(L + L_0)^\perp = J(L_0 \cap L_1)$, we conclude the result. ■

Since the set of Fredholm self-adjoint operators is open in the space of all self-adjoint operators, we conclude that $\mathcal{F}_{L_0}(\Lambda) \subset \Lambda$ is open. Before proving our principal result, we need a simple lemma which says that a Fredholm pair of lagrangians is a complementary pair modulo a compact perturbation.

LEMMA 2.6 *If (L_0, L_1) is a Fredholm pair of lagrangians then there exists $L'_1 \in \Lambda$, $L'_1 \sim L_1$, such that (L_0, L'_1) is a complementary pair.*

PROOF. Let L_2 be a lagrangian complementary to L_1 and L_0 and $S \in \mathcal{L}(L_0, L_2)$ such that $\operatorname{Gr}(S) = L_1$. By lemma 2.5, $T = \varphi_{L_0, L_2}(L_1)$ is a Fredholm operator in L_0 . Defining $T' = T + P_{\ker T}$ and $S' = (P_{L_0}J|_{L_2})^{-1}T'$, we obtain that T' is an isomorphism and $L'_1 = \operatorname{Gr}(S')$ is a lagrangian complementary to L_0 . Since S' is a compact perturbation of S , we have that $L'_1 = \operatorname{Gr}(S') \sim \operatorname{Gr}(S) = L_1$. This proves the result. ■

The next theorem is the principal result of this article.

THEOREM 2.7 *There exists a homotopy equivalence $\mathcal{F}_{L_0}(\Lambda) \approx U(\infty)/O(\infty)$.*

PROOF. Consider the restriction $\chi = \pi|_{\mathcal{F}_{L_0}(\Lambda)} : \mathcal{F}_{L_0}(\Lambda) \rightarrow \pi(\mathcal{F}_{L_0}(\Lambda))$ of the fiber bundle $\pi : \Lambda \rightarrow \Lambda_e$. Since $\mathcal{F}_{L_0}(\Lambda)$ is stable under compact perturbations, the map χ defines a fiber bundle with typical fiber $\mathcal{G}_c(L_0)$. The previous lemma shows that $\pi(\mathcal{F}_{L_0}(\Lambda)) = \pi(\Lambda_0(L_0))$. The set $\Lambda_0(L_0)$ is homeomorphic to $\mathcal{L}^{\text{sa}}(L_0^\perp)$ via the usual map

$$\begin{aligned} \varphi_{L_0^\perp, L_0} : \Lambda_0(L_0) &\rightarrow \mathcal{L}^{\text{sa}}(L_0^\perp) \\ L &\mapsto JS, \end{aligned}$$

where $S : L_0^\perp \rightarrow L_0$ has graph $\operatorname{Gr}(S) = L$ and $\mathcal{L}^{\text{sa}}(L_0^\perp)$ denotes the Banach space of bounded self-adjoint operators in L_0 . Now, the map

$$\begin{aligned} \pi(\Lambda_0(L_0)) \times [0, 1] &\rightarrow \pi(\Lambda_0(L_0)) \\ (\pi(\varphi_{L_0^\perp, L_0}^{-1}(T)), t) &\mapsto \pi(\varphi_{L_0^\perp, L_0}^{-1}(tT)) \end{aligned}$$

is well-defined, continuous and gives a homotopy between the identity function on $\pi(\Lambda_0(L_0))$ and a constant. So, $\pi(\Lambda_0(L_0))$ is contractible, and the long exact sequence of the fibration applied to $\chi : \mathcal{F}_{L_0}(\Lambda) \rightarrow \pi(\Lambda_0(L_0))$ implies that exists a weak homotopy equivalence between $\mathcal{F}_{L_0}(\Lambda)$ and $\mathcal{G}_c(L_0)$. The theorem follows by [8, Theorem 15] and proposition 2.2. ■

We end with a result about the Maslov index for curves in $\mathcal{F}_{L_0}(\Lambda)$. Given a curve $\gamma : [0, 1] \rightarrow \Lambda_0(L_1)$, the Maslov index of γ with respect to L_0 is defined as

$$\mu_{L_0}(\gamma) = \text{sf}(\varphi_{L_0, L_1} \circ \gamma),$$

where $\text{sf}(\varphi_{L_0, L_1} \circ \gamma)$ denotes the spectral flow of the curve of Fredholm self-adjoint operators $\varphi_{L_0, L_1} \circ \gamma$, as defined in [9]. A straightforward calculation show that this definition is coherent with the open covering $\{\Lambda_0(L_1)\}_{L_1 \in \Lambda}$, that is, if γ has image in $\Lambda_0(L_1) \cap \Lambda_0(L_2)$, then $\text{sf}(\varphi_{L_0, L_1} \circ \gamma) = \text{sf}(\varphi_{L_0, L_2} \circ \gamma)$. The integer $\mu_{L_0}(\gamma)$ is additive by concatenation and invariant by a homotopy fixing the endpoints. Given any curve $\gamma : [0, 1] \rightarrow \mathcal{F}_{L_0}(\Lambda)$, we obtain a partition $0 = x_0 < \dots < x_n = 1$ and $L_1, \dots, L_n \in \Lambda$ such that (L_0, L_j) are complementary and $\gamma_j = \gamma|_{[x_{j-1}, x_j]}$ has image in $\Lambda_0(L_j)$, for $j = 1, \dots, n$. We define the Maslov index of γ with respect to L_0 as

$$\mu_{L_0}(\gamma) = \sum_{j=1}^n \text{sf}(\varphi_{L_0, L_j} \circ \gamma_j).$$

Homotopy invariance and additivity properties show that the Maslov index is well-defined and gives a homomorphism of the fundamental grupoid¹ of $\mathcal{F}_{L_0}(\Lambda)$ with \mathbb{Z} . For more details about the construction of the Maslov index, see [3], [4], [6]. The next proposition shows that the restriction of μ_{L_0} to the fundamental group of $\mathcal{F}_{L_0}(\Lambda)$ is a isomorphism.

PROPOSITION 2.8 *The Maslov index with respect to a fixed lagrangian L_0 is a isomorphism $\mu_{L_0} : \pi_1(\mathcal{F}_{L_0}(\Lambda)) \rightarrow \mathbb{Z}$.*

PROOF. Note that by theorem 2.7, $\mathcal{F}_{L_0}(\Lambda)$ is path-connected and $\pi_1(\mathcal{F}_{L_0}(\Lambda)) = \mathbb{Z}$, so it is sufficient to prove that μ_{L_0} is onto. Given $n \in \mathbb{Z}$, take a loop α in $\mathcal{F}^{\text{sa}}(L_0^\perp)$ with spectral flow n ; the curve $\gamma = \varphi_{L_0^\perp, L_0}^{-1} \circ \alpha$ is a loop in $\mathcal{F}_{L_0}(\Lambda)$ with Maslov index n , as desired. ■

References

- [1] ABBONDANDOLO, A., MAJER, P., *Infinite Dimensional Grassmannians*, arXiv: math.AT/0307192.
- [2] BOOSS-BAVNBEK, B., FURUTANI, K., *Symplectic Functional Analysis and Spectral Invariants*, Contemporary Mathematics 242, (1999), 53-83.
- [3] EIDAM, J. C. C., PICCIONE, P., *Partial Signatures and the Yoshida-Nicolaescu Theorem*, arXiv:math.DG/0502459.

¹The fundamental grupoid of a space X is the set of all homotopy classes, with fixed endpoints, of curves in X .

- [4] FURUTANI, K., *Fredholm-Lagrangian-Grassmannian and the Maslov Index*, J. Geom. Phys. 51 (2004), 269-331.
- [5] KATO, T., *Perturbation Theory for Linear Operators*, Springer-Verlag, 1976.
- [6] NICOLAESCU, L., *The Maslov Index, the Spectral Flow and Decompositions of Manifolds*, Duke Math. Journal 80 (1995), 485-533.
- [7] PALAIS, R., *On the Homotopy Type of certain Groups of Operators*, Topology 3 (1965), 271-279.
- [8] PALAIS, R., *Homotopy Theory in Infinite Dimensional Manifolds*, Topology 5 (1966), 1-16.
- [9] PHILLIPS, J., *Self-adjoint Fredholm Operators and Spectral Flow*, Canad. Math. Bull. 39 (4) (1996), 460-467
- [10] RIESZ, F., SZ.-NAGY, B., *Functional analysis*, Frederick Ungar Publishing Co., New York, 1955.

DEPARTAMENTO DE MATEMÁTICA,
 INSTITUTO DE MATEMÁTICA E ESTATÍSTICA
 UNIVERSIDADE DE SÃO PAULO, BRASIL.
 E-MAIL ADDRESSES: zieca@ime.usp.br, piccione@ime.usp.br