

Ground State Rotational Bands of Deformed $e-e$ Nuclei

Mohamed E. Kelabi^{*}
A. Y. Ahmed^{*}
Vikram Singh[†]

Abstract

In the frame work of the hydrodynamical model, a new model of the ground state rotational bands of deformed $e-e$ nuclei is developed by introducing the variable moment of inertia, and the effect of β - and γ - vibrational bands. The model is applied to calculate the energies of the ground state band of ^{158}Dy . The results of our calculations are in close agreement with data compared with other existing models.

Introduction

Recently Abzouzy and Antony^[1] have shown that the semi-empirical expression of Sood^[2] works well for Th and U nuclei, which is obtained from the power series of angular momentum factor $I(I+1)$ as

$$\frac{C}{B} = \frac{D}{C} = \dots = N \frac{B}{A} \quad (1)$$

where A, B, C, D, \dots are the coefficients of successive order correction terms^[3]. They have then assigned N to depend on two parameters

$$N = a - bI \quad (2)$$

and deduced their values^[1]: $a = 2.55$ and $b = 0.05$ for the best fit.

Theory and formalism

A more systematic calculations can be carried out by considering only the Rotational-Vibrational model RVM of Bohr and Mottelson^[4]. This was done by adding the effect of nuclear rotation^[2] on the moment of inertia, thus one writes

$$E(J) = A(J)J(J+1) - B(J)J^2(J+1)^2 \quad (3)$$

where

$$A(J) = \frac{\hbar^2}{2J(J)} \quad (4)$$

^{*} Physics Department, Al-Fateh University, Tripoli, LIBYA

[†] B-1/1051, Vasant Kunj, New Delhi 110070.

$$B(J) = \frac{4A(J)^3}{(\hbar\omega_\gamma)^2} + \frac{12A(J)^3}{(\hbar\omega_\beta)^2} \quad (5)$$

here $\hbar\omega_\beta$ and $\hbar\omega_\gamma$ are the head energies of the β - and γ -vibrations^[5], respectively. For convenience, we rewrite Eq. (3) as

$$E(J) = A(J)J(J+1) \left(1 - \frac{B(J)}{A(J)} J(J+1) \right) \quad (6)$$

where the ratio $\frac{B(J)}{A(J)}$ takes the form

$$\frac{B(J)}{A(J)} = \frac{\hbar^4}{J(J)^2} \left[\frac{1}{(\hbar\omega_\gamma)^2} + \frac{3}{(\hbar\omega_\beta)^2} \right]. \quad (7)$$

In order to write $J(J)^{-1}$ and $J(J)^{-2}$ in terms of the ground state value J_0 ^[6], which corresponds to $J=0$, we make Taylor series expansion, giving

$$\frac{1}{J(J)} = \frac{1}{J_0} \frac{1}{1 + \sigma_1 J + \sigma_2 J^2 + \dots} \quad (8)$$

$$\frac{1}{J(J)^2} = \frac{1}{J_0^2} (1 - 2\sigma_1 J - 2\sigma_2 J^2 + 3\sigma_1^2 J^2 + 6\sigma_1\sigma_2 J^3 - \dots) \quad (9)$$

where, we have introduced the Softness parameter^[7], defined by

$$\sigma_n = \frac{1}{n!} \frac{1}{J_0} \left. \frac{\partial^n J(J)}{\partial J^n} \right|_{J=0}. \quad (10)$$

Keeping only the first order terms in σ_n , Eqs. (8) and (9) become

$$\frac{1}{J(J)} = \frac{1}{J_0} \frac{1}{1 + \sigma_1 J} \quad (11)$$

$$\frac{1}{J(J)^2} = \frac{1}{J_0^2} (1 - 2\sigma_1 J). \quad (12)$$

Inserting Eqs. (11) and (12) into (4) and (7), respectively, and rearranging Eq. (6), one easily obtains

$$E(J) = \frac{A_0}{1 + \sigma_1 J} J(J+1) \left[1 - \frac{B_0}{A_0} (1 - 2\sigma_1 J) J(J+1) \right] \quad (13)$$

which can be simplified to give

$$E(J) = \frac{A_0}{1 + \sigma_1 J} J(J+1) - c A_0^3 \frac{(1 - 2\sigma_1 J)}{1 + \sigma_1 J} J^2(J+1)^2 \quad (14)$$

where, $A_0 = \frac{\hbar^2}{2\mathcal{J}_0}$ and $B_0 = cA_0^3 = \left[\frac{4}{(\hbar\omega_\gamma)^2} + \frac{12}{(\hbar\omega_\beta)^2} \right] A_0^3$.

This third order nonlinear expression, Eq. (14), can be used to calculate the ground state rotational bands of deformed $e-e$ nuclei.

Results

We have applied Eq. (14) to calculate the energies of the ground state band of ^{158}Dy . In these calculations, we have treated A_0 , σ_1 and c as three free parameters, which can be determined by fitting the first excited states 2^+ , 4^+ , and 6^+ with the available data. In Fig. 1 we schematically present the results of our calculations along with the experimentally observed energies and the prediction of other existing models^[8]. The results of our model are in close agreement with data than the other models.

$$A_0 = 17.042 \text{ KeV} \quad \sigma_1 = 13.806 \quad c = 3.440 \times 10^{-6} \text{ KeV}^{-2}$$

| J^π | Exp | VMI | VAVM | GVM | Present work |
|------------|--------|--------|--------|--------|--------------|
| 18^+ ——— | 3781.7 | 3943.0 | 3913.3 | 4030.3 | 3866.3 |
| 16^+ ——— | 3190.7 | 3263.8 | 3248.5 | 3329.3 | 3221.0 |
| 14^+ ——— | 2612.6 | 2628.6 | 2623.7 | 2674.9 | 2613.3 |
| 12^+ ——— | 2049.2 | 2041.7 | 2043.5 | 2071.9 | 2043.1 |
| 10^+ ——— | 1519.9 | 1508.8 | 1513.5 | 1526.2 | 1516.2 |
| 8^+ ——— | 1044.1 | 1036.9 | 1041.4 | 1045.0 | 1043.1 |
| 6^+ ——— | 637.87 | 637.87 | 637.87 | 637.87 | 637.87 |
| 4^+ ——— | 317.26 | 317.26 | 317.26 | 317.26 | 317.26 |
| 2^+ ——— | 98.94 | 98.94 | 98.94 | 98.94 | 98.94 |
| 0^+ ——— | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |

Fig.1. Experimental and calculated energy levels of ^{158}Dy in [KeV].

Conclusion

The present model Eq. (14) is practically fit to predict the ground state rotational bands of almost all deformed $e-e$ nuclei, and can also be applied to nuclei where the energies of β - and γ - vibrations are experimentally available. With the known values of $\hbar\omega_\beta$ and $\hbar\omega_\gamma$, contained in parameter c of Eq. (14), the model reduces to only two parametric expression, hence, the higher spin states can be evaluated by fitting only the first excited states 2^+ and 4^+ with experimental data.

References

- 1) A. Abzouzi and M. S. Antony, Phys. Rev. C37 (1988) 401.
- 2) P. C. Sood, Phys. Rev. 161 (1967) 1063.
- 3) O. Nathan and S. G. Nilson, Alpha, Beta, and Gamma Ray Spectroscopy, Ed. K. Siegbahn, North Holland (1965) 601.
- 4) A. Bohr and B. R. Mottelson, Nuclear Structure Vol-11, Nuclear Deformation, Benjamin, New York (1984).
- 5) R. K. Sheline, Rev. Mod. Phys. 32 (1960) 1.
- 6) J. S. Batra and R. K. Gupta, Phys. Rev. C43 (1991) 43.
- 7) H. Morinaga, Nucl. Phys. 75 (1966) 385.
- 8) Dennis Banatsos and Abraham Klein, Atomic Data and Nuclear Data Tables 30 (1984) 271.