

Three Years of Mira Variable Photometry: What Has Been Learned?

Dale E. Mais
Palomar Community College
dmais@ligand.com

David Richards
Aberdeen & District Astronomical Society
david@richweb.f9.co.uk
&
Robert E. Stencel
Dept. Physics & Astronomy
University of Denver
rstencdel@du.edu

Abstract

The subject of micro-variability among Mira stars has received increased attention since DeLaverny et al. (1998) reported short-term brightness variations in 15 percent of the 250 Mira or Long Period Variable stars surveyed using the broadband 340 to 890 nm “Hp” filter on the HIPPARCOS satellite. The abrupt variations reported ranged 0.2 to 1.1 magnitudes, on time-scales between 2 to 100 hours, with a preponderance found nearer Mira minimum light phases. However, the HIPPARCOS sampling frequency was extremely sparse and required confirmation because of potentially important atmospheric dynamics and dust-formation physics that could be revealed. We report on Mira light curve sub-structure based on new CCD V and R band data, augmenting the known light curves of Hipparcos-selected long period variables [LPVs], and interpret same in terms of [1] interior structure, [2] atmospheric structure change, and/or [3] formation of circumstellar [CS] structure. We propose that the alleged micro-variability among Miras is largely undersampled, transient overtone pulsation structure in the light curves. © 2005 Society for Astronomical Science.

1. Introduction

From European Space Agency's High Precision Parallax Collecting Satellite, HIPPARCOS (ESA, 1997) mission data, deLaverny et al. (1998) discovered a subset of variables (15 percent of the 250 Mira-type variables surveyed) that have exhibited abrupt short-term photometric fluctuations, within their long period cycle. All observations were made in a broadband mode, 340 to 890 nm, their so-called Hp magnitude. They reported variation in magnitude of 0.23 to 1.11 with durations of 2 hours up to almost 6 days, preferentially around minimum light phases. Instrumental causes could not be identified to produce this behavior. Most of these variations are below the level of precision possible with purely visual estimates of the sort collected by AAVSO, but may contribute to some of the scatter in visual light curves. 51 events in 39 M-type Miras were

detected with HIPPARCOS, with no similar variations found for S and C-type Miras. These short-term variations were mostly detected when the star was fainter than Hp = 10 magnitude including one star at Hp = 13 magnitude and one at Hp = 8.3. For 27 of the original 39 observations, the star underwent a sudden increase in brightness. From their study, deLaverny et al. found that 85% of these short-term variations were occurring around the minimum of brightness and during the rise to the maximum, at phases ranging from 0.4 to 0.9. No correlation was found between these phases and the period of the Miras, but that brightness variations do occur preferentially at spectral types later than M6 and almost never for spectral types earlier than M4. Similar results were reported by Maffei & Tosti (1995) in a photographic study of long period variables in M16 and M17, where 28 variations of 0.5 mag or more on timescales of days were found among spectral types later than M6. Schaeffer (1991) collected reports on fourteen cases of flares

on Mira type stars, with an amplitude of over half a magnitude, a rise time of minutes, and a duration of tens of minutes. In analogy to the R CrB phenomenon, brightness variation could also be consequence of dust formation (fading) and dissipation (brightening) in front of a star's visible hemisphere. Future narrow band infrared interferometric observations will help resolve this.

Recently, Wozniak, McGowen and Vestrand (2004) reported analysis of 105,425 I-band measurements of 485 Mira-type galactic bulge variables sampled every other day, on average, over nearly 3 years as a subset of the OGLE project. They failed to find any significant evidence for micro-variability, to a limit of 0.038 I-band events per star per year. They conclude that either Hipparcos data are instrumentally challenged, or that discovery is subject to metallicity or wavelength factors that minimize detection in I-band among galactic bulge objects. In contrast, Mighell and Roederer [2004] report flickering among red giant stars in the Ursa Minor dwarf spheroidal galaxy, including detection of low-amplitude variability in faint RGB stars on 10 minute timescales! However, Melikian [1999] provides a careful analysis of the light curves for 223 Miras based on Hipparcos data, finding that 82 stars [37%] show a post-minimum hump-shaped increase in brightness on the ascending branch of the light curve. Melikian advocates that differing physical processes and perhaps stellar properties – e.g. later spectral types, longer periods and higher luminosity - differentiate behaviors among these stars.

The purpose of this report is to provide V and R-band photometry of objects related to the deLaverny et al. results, with dense temporal sampling. We find a similar lack of micro-variability as noted by Wozniak et al., but do confirm facets of the Melikian report. This suggests that these phenomenon can be placed in a larger context of pulsational variations and episodic dust formation, with implications for ongoing spectroscopic and interferometric observations of mid-infrared studies of LPV stars.

2. Observations and Data Reduction

Our target list primarily was drawn from the objects listed by deLaverny et al. (1998), although limited to the northern sky. Of the 39 M type Mira's described therein, 20 are relatively bright and visible from the northern hemisphere. Because of the efficiency of automated sampling, we

augmented this list with additional M type Miras and the brightest C, CS and S stars where one can obtain good signal to noise with low to moderate resolution spectroscopy on a small telescope. These stars and associated characteristics are detailed in Table 1.




Figure 1. Dale Mais and his photometry machine co-located near Valley Center, California.

These along with a variety of brighter S and C type stars were also chosen. Brighter stars were chosen since they represented stars with magnitudes such that moderate resolution spectroscopy could be performed as part of the monitoring process. To accomplish this in a semi-automated manner, the telescope, camera and filter wheel are controlled by a single computer using Orchestrate software (www.bisque.com). Once the images are reduced, a script written by one of the authors (David Richards) examines the images performing an image link with TheSky software (www.bisque.com). The images obtained in this manner are stamped both with the name of the variable star, since this was how Orchestrate was instructed to find the object, and the position of the image in the sky. This allows TheSky to quickly perform the links with its USNO database. Once the astrometric solution is accomplished, the program reads through a reference file with the pertinent data such as reference star name and magnitudes along with variable star of interest. The input file is highly flexible, stars and filter magnitudes of reference stars can be added freely as image data require. This file only needs to be created once, especially convenient for a set of program stars, which will have continuous coverage over time. There is no need for entering magnitude information of reference stars in a repeated manner. The results file is readily imported to spreadsheet software, where the various stars and their magnitudes can be plotted, almost in real time. This is an important aspect of this project, the ability to see changes

(flare-ups) quickly and as a result respond to these changes with spectroscopic observations.


(a) XZ Herculis



(b) S Serpentis

Figures 2: Representative light curves. Additional examples can be found in Mais et al. 2005 SAS Conference Proceedings. These data will be submitted to the AAVSO archives.

Photometry was conducted with an Astrophysics 5.1-inch f/6 refractor located in rural San Diego county, California, using an ST-10XME camera and 2x2 binned pixels and the Johnson V and R filters. Images were obtained in duplicate for each band and two reference stars used per variable star for analysis. Image reduction was carried out with CCDSOFT (www.bisque.com) and Source



Extractor (Bertin and Arnout, 1996) image reduction groups and specially written scripts for magnitude determinations, which allowed for rapid, nearly real time magnitudes to be found (see below).

The project has been underway since 2003 and involves a total of 96 stars, 20 M type Miras, 19 S types and the remainder C types. While there are certainly many more of these type stars, only those that had a significant part of their light curve brighter than visual magnitude 8 were considered, due to magnitude limitations in the spectroscopy part of the project. Fortunately, these stars are much brighter in the R and I bands, often by 2-4 magnitudes when compared to their V magnitudes, and many of the interesting molecular features are found in this region of the spectrum. The photometric analysis involves using 2 different reference stars. Their constant nature is readily discerned over the time period by the horizontal slope of their light curves, both in the V and R bands. After considerable effort, magnitudes are now determined at the 0.02 magnitude level. Thus any flare-ups in the range of 0.1 magnitude and brighter should be readily discerned.

Early on it was felt that semi-automating the process was the best way to proceed. The use of a precision, computer-controlled mount (Paramount, www.bisque.com) along with the suite of software by Software Bisque got the project rapidly underway. TheSky in conjunction with CCDSOFT lends itself to scripting, and a script was put together that automated the magnitude determinations. To give an example of how this has streamlined the effort, on a typical night, initially using Orchestrate and later using a script developed by coauthor David Richards to control the telescope, camera and filter wheel, 40 stars, visible at the time, are imaged in duplicate in each of the V and R bands. This takes about 1 hour. Reduction of the images using image reduction groups in CCDSOFT takes another 5 minutes. The script that determines the magnitudes takes about 10 minutes to churn its way through all the images. Within another 20 minutes, the data, via spreadsheet, has been added to each variable stars growing light curve. Thus in less than 2 hours all of the program stars have been observed and their results tallied. Until more of program stars rotate into view, one is free to pursue spectroscopic examination of the program stars, establishing baseline observations. Another portion of this effort included standardizing the reference stars in each of the fields using the Landolt standards (Landolt, 1983). Once this is done all previous and subsequent observations of the variable stars will have their magnitudes expressed in absolute Johnson-Cousins magnitudes.

3. Results & Analysis

3a. Evidence for flares?

Three years of monitoring of 96 Mira-like variables, including all the northern objects included




Figure 3a. variation near minimum light, RR Boo

in the Hipparcos report [deLaverny et al. 1998] has yielded minimal evidence for flare-like changes in V or R band. Best cases include 0.2 and 0.3 magnitude increases near minimum light in RR Boo. This object was reported by Guenther and Henson [2001] to have shown a one-time 0.8 mag flare. Other marginal cases in our data include CE Lyr, X CrB and DH Lac. In the case of CE Lyr, the proximity of a faint stellar companion could contribute to jumps in automated photometry as the variable changes around minimum. Otherwise, most variables show smooth light variation with no hint of flare-like fluctuations at a level of 0.01 mag.

3b. Light curve “bump” phenomena

In contrast to high-frequency events like flares [hours or days], examining the light curves for the stars observed between 2003 and 2006 revealed persistent low frequency changes on timescales of weeks. Following the discussion of these by Melikian [1999], we label these “bumps” in the Mira light curves. Good examples of this are seen in the light curves of RT Boo, R CMi, X CrB, U Cyg, XZ Her [Fig.2a], S Ser [Fig.2b], RU Her, U Cyg and R Lyn. Some of these are seen in visual light curves compiled by AAVSO and AEFOV, but others are seen at levels below the ~0.1 mag precision typical of visual observations. This is one

of the important benefits of high precision photometry. Most light curve bumps are non-recurring and seem to appear after especially deep minimum light. The correlation of bumps with Mira properties deserves further attention. A few extremes in our sample are noted: double maxima in T Cam, S Cas, RR Her, S Cep, RS Cyg, RR Her and Y Per. Two stars show the bump feature post-maximum light: V CrB and T Dra.

3c. Period determination

Period finding was performed using PerAnSo software suite by Tonny Vanmunster, [http://users.skynet.be/fa079980/peranso/index.htm] which reports best fits using the ANOVA method. Initial application shows agreement with literature periods for most variables, within a few percent and with errors in fit of 1 to 20 days, depending on the data interval and phase coverage thus far. More work is in progress.




Figure 3b: correlation between M-type Miras newly derived periods and older GCVS determinations.




Figure 3c: correlation between C&S-type Miras newly derived periods and older GCVS determinations.

4. Conclusions

Among our conclusions based on V and R band measurements, with ~ 10 millimag precision, of nearly one hundred brighter Mira type stars are:

[1] flare events are rare, and statistically similar to the OGLE result for I band monitoring of 0.038 events per star per year, with some evidence that “flares” are bluer in color;

[2] we are confirming indications of correlations between depth of minima and occurrence of a “bump” or change of slope on the ascending branch of some light curves [cf. Melikian 1999];

[3] our coverage of approximately 3 cycles is sufficient to confirm the majority of previously published periods;

[4] we hypothesize that bump phasing and contrast varies with internal structure and opacity in analogy with similar phenomena among the “bump Cepheids” and deserves further study.

5. Acknowledgements

The authors wish to acknowledge the assistance of Thomas Bisque of Software Bisque for many useful discussions and help with the scripting, and of the AAVSO for its compilation of variable star visual light curves referenced in this paper, and the estate of William Herschel Womble for partial support of University of Denver astronomers participating in this effort.

6. References

- Buchler, J. R., Moskalik, P., & Kova'cs, G. 1990, ApJ, 351, 617 “A survey of Cepheid bump mode pulsations”
- de Laverny, P.; Geoffray, H.; Jorda, L.; Kopp, M. 1997 A&AS 122: 415 “Long-term UBVRI monitoring of 12 southern hemisphere Long Period Variables”
- deLaverny, P., Mennessier, M., Mignard, F., & Mattei, J. 1998, A & A 330: 169, “Detection of short-term variations in Mira-type variables from HIPPARCOS photometry”
- Guenther, H. D.; Henson, G. D., 2001, BAAS 199: 90-07 “Monitoring Select Mira Stars for Short-term Variability”
- Hawkins, G., Mattei, J. & Foster, G. 2001 PASP 113:501, “R Centauri: An Unusual Mira Variable in a He-Shell Flash”
- Maffei, P. and Tosti, G. 1995 AJ 109: 2652, “Rapid Variations in Long Period Variables”
- Marsikova, V. 1997 JAAVSO 27: 141, “Fast Variations of the Mean brightness and Light Curve Parameters of the Carbon Mira-Type Star S Cep”
- Melikian, N. 1999 Astrophysics 42: 408, “Classification of light curves of Mira variables”
- Mighell, K. and Roederer, I. 2004 ApJ 617: L41 “Flickering Red Giants in the Ursa Minor Dwarf Spheroidal Galaxy: Detection of Low-Amplitude Variability in Faint Red Giant Branch Stars on 10 Minute Timescales” & San Diego AAS poster 54.18
- Schaefer, B. 1991, Ap.J. 366: L39, “Flares on Mira stars?”
- Stencel, R., Phillips, A., Jurgenson, C. and Ostrowski-Fukuda, T. 2002 DC-AAS poster 92.19 “Efforts to Verify Microvariability among Hipparcos-selected AGB Stars”
- Teets, W. and Henson, G. 2004 Atlanta-AAS poster 8.03 “Multiple Wavelength Monitoring of Mira Stars for microvariability”
- Vardya, M. 1988 A&AS 73:181, “Classification of Mira variables based on visual light curve shape”
- Wozniak, P., McGowan, K. and Vestrand, W. 2004 ApJ 610:1038 “Limits on I-band Microvariability of Galactic Bulge Mira variables” [< 0.038 I-band events per star per year]

Tables

Table 1a: M type Mira Stars monitored

Star*	Sp.Type, Vmax	Period, Epoch* GCVS IV	Period, Epoch* Anova/Peranso	Notes
SV And**	M5e, 7.7	316.21, 42887	308.64, 52892.3	
R Boo	M3e, 6.2	223.40, 44518	224.22, 53465.3	
RR Boo**	M2e, 8.3	194.70, 43047	194.74, 53197.7	
RT Boo**	M6.5e, 8.3	273.86, 42722	276.12, 53225.5	
W Cnc	M6.5e, 7.4	393.22, 43896	406.36, 53336.8	
R CVn	M5.5e, 6.5	328.53, 43586	335.14, 53425.5	
V CVn	M4e, 6.5	191.89, 43929	189.21, 53210.1	
R Cas	M6e, 4.7	430.46, 44463	385.8, 53533.9	dP/dt?
T Cas	M6e, 6.9	444.83, 44160	447.18, 53289.3	
V Cas	M5e, 6.9	228.83, 44605	232.71, 53285.6	
T Cep	M5.5e, 5.2	388.14, 44177	372.67, 53287.2	
R Cet**	M4e, 7.2	166.24, 43768	163.30, 53221.6	
X CrB**	M5e, 8.5	241.17, 43719	240.17, 52865.3	
AM Cyg**	M6e, 11.3	370.6, 30075	374.47, 53345.2	
T Eri**	M3e, 7.2	252.29, 42079	--, --	
RU Her**	M6e, 6.8	484.83, 44899	--, --	
SS Her**	M0e, 8.5	107.36, 45209	106.44, 53245.4	
XZ Her**	M0, 10.5	171.69, 33887	171.15, 52939.3	
R Hya	M6eTc, 3.5	388.87, 43596	--, --	
T Hya**	M3e, 6.7	298.7, 41975	292.44, 53492.1	
X Hya**	M7e, 7.2	301.10, 41060	287.91, 53113.4	dP/dt?
DH Lac**	M5e, 11.6	288.8, 41221	292.45, 52929.6	
R Lmi	M6.5eTc:, 6.3	372.19, 45094	390.02, --	
W Lyr	M2e, 7.3	197.88, 45084	201.57, 53575.0	
CE Lyr**	--, 11.7	318, 25772	315.90, 53259.0	
HO Lyr**	M2e, 11.4	100.4, 30584	99.75, 53235.6	
V Mon**	M5e, 6.0	340.5, 44972	342.51, 53312.7	
RX Mon**	M6e, 9.6	345.7, 35800	351.62, 53192.9	
Z Oph	K3ep, 7.6	348.7, 42238	347.52, 53450.6	
R Peg	M6e, 6.9	378.1, 42444	358.17, 53350.0	dP/dt?
RT Peg	M3e, 9.4	215.0, 45599	216.76, 53315.5	
SW Peg	M4e, 8.0	396.3, 38750	398.43, 53482.9	
S Per	M3lab, 7.9	822, --	775.33, 53058.8	
S Ser**	M5e, 7.0	371.84, 45433	371.69, 53234.1	
AH Ser**	M2, 10.0	283.5, 36682	284.83, 53413.8	

*Stars are ordered by the traditional nomenclature: alphabetical by constellation, then by single letter R...Z, then double letters RR...ZZ, then AA...QQ and finally V###; MJD= JD – 2,400,000; GCVS: Gen.Catalog Var.Stars 4th ed., 1985 & <http://www.sai.msu.su/groups/cluster/gcvs/gcvs/>

** Hipparcos “flare stars” included in deLaverny et al. [1998]

Table 1b: C & S type Mira Stars monitored

Star* Sp.Type,	Period, Epoch* V [max]	Period, Epoch* GCVS IV	Period, Epoch* Anova/Peranso	Notes
W And	S6.1e, 6.7	395.93, 43504	406.72, 53221.6	
RR And	S6.5,2e, 8.4	328.15, 43390	327.17, 52917.8	
ST And	C4.3e, 7.7	328.34, 38976	324.59, 53424.4	
SU And	C6.4N8,8.0	--, --	282.83, 53235.0	dP/dt?
V Aql	C5.4N6,6.6	353, --	524.43, 53608.2	dP/dt?
W Aql	S3.9e, 7.3	490.43, 39116	484.53, 53157.9	
UV Aql	C5.4N4,11.1	385.5, 30906	--, --	
S Aur	C4-5N3,8.2	590.1, 42000	554.59, 53169.9	dP/dt?
V Aur	C6.2eN3e,8.5	408.09, 43579	356.18, 53287.1	dP/dt?
TX Aur	C5.4N3, 8.5	--, --	163.30, 53370.8	
EL Aur	C5.4N3,11.5	--, --	234.60, 53064.8	
FU Aur	C7.2N0,11.0	--, --	157.04, 53265.2	
R Cam	S2.8e, 6.97	270.22, 43978	270.53, 53153.5	
S Cam	C7.3eR8,7.7	327.26, 43360	331.74, 53221.6	
T Cam	S4.7e, 7.3	373.20, 43433	370.74, 53160.7	
U Cam	C3.9N5, 11.0	--, 43060	--, --	
RU Cam	C0.1K0, 8.1	22, --	313.48, --	
ST Cam	C5.4N5, 9.2	300:, --	160.30, 53214.9	
UV Cam	C5.3R8, 7.5	294:, --	--, --	
W Cma	C6.3N, 6.35	--, --	295.53, --	
R Cmi	C7.1eJ, 7.25	337.78, 41323	335.77, 53114.8	
T Cnc	C3.8R6, 7.6	482, --	495.05, 53396.9	
V Cnc	S0e, 7.5 272.13,	43485	266.19, 53284.3	
S Cas	S3.4e, 7.9	612.43, 43870	636.07, 53059.6	
U Cas	S3.5e, 8.0	277.19, 44621	280.90, 52913.8	
W Cas	C7.1e, 7.8	405.57, 44209	404.94, 53143.7	
ST Cas	C4.4N3, 11.6	--, --	408.53, 53307.0	
V365 Cas	M5S7.2,10.2	136, --	325.73, 52844.8	
S Cep	C7.4eN8,7.4	486.84, 43787	488.08, 52976.0	
V CrB	C6.2eN2, 6.9	357.63, 43763	351.86, 53449.8	
R Cyg	S2.5eTc, 6.1	426.45, 44595	425.24, 53550.6	
U Cyg	C7.2eNp,5.9	463.24, 44558	488.61, 52919.2	dP/dt?
V Cyg	C5.3eNp, 7.7	421.27, 44038	403.23, 53248.1	
RS Cyg	C8.2eN0p,6.5	417.39, 38300	419.58, 53254.4	
RV Cyg	C6.4eN5,10.8	263, --	458.72, 53592.6	
RY Cyg	C4.8N, 8.5	--, --	307.22, 53233.4	
SV Cyg	C5.5N3, 11.7	--, --	415.47, 53164.7	
TT Cyg	C5.4eN3,10.2	118, --	129.03, 53255.9	
YY Cyg	C6.0evN,12.1	388, 298261	384.62, 52941.8	
AW Cyg	C4.5N3,11.0	--, --	557.88, 52875.9	
AX Cyg	C4.5N6, 7.85	--, --	461.75, 52845.1	
V460 Cyg	C6.4N1, 5.6	180:, --	158.31, 53183.8	
T Dra	C6.2eN0, 7.2	421.6, 43957	410.17, 53283.2	
UX Dra	C7.3N0, 5.9	168, --	314.27, 53246.2	
R Gem	S2.9eTc, 6.0	369.91, 43325	374.13, 53359.8	
T Gem	S1.5e, 8.0	287.79, 44710	289.15, 53153.5	
TU Gem	C6.4N3, 9.4	230, --	431.03, 53100.2	
NQ Gem	C6.2R9ev,7.4	70:, --	256.72, --	
S Her	M4Se, 6.4	307.28, 45054	307.33, 53638.5	
RR Her	C5.7eN0, 8.8	239.7, --	233.47, 53126.3	
R Lep	C7.6eN6, 5.5	427.01, 42506	456.19, 53009.3	

SZ Lep	C7.3R8, 7.4	--, --	--, --
R Lyn	S2.5e, 7.2	378.75, 45175	382.52, 53124.2
T Lyr	C6.5R6, 7.8	--, --	430.79, 52892.2
U Lyr	C4.5eN0, 8.3	451.72, 42492	--, --
HK Lyr	C6.4N4, 7.8	--, --	354.11, 53182.0
V614 Mon	C4.5JR5, 7.0	60:, --	--, --
V Oph	C5.2N3e, 7.3	297.21, 45071	302.58, 52826.6
TW Oph	C5.5N, 11.6	185:, --	459.35, --
RT Ori	C6.4Nb, 9.7	321, --	--, --
BL Ori	C6.3NbTc, 7.9	--, --	316.01, 53036.8
RX Peg	C4.4JN3, 9.7	629:, --	495.24, 52987.9
RZ Peg	C9.1eNTc, 7.6	438.7, 54248	433.93, 53096.8
HR Peg	S5.1M4, 6.1	50:, --	--, --
Y Per	C4.3eR4, 8.1	248.6, 45245	255.81, 53083.3
V466 Per	C5.5N5, 10.9	--, --	358.8, 53288.8
T Sgr	S4.5e, 7.1	394.66, 44897	382.65, 53049.7
ST Sgr	S4.3e, 7.2	395.12, 40463	401.07, 53046.6
AQ Sgr	C7.4N3, 9.1	199.6, --	--, --
V1942 Sgr	C6.4N2R8, 6.7	--, --	209.68, 52879.6
FO Ser	C4.5R6, 8.5	--, --	--, --
TT Tau	C4.2N3, 10.2	166.5, --	163.30, 53010.9
SS Vir	C6.3eN, 6.0	364.14, 45361	361.85, 53231.7
BD Vul	C6-7Ne, 9.3	430, 25758	--, --

*Stars are ordered by the traditional nomenclature: alphabetical by constellation, then by single letter R...Z, then double letters RR...ZZ, then AA...QQ and finally V###; MJD= JD – 2,400,000; GCVS: Gen.Catalog Var.Stars 4th ed., 1985 & <http://www.sai.msu.su/groups/cluster/gcvs/gcvs/>