

CRYPTOPLANET UPDATE

Robert L. Kurucz

Harvard-Smithsonian Center for Astrophysics

February 15, 2007

Abstract

We have had several talks recently reviewing 11 years of exoplanet discoveries through radial velocity variations, or from transits, or from microlensing. More than 200 exoplanets have been found, including some around pulsars that we do not discuss here.

My physical definition for a planet is a roughly spherical, self-gravitating body more massive than 10^{26} g formed from the leftover material in a protostellar disk after the protostar forms. Radiation from the protostar pushes the inner wall of the disk outward. The material agglomerates and forms planets in radial sequence. The outer planets are formed slowly by classical dynamical mechanisms acting in the snow zone. Planets have dense cores because of agglomeration. Planets bound around other stars were formed and behave like the planets in our solar system.

Not one of the exoplanets discovered thus far is a planet. They are cryptoplanets formed from matter ejected by protostars. When protostars have excessive infall at high latitudes, they partially balance angular momentum through outflow at the equator as they spin up. The ejected matter is trapped in the magnetic torus formed between the star and the disk, like a tokamak. The tokamak eventually reconnects and magnetic compression forms self-gravitating remnants trapped and compressed by a closed spherical magnetic field, spheromaks. Cooled spheromaks are cryptoplanets. They orbit near the star. They can merge with each other or fall into the star or be ejected. They can grow by accreting gas. They have a low density core and abundances characteristic of the protostar. Their masses, radii, densities, and orbits are random, and are inconsistent with the parameters for planets. They tend to have lower density than planets.

More than 20 disks around protostars have been observed and are claimed to be pre-planetary. In fact they are post-planetary. Disks are not observable until the central Jupiter-orbit-size region has been cleared out and the inner planets have been formed out through “Jupiter”.

CRYPTOPLANET UPDATE

ROBERT L. KURUCZ

**HARVARD-SMITHSONIAN
CENTER FOR ASTROPHYSICS**

FEBRUARY 15, 2007

CRYPTOPLANET UPDATE

CONSERVATION OF ANGULAR MOMENTUM

CONVECTION

UNIFORMITARIANISM

FORMATION OF PLANETS

FORMATION OF CRYPTOPLANETS

CRYPTOPLANET CHARACTERISTICS

CONSERVATION OF ANGULAR MOMENTUM

SHU (1982)




Figure 3.23. The spin-up of an ice-skater. When the twirling skater has her arms out, her moment of inertia I is large and her rate of spin ω is small for a given angular momentum $J = I\omega$. When she pulls her arms in, her moment of inertia I decreases, and the conservation of angular momentum implies that her rate of spin ω increases.

**HOWEVER, THIS CAPTION DOES NOT
EXPLAIN THE FIGURE**

MASSES:

SKATER 50 KG, HAIR 50G, SKIRT 500G

1/1000 OF MASS MOVES OUTWARD IN HAIR

1/100 OF MASS MOVES OUTWARD IN SKIRT

**ADD 1 KG LEAD DRAPERY WEIGHTS TO HEM
OF SKIRT. NOW 3/100 OF MASS.**

ENOUGH TO REDUCE SPINUP


ADD TO CAPTION:

**A ROTATING BODY TRIES TO MINIMIZE
ITS SPINUP BY MAXIMIZING ITS MOMENT
OF INERTIA**




CONVECTION

SURFACE




PLUME CONVECTION

SURFACE



CONVECTION IN ATMOSPHERE
AND ENVELOPE OF DWARF
LATE TYPE STAR

IN FALL



HEAT

HEAT

INFALL STOPS RADIATIVE
COOLING AND DOWNWARD
FLOW BUT NOT UPWARD
FLOW

UNIFORMITARIANISM IN GEOLOGY

**100 YEARS AGO GEOLOGISTS WERE
UNIFORMITARIANS. THERE WERE SLOW
UPLIFTS, SLOW EROSION, SLOW LAYERING,
AND THEN THE DEPOSITS WERE UPLIFTED
AND THE CYCLE WAS REPEATED**

**ALL IMPORTANT CONSTRUCTION TOOK A
LONG TIME AT MM/YEAR**

**IN 1915 WEGENER PROPOSED CONTINENTAL
DRIFT THAT REQUIRED MOTIONS OF CM/YEAR
AND UNKNOWN PHYSICS**

**THE UNIFORMITARIANS RIDICULED THE
THEORY, SUPPRESSED RESEARCH ON IT,
AND PREVENTED PROGRESS FOR MORE
THAN A GENERATION**

**THE UNIFORMITARIANS ALSO SUPPRESSED
CATASTROPHIC EXPLANATIONS FOR EVENTS,
EVEN VIOLATING COMMON SENSE AND
EVERYDAY EXPERIENCE**

**UNIFORMITARIANISM AND CONTINENTAL
DRIFT WERE CONSISTANT AT ONE TIME
WHEN ALL THE CONTINENTS MERGED INTO
ONE, PANGEA, AND THERE WAS ONE OCEAN.
THE MOUNTAINS WORE DOWN. THERE WAS
NOT MUCH WEATHER AND NOT MUCH
WEATHERING. EVERYTHING SLOWED DOWN.**

**FORTUNATELY, THE CORE IS RADIOACTIVELY
HEATED AND CONVECTION EVENTUALLY
BROKE UP PANGEA OR WE WOULD NOT BE
HERE TODAY**

UNIFORMITARIANISM IN BIOLOGY

BIOLOGISTS USED TO THINK THAT EVOLUTION OCCURRED AT THE MUTATION RATE, RANDOMLY, SLOWLY, AND UNIFORMLY.

BUT, ACTUALLY, 90% OF EVOLUTION TAKES PLACE IN 10% OF THE TIME. "PUNCTUATED EVOLUTION".

EVOLUTION IS DRIVEN BY CATASTROPHES THAT PRODUCE LIFE AND DEATH SITUATIONS. SPECIES MUST EVOLVE OR DIE OUT WHEN THEIR ECOLOGICAL NICHE CHANGES OR DISAPPEARS.

**NATURE DOES NOT BELIEVE IN SLOW
EVOLUTION OF ANY SYSTEM**

**LONG BORING INTERVALS ARE ALWAYS
PUNCTUATED BY EXCITING CATASTROPHES**

**THE CATASTROPHES ARE BUILT INTO
SLOWLY EVOLVING SYSTEMS OR COME
FROM OUTSIDE**

UNIFORMITARIANISM IN ASTRONOMY

CONVECTION IS UNIFORMITARIANISM AT WORK. IT JUST GOES ON AND ON FOR BILLIONS OF YEARS FROM PROTOSTAR TO THE MAIN SEQUENCE AND UPWARD. BORING.

UNLESS THE MAGNETIC FIELD IN THE INTERIOR OF THE STAR HAS A SECRET LIFE, AT INTERVALS LONGER THAN PEOPLE HAVE BEEN OBSERVING.

FORMATION OF PLANETS

THERE ARE SIMPLE, LOGICAL RULES THAT ARE THE SAME FOR EVERY UNARY LATE-TYPE POP I PROTOSTAR IN THE UNIVERSE

PLANETS ARE SELF-GRAVITATING BODIES $> 10^{26}$ G FORMED FROM PROTOSTELLAR DISKS AFTER PROTOSTARS FORM

ALL UNARY DWARF LATE-TYPE STARS ARE FORMED FROM DISKS AND HAVE PLANETARY SYSTEMS LIKE OURS UNLESS AN OB STAR, OR A SUPERNOVA, OR A COLLISION HAS DISRUPTED THE DISK

BINARIES PROBABLY HAVE PLANETS AS WELL BUT I DO NOT DISCUSS THAT HERE.

QUESTIONS

HOW ARE BINARIES MADE?

WHAT IS THE PURPOSE OF PLANETS?

WHY DOES A PROTOSTAR STOP GROWING?

WHAT HAPPENS TO THE DIPOLE MAGNETIC FIELD THAT COUPLES THE PROTOSTAR TO THE DISK?

WHAT HAPPENS TO INFALLING GAS ONCE THE MAGNETIC COUPLING IS BROKEN?

WASTE MANAGEMENT

**EVERY UNARY POP I PROTOSTAR HAS THE
SAME WASTE MANAGEMENT PROBLEM**

DISK WASTE PROPERTIES:

RADIOACTIVE, SELF-HEATING

MAGNETIC, FE ~ SI

CARBONACEOUS

**DAMAGE BY RADIOACTIVITY, RADIATION,
AND PARTICLES**

**FREE RADICALS, BROKEN MOLECULES,
POSITIVE AND NEGATIVE CHARGES**

CARCINOGENIC

GENERALLY STICKY

FAR FROM STAR, ICY

DISPOSAL METHOD: DISPERSAL

NEARBY OB STARS, OR SUPERNOVAS, OR COLLISIONS BLOW EVERYTHING AWAY.

IT IS FREE ENERGY.


DISK IS WIPE BEFORE STAR IS COMPLETED OR BEFORE PLANETS ARE COMPLETED

"URANUS" AND "NEPTUNE" ARE MOST LIKELY TO BE LOST BECAUSE THEY TAKE SO LONG TO MAKE

DISPOSAL METHOD: COMPACTION

FIRST STEPS ARE TO PREVENT INFALL OF NEW WASTE AND TO GET WASTE IN DISK TO AGGLOMERATE SO IT IS EASIER TO HANDLE


PULLS THE DIPOLE MAGNETIC FIELD OUT OF THE DISK TO BREAK THE COUPLING




THE PROTOSTAR ROTATES ONCE PER DAY
DIFFERENTIAL ROTATION AS IN THE SUN
PROTOSTAR IS SLIGHTLY OBLATE

(23)

RADIATION PRESSURE PUSHES DISK MATERIAL AWAY AND FORMS A WIND OF DENSE, WARM GAS AND DUST THAT ABSORBS 10 PERCENT OF THE RADIATION FROM THE STAR.



MAGNETIC FIELD IS STRETCHED OUTWARD WITH THE DISK SINCE IT IS PINNED TO THE DISK.



21/2

WHAT HAPPENS TO THE DIPOLE MAGNETIC FIELD?

WHEN THE DIPOLE FIELD PULLS OUT OF THE DISK TO FORM THE SUPER-CME, IT FILLS THE SURFACE OF THE STAR FROM ABOUT +70 TO -70 DEG LAT

SUPER-CME ILLUMINATES BOTH SIDES OF THE DISK AS WELL AS THE INNER WALL

RADIATION AND PARTICLES BLOW AWAY ALL THE INFALLING GAS OUT TO ~ SNOW LINE AND PUSH INWARD ON BOTH SURFACES OF THE DISK SO THAT DUST AGGLOMERATES. VAPORIZES AT SURFACE

OPTICAL DEPTH OF POPCORN IS MUCH LESS THAN THAT OF DUST FROM WHICH IT IS MADE

GAS INTAKE

DUST





FILLED WITH RADIATION + WIND

RADIATION + WIND



SUPER CME





INNER
WALL



PLANETS FORM SEQUENTIALLY OUTWARD

RADIATION AND WIND FROM STAR HEAT

POPCORN AND PUSH IT OUTWARD

ORBITS MOVING OUTWARD CROSS STATIONARY

ORBITS AND ORBITS MOVING INWARD AT


AT SHALLOW ANGLE SO SLOW COLLISIONS

WALL OF AGGLOMERATED POPCORN

(= POPCORN BALLS) FORMS THAT ABSORBS

3-5% OF THE RADIATION FROM THE STAR

NEAR STAR VOLATILES ARE COOKED OUT



**POPCORN AGGLOMERATES AND CONSOLIDATES
UNTIL PLANETESIMALS FORM AND THE WALL
BECOMES OPTICALLY THIN**

**A NEW WALL FORMS FURTHER OUT WHERE
THE DISK IS INTACT AND THE PROCESS
REPEATS**

**AS THE RADIUS INCREASES THERE IS LESS
"COOKING" OF THE POPCORN**

POPCORN

STICKY

CRUSHABLE

INSULATING

RADIOACTIVELY SELF-HEATING

WARMED AND ANNEXED BY STAR

NOT LIKE PARTICLES IN


MANT BODY CALCULATIONS



LOW VELOCITY
COLLISION
STICKS



MODERATE
VELOCITY COLLISION
CRUSHES





COLLISION WITH SMALL DENSE
FAST PARTICLES PENETRATE

WHEN BODY GROW LARGER
HEAT FROM RADIOACTIVITY IS
TRAPPED IN THE CENTER AND
DIFFERENTIATION BEGINS.

WHEN GROWS TO SELF-
GRAVITATING PLANETESIMAL
CRUSHES UNDER OWN WEIGHT.
DIFFERENTIATION INCREASES.
MELTING BEGINS.

DIFFERENTIATION IS MAINTAINED
WHEN PLANETESIMAL MERGE.







**PLANETESIMALS FORMED IN THE WALL ARE
IN NEARBY ORBITS**

**THE DISPERSION OF THE DOTS IN THE
CARTOONS IS GREATLY EXAGGERATED**

**THEY RAPIDLY GRAVITATIONALLY COLLAPSE
INTO A PLANET**






**NOT ENOUGH MASS IN ASTEROID WALL TO
FORM A PLANET. LEFT WITH PLANETESIMALS**

SNOW BELT

**ICE IS ADDED TO THE POPCORN AND DUST
IN JUPITER'S WALL. A LARGE CORE FORMS
THAT IS ABLE TO PULL IN GAS**



UNIFORMITARIANISM WORKS FOR THE OUTER PLANETS

**SINCE RADIATION FROM THE STAR IS SO
WEAK AT THOSE DISTANCES, THEY ARE
ACCRETED SLOWLY BY UNASSISTED GRAVITY**

REMINDER: CFA Colloquium 4 May 2006

SPEAKER: David Wilner (CfA)

TITLE: Protoplanetary Disks:
Analogues of the Early Solar System

ABSTRACT: [edited]

The nearly circular, coplanar orbits of the planets in our Solar System have led astronomers for centuries to consider an origin within a rotating disk.

Advances in observational resolution and sensitivity, especially at millimeter wavelengths, now provide direct evidence for disks of gas and dust surrounding nearby pre-main-sequence stars, many of which have physical properties appropriate to the formation of Solar Systems like our own.

I will review briefly how we know what we know about these "protoplanetary" disks, and I will discuss how new observations are starting to capture key aspects of the planet formation process, including

- (1) dust evolution towards "pebbles", and
- (2) inner disk holes opened by disk-protoplanet interactions.

DAVE WILNER SHOWED IMAGES OF THE 20
CLOSEST "PROTOPLANETARY" DISKS
THAT ARE VISIBLE THROUGH SURROUNDING
GAS AND DUST

THE DISKS HAD UNRESOLVED CENTRAL
HOLES WITH DIAMETERS > 20 AU. SMALLER
HOLES ARE NOT VISIBLE

IN FACT, THESE ARE "POST-PLANETARY"
DISKS. PLANETS HAVE ALREADY FORMED
RAPIDLY OUT THROUGH "JUPITER"

SPEAKER: Renu Malhotra (University of Arizona)

TITLE: Bombardment History of the Terrestrial Planets

ABSTRACT: [edited]

Analysis of the impact crater record of the terrestrial planets and the Moon implicates two populations of impactors that are distinguishable by their size distributions.

The old population, responsible for an intense period of bombardment that ended 3.8 Gy ago, is virtually identical in size distribution to the present main belt asteroids;

the second population, responsible for craters younger than ~3.8 Gy, matches closely the size distribution of the near earth asteroids.

An inner Solar System impact cataclysm occurred ~3.9 Gy ago,

Many asteroids were ejected from the main belt on a short timescale, $\sim 10^7$ y, possibly due to a major dynamical instability owed to the orbital migration of the giant planets in early Solar System history.

FORMATION OF OUTER PLANETS

**RENU MALHOTRA SHOWED THAT URANUS
AND NEPTUNE TOOK ~600 MILLION YEARS
TO FORM AND MIGRATE INTO PLACE**

**THEIR FORMATION AND MIGRATION PRODUCED
RESONANCES IN THE ASTEROID BELT THAT
EITHER DROVE ASTEROIDS INWARD TO
CRATER THE INNER PLANETS, OR EJECTED
THEM FROM THE SOLAR SYSTEM**

FORMATION OF CRYPTOPLANETS

DATE: MONDAY, November 20

TIME: 11:30

ROOM: Pratt

SPEAKER: Frederic Pont (Geneva Observatory)

"Transiting Extra-solar Planets:
Towards Exo-planetology"

Abstract [edited]

It is now "Year 11" for extrasolar planet studies, and the frantic pace of progress shows no sign of slowing down.

Recent landmarks since the last Cool Star meeting include the detection of planets of a few Earth masses by radial velocity and microlensing surveys, and the characterization of several transiting hot gas giants.

Transiting planets have served as the Rosetta Stone to connect the abstract language of exoplanet orbits to the detailed knowledge of solar system planets, providing an empirical foundation to the nascent field of "exo-planetology".

Precise masses and radii are known for more than a dozen transiting gas giants, some discovered by wide-field photometric monitoring, others by radial velocity surveys.

For the brightest of these, infrared emission from the planet itself could be detected at several wavelengths with the Spitzer Space Telescope.

The Corot satellite is expected to discover dozens of transiting planets, possibly including ice giants and rock giants.

PONT'S TRANSITING PLANET TABLE

NAME	Mplanet	Rplanet	Period	a [AU]
OGLE-TR-10	0.61	1.22	3.101278	0.04162
OGLE-TR-56	1.29	1.30	1.211909	0.0225
OGLE-TR-111	0.52	1.01	4.0144479	0.0467
OGLE-TR-113	1.35	1.09	1.4324757	0.0229
OGLE-TR-132	1.14	1.18	1.689868	0.0299
HD189733	1.15	1.156	2.2185733	0.031
HD149026	0.330	0.726	2.87598	0.042
TrES-1	0.76	1.081	3.0300737	0.0393
TrES-2	1.28	1.24	2.47063	0.0367
HD209458	0.657	1.320	3.52474859	0.047
XO-1	0.90	1.184	3.941534	0.0488
HAT-P-1	0.53	1.36	4.46529	0.0551
WASP-1	0.867	1.443	2.519961	0.0382
WASP-2	0.88	1.038	2.152226	0.0307

NAME	Mstar	Rstar	Teff	[Fe/H]
OGLE-TR-10	1.10	1.14	6075	0.28
OGLE-TR-56	1.17	1.32	6119	0.19
OGLE-TR-111	0.81	0.831	5044	0.19
OGLE-TR-113	0.78	0.77	4804	0.15
OGLE-TR-132	1.26	1.34	6210	0.37
HD189733	0.82	0.753	5050	-0.03
HD149026	1.3	1.45	6147	0.36
TrES-1	0.89	0.811	5250	-0.02
TrES-2	1.08	1.00	5960	0
HD209458	1.101	1.125	6117	0.02
XO-1	1.0	0.928	5750	0.015
HAT-P-1	1.12	1.15	5975	0.13
WASP-1	1.15	1.453	6200	?
WASP-2	0.79	0.813	5200	?

PONT SAID, IN JEST, THAT EXOPLANETS SEEM
TO BE DIFFERENT FROM OUR PLANETS AND
THAT OUR PLANETS WOULD TURN OUT TO BE
WEIRD

THAT IS WHY I DECIDED TO GIVE THIS TALK

SAO detects 'puffy' new planet

BY CAROLYN MARTIN
OPA Staff Writer

The discovery of a huge, bizarre, "puffy" new planet and the announcement of a new geological identification chart for planets garnered international headlines in September, after scientists from the Smithsonian Astrophysical Observatory briefed journalists at a news conference in the Castle Commons.

The newly discovered planet, called HAT-P-1, is the largest ever found inside or outside our solar system, and it has the lowest density of any planet ever measured. It orbits a distant star about 450 light-years away in the constellation Lacerta.

Robert Noyes, SAO research astrophysicist, and Gaspar Bakos, Hubble fellow at SAO, discovered the planet using a network of small, automated telescopes.

HAT-P-1 has a radius 1.38 times Jupiter's, but its mass is only half that of Jupiter, and it is less dense than water—astronomers affectionately describe it as "puffy," Noyes explained. "If you put this planet in a giant tub of water, it would float on the surface like a cork ball."


The new planet is named for the telescopes scientists used to discover it—the HAT network of six small,

automated telescopes at Smithsonian facilities in Arizona and Hawaii. The telescopes search for planets by watching for stars that dim slightly when an orbiting planet crosses directly in front of the star as viewed

See 'Planet,' Page 6



HAT-P-1, the newly discovered planet, orbits a star 450 light-years away. Although it is the largest planet ever measured, it would float if there was a bathtub large enough to hold it. (Image by David Aguilar)



This graph plots the masses and radii of 14 known transiting exoplanets, along with Jupiter and Saturn. At least four planets, and possibly as many as five, have radii that are much larger than theory predicts. The lines represent theoretical models for close-in planets with a 20-Earth-mass core and with no core.

FROM DAVE CHARBONNEAU

"I think we're missing something fundamental about the interior structure or atmospheres of hot Jupiters."

--- Josh Winn

DATE: MONDAY, October 30
TIME: 12:00
ROOM: Pratt
SPEAKER: Jason Wright (UC Berkeley)

"Characterizing Exoplanetary Systems:
The Search for Solar System Analogs"

Abstract [edited]

For 10 years the California & Carnegie Planet Search has been collecting precision radial velocities of hundreds of Sun-like stars with ever-improving precision.

Today, this temporal baseline allows us to detect exoplanets at orbital distances of ~ 5 AU including many in systems already known to harbor an inner exoplanet.

Analysis of incomplete orbits of longer-period exoplanets allows us to peek beyond this 5 AU curtain, and recent hardware and software upgrades bring our RV precision to 1 m/s, allowing us to detect exoplanets of only a few earth masses.

Together, these achievements have helped bring the number of known multiple-planet systems up to 20 (and counting) and test theories of planet formation and migration.

Although it appears that multiplicity among exoplanets is common, the search for a true Solar System analog continues.

EXOPLANETS

**THE ONLY CONNECTION BETWEEN PLANETS
AND EXOPLANETS IS THAT THEY BOTH ORBIT
STARS**

IN GENERAL:

**EXOPLANETS ORBIT CLOSE TO THE STAR
EXOPLANETS HAVE MORE ECCENTRIC ORBITS
EXOPLANETS ARE MORE MASSIVE
EXOPLANETS ARE LESS DENSE**

**NOT A SINGLE EXOPLANET CAN BE
MATCHED TO A PLANET**

**ANYONE WITH COMMON SENSE WOULD
THINK THAT THEY ARE TWO SEPARATE
CLASSES OF OBJECT**

**A NEW CLASS OF OBJECT REQUIRES
AN EXPLANATION AND IMAGINATION**

**I DID NOT BELIEVE THAT THE FIRST
EXTRA SOLAR GIANT PLANET WAS A
PLANET. NEITHER DID DAVID BLACK.**

**I MADE UP A NAME FOR THE CLASS:
CRYPTOPLANETS**

**WHEN OBSERVATIONS IMPROVE THERE
WILL EVENTUALLY BE EXOPLANETS THAT
ARE PLANETS, SO EXOPLANET IS NOT A
VALID NAME FOR THE CLASS**

CRYPTOPLANETS

WHAT ARE THEY?

PLANETS ARE WASTE DUMPS FROM THE DISK

**MAYBE THEY ARE WASTE DUMPS, BUT NOT
FROM THE DISK**

WASTE INFALL?

WASTE ANGULAR MOMENTUM?

WASTE MAGNETIC FIELD?

**SINCE CRYPTOPLANETS ARE CLOSE TO THE
STAR, THEY ARE PROBABLY ALL THREE AND
FORM BEFORE PLANETS**


**START WITH THE SAME PROTOSTAR
MACHINERY AS BEFORE**

**STANDARD SCENARIO ASSUMES ISOLATED
EVOLUTION BUT STARS ARE FORMED IN
CLUSTERS**

**IF THERE IS A COLLISION WITH A DENSE
CLOUD, OR WITH A STAR, OR IF AN OB STAR
FORMS NEARBY, OR THERE IS A SUPERNOVA,
THERE CAN BE A PULSE OF ENHANCED
DENSITY**

**LET THE COMPLEX MACHINERY NOT WORK
PERFECTLY SO THAT THERE IS EXCESS
INFALL THAT THE DIPOLE MAGNETIC FIELD
CANNOT CONTROL**


**INFALL GAS FLOWS OVER THE SURFACE
OF THE STAR TO THE EQUATOR AND THEN
OUTWARD**





IN FALL

SURFACE



IN FALL STOPS RADIATIVE
COOLING AND DOWNWARD
FLOW BUT NOT UPWARD
FLOW


EXCESS INFALL BLOCKS CONVECTION

**ENVELOPE AND ATMOSPHERE OUTSIDE
THE MAGNETIC CYLINDER PUFFS UP,
DECOUPLES, AND SPINS UP TO
INCREASE THE MOMENT OF INERTIA**

**COMBINES WITH INFALL GAS THAT IS
ALSO MOVING OUTWARD FROM EQUATOR**

**INNER CYLINDER IS A DWARF AND
OBLATE PART IS A SUBGIANT**

**THE MATERIAL CAN NEVER GO BACK
SINGLE EVENT OR MULTIPLE EVENTS**




**THE PROTOSTAR, THE MAGNETIC FIELD,
AND THE DISK CONTINUE TO EVOLVE
AS BEFORE FOR PLANET FORMATION**

**THE COUPLING BETWEEN THE STAR AND
DISK WEAKENS AND THE FIELD LINES WRAP**

64
(4½)

MORE TO SCARE




**IN PLANET FORMATION, FIELD LINES
PULL OUT OF THE DISK AND FORM A TORUS
FILLED WITH LOW DENSITY IONS THAT
COLLAPSE INTO A SUPER-CME**

**IN CRYPTOPLANET FORMATION, THE FIELD
LINES PULL OUT OF THE DISK AND FORM A
TORUS FILLED WITH DENSE GAS**


**INITIALLY THE TOROIDAL MAGNETIC FIELD
ROTATES MORE SLOWLY THAN THE GAS INSIDE
WHICH HAS HIGH ANGULAR MOMENTUM**


**THE WHOLE APPARATUS MUST BE HIGHLY
UNSTABLE. WHEN A THIN SPOT FORMS
FROM OSCILLATIONS OR WHATEVER, THE
MAGNETIC FIELD LINES CAN RECONNECT**

**PART OF THE FIELD FALLS BACK TO MAKE
TO MAKE A SUPER-CME
PART FORMS A FREE-SPINNING
TOROIDAL RING, A TOKAMAK**




66
4 1/2










**RING IS HOT AND DENSE AND HAS A LARGE
SURFACE AREA**


**RING AND SUPER-CME ARE VERY LUMINOUS.
SUPER-CME PRODUCES STRONG PARTICLE
OUTBURSTS**

**RADIATION AND PARTICLES BLOW AWAY
ALL INFALLING GAS OUT TO ~ SNOW LINE
AND PUSH INWARD ON BOTH SURFACES
OF THE DISK SO THAT DUST AGGLOMERATES.
VAPORIZES AT SURFACE**

**SINCE RING IS EQUATORIAL, IT MAY DESTROY
MORE OF THE INNER DISK AND REDUCE INNER
PLANET FORMATION**

**RING IS UNSTABLE AND RECONNECTS IN
SECTIONS**





**HOT DENSE PLASMA BLOBS ARE
ENCAPSULATED IN SPHERICAL MAGNETIC
BOTTLES, SPHEROMAKS.**

**BLOBS ARE LARGE ENOUGH AND DENSE
ENOUGH TO BE GRAVITATIONALLY BOUND
WHEN THEY COOL, CRYPTOPLANETS**

**MAGNETIC BOTTLES SHIELD THE
CRYPTOPLANETS FROM ABLATION BY THE
STELLAR WIND**

**THE NUMBER OF CRYPTOPLANETS IS RANDOM,
UP TO A DOZEN**





**SOME OF THE CRYPTOPLANETS ARE
EJECTED FROM THE SYSTEM,
OR FALL INTO THE STAR,
OR EVAPORATE,
OR MERGE,
OR CONTINUE TO ORBIT.**

**CRYPTOPLANETS CAN ACCRETE NEUTRAL GAS
MASS CAN VARY UP TO BROWN DWARF MASS
BROWN DWARFS CAN BE CRYPTOPLANETS,
FREE OR BOUND**

SPHEROMAKS

A SPHEROMAK IS A SELF-GRAVITATING SPHERE
OF HOT IONS OF MOSTLY H AND HE
SURROUNDED BY A SPHERICALLY WOUND
MAGNETIC FIELD THAT PREVENTS IONS
FROM ESCAPING OR ENTERING

SPHEROMAKS DO NOT HAVE EXTERNAL
MAGNETIC FIELDS

DO NOT ROTATE MUCH

LARGE ORBITAL ANGULAR MOMENTUM

ABUNDANCES UNKNOWN, NOT FROM DUST
PROBABLY SUB-SOLAR LIKE YOUNG A-STAR
VEGA, LAMBDA BOO

COOLS RAPIDLY BY RADIATION AND CONTRACTS
PRESSURE AND DENSITY KEEP INCREASING

FORMS HeH^+ , H_2 , H_2^+ , H_3 , H_3^+ , HYDRIDES
CARBIDES, NITRIDES, OXIDES, AND THEIR IONS

CRYPTOPLANET CHARACTERISTICS

SPHEROMAKS EVENTUALLY COOL AND
BECOME CRYPTOPLANETS

COOL FOR A CRYPTOPLANET IS STILL $> 1000\text{K}$.

FORM TYPICAL PLANETARY MOLECULES BUT
WITH NON-PLANETARY ABUNDANCES

OPACITY IS NOT PLANETARY

INTERIOR MODELS ARE WRONG

CORE NOT AGGLOMERATED BUT LOW DENSITY

SURFACE MODELS ARE WRONG

PREDICTED ENERGY DISTRIBUTION IS WRONG

CANNOT ASSUME SED IN INTERPRETING

SPITZER OBSERVATIONS

**CRYPTOPLANETS ARE A GENERAL
PHENOMENON OF POP I UNARY LATE-TYPE
STAR FORMATION**

**THEY INDICATE THAT SOMETHING WENT
WRONG WITH THE FORMATION PROCESS
SO THAT THERE WAS TOO MUCH INFALL**

THIS MUST HAPPEN MORE THAN 1% OF THE TIME

**THE CHANCE THAT ONE OR MORE
CRYPTOPLANETS WILL CONTINUE TO ORBIT
THE STAR IS RANDOM**

**THE MASSES OF CRYPTOPLANETS ARE
RANDOM**

**CRYPTOPLANETS MAKE IT DIFFICULT
TO DETECT PLANETS**

**MODULATE RADIAL VELOCITIES
BRIGHTER THAN PLANETS**

**IN ECCENTRIC ORBITS THAT MAY SWALLOW
INNER PLANETS**

**IF THE PURPOSE OF SEARCHING FOR PLANETS
IS SEARCHING FOR LIFE, CRYPTOPLANETS
ARE NOT GOOD CANDIDATES**

WHEN THERE IS TOO MUCH INFALL, THE
INFALL FLOWS OUTWARD AT THE EQUATOR
AND HITS THE INNER WALL OF THE DISK.

THERE IS AN IMMEDIATE CATASTROPHE AND
PROBABLY MATTER IS FLUNG OUTWARD.


FU ORI STARS?

THE BIRTH OF AN FGK STAR IS DEFINED AS
THE PULLOUT OF THE DIPOLE FIELD FROM THE
DISK SO THAT INFALL STOPS AND THE
PROTOSTAR STOPS GROWING.

THE BIRTHRATE FOR FGK STARS IN OUR
GALAXY IS BETWEEN 1/DECADE AND 1/CENTURY.

IT IS A ONCE IN A LIFETIME EXPERIENCE.

THE FORMATION OF A STAR WITH
CRYPTOPLANETS HAPPENS ONCE IN A
MILLENNIUM




Infrared Spectrum of HD 209458b

NASA / JPL-Caltech / M. R. Swain (JPL/Caltech)

Spitzer Space Telescope • IRS

ssc2007-04b



Infrared Spectrum of HD 209458b
NASA / JPL-Caltech / M. R. Swain [JPL/Caltech]

Spitzer Space Telescope • IRS
ssc2007-04b

DE-OBFUSCATED SPECTRA
EITHER ANTI-CORRELATED
OR NOISE