

Chapter 3

The Floyd-Warshall Algorithm, the AP and the TSP

Section 3.1 Introduction

Let s and t be two vertices of a connected weighted graph G represented by the matrix M_G . The shortest path problem finds a path between s and t whose total edge weight is minimum.

Generally, edge-weight is taken to mean distance but the word is used loosely and may represent some other measurable quantity. Dijkstra's algorithm [1] finds the distance between s and all of the other vertices of G . However, it assumes that all edge weights are non-negative. The Floyd-Warshall algorithm [2], [5], finds the shortest paths between all pairs of nodes. Furthermore, unlike the Dijkstra algorithm, it allows arc weights to be negative. In this paper, we use the version of the F-W algorithm given in [4]. Of particular interest to us, this algorithm allows us to find all cycles of smallest total arc-weight. The running time of the F-W algorithm is $O(n^3)$. In this paper, given a random n -cycle, D , obtained from the symmetric group, S_n , we apply D^{-1} to permute the columns of M_G . Thus, if the arc $(a, D_0(b))$ has the weight $w(a, D_0(b))$ in M_G , its weight is transformed into $w(a, b)$ in $D^{-1}M_G$. It follows that a negative-weighted cycle, C , in $D^{-1}M_G$, say (a_1, a_2, \dots, a_i) , has the property that the total edge weight of DC is less than that of D . Furthermore, as long as no pair of consecutive vertices of C - $a_j a_{j+1}$ - has the property that $a_{j-1} a_j$ is an arc of D , DC will be a derangement, i.e., a permutation that moves all points in $V = \{1, 2, \dots, n\}$. In chapter 1, we used H -admissible permutations to obtain hamilton circuits in graphs. In the algorithm given here, we use admissible permutations as defined in chapter 2. Starting with a random n -cycle, they were used there to obtain a spanning set of $\lceil \log n \rceil$ disjoint cycles. Here they are used to obtain a rough approximation to an optimal solution of the edge-distance assignment problem defined by the entries in M_G . This approximation yields a weight

matrix in which the negative entries are generally fewer in number and smaller in absolute value than those in M_G . Theorem 3.1 and its corollary allow us to use considerably fewer trials than generally used in Floyd-Warshall. Once we have obtained an optimal solution, σ_{APOPT} , to the assignment problem, the only cycles we can obtain in $\sigma_{APOPT}^{-1} M_G$ are all non-negative. We then use the above theorems and the F-W algorithm to obtain the smallest positive cycles of total edge weight less than a fixed positive number, say N . As we proceed, we test cycles and sets of disjoint cycles to see if any product is an n -cycle. The n -cycle of smallest weight that we obtain is an approximate optimal solution to the TSP. The reason that it is not exact is that the F-W algorithm allows us to obtain cycles of smallest weight constructed by always using the shortest distance between any pair of vertices. It is possible that a cycle, C , may be a disjoint cycle of a permutation, s , such that $\sigma_{APOPT} s = \sigma_{TSOPT}$ where C contains a subpath $[a, \dots, b]$ that is not the shortest path between a and b . However, we do give conditions where our version of the F-W algorithm yields σ_{TSOPT} . Furthermore, we give necessary conditions in theorem 3.6 that cycles of s not obtainable by F-W must satisfy.

Section 3.2 Theorems

Theorem 3.1 *Let C be a cycle of length n . Assume that the weight $w(i, C(i))$ ($i=1,2,\dots,n$) corresponds to the arc $(i, C(i))$ of C . Then if*

$$W = \sum_{i=1}^{i=n} w(i, C(i)) < 0$$

there exists at least one value $i = i'$ with $1 \leq i' \leq n$ such that

$$\sum_{j=0}^{j=m} w(i' + j, C(i' + j)) < 0 \quad (A)$$

where $m = 0,1,2,\dots,n-1$ and $i' + j > n$ is represented by its value modulo n .

Proof. The proof is by induction. Let $n = 2$. Then one of two cases occurs. Either both arcs have negative weights or one is negative and one is positive. In the latter case, since the sum is

negative, if our first arc is negative, the theorem is proved. Assume that the theorem is true when the cycle has k arcs. We now consider the case when it has $k+1$ arcs. In this case, there must exist at least one case, say $i = i^*$, where $w(i^*) + w(i^* + 1) < 0$. Suppose this wasn't the case. Then as we traversed the cycle in a clockwise manner, the sum of the weights of any two consecutive arcs would always be non-negative. Therefore, the total sum of the weights of the cycle couldn't be negative. We now replace the nodes i^* and $i^* + 1$ by the node i^{**} whose weight is defined as $w(i^{**}) = w(i^*) + w(i^* + 1)$. The cycle C having nodes has now been replaced by the cycle C^* having k nodes. By induction, the theorem is valid for C^* . Let i' be the node of C^* defined in theorem 3.1 and $i' + j^* = i^{**}$. Then

$$\sum_{j=0}^{j=j^*} w(i' + j, C(i' + j)) < 0$$

Now consider the sum

$$\sum_{j=0}^{j=j^*-1} w(i' + j, C^*(i' + j)) + w(i^*, C(i^*))$$

The summation on the left is negative due to our inductive assumption that the theorem holds for

C^* . On the other hand, we have assumed that $W = \sum_{i=1}^{i=k} w(i, C(i)) < N$ and $w(i^*, C(i^*))$ is

negative. Therefore, the sum of these two quantities is also negative. Furthermore,

$$\sum_{j=0}^{j=j^*} w(i' + j, C(i' + j)) = \sum_{j=0}^{j=j^*-1} w(i' + j, C^*(i' + j)) + w(i^*, C(i^*)) + w(i^* + 1, C(i^* + 1)) < 0$$

Thus, if we now go back to C , using the same node i' , we can replace the node i^{**} of C^* by the nodes i^* and $i^* + 1$ of C and still satisfy condition A of the theorem, concluding the proof by induction.

Corollary 3.1. Suppose that C is a cycle such that

$$W = \sum_{i=1}^{i=n} w(i, C(i)) < N \quad (B)$$

Then there exists a value $i = i'$ such that each partial sum, S_j , has the property that

$$S_j = \sum_{j=0}^{j=m} w(i' + j, C(i' + j)) < N \quad (C)$$

always holds.

Proof. Subtract N from both sides of (B). Now let the weight of arc $(n, C(n))$

become $w(n, C(n)) - N$. From theorem 3.1,

$$W^* = W - N = \sum_{i=1}^{i=n} w(i, C(i)) < 0 \quad (D)$$

has the property that every partial sum is less than 0 where each partial sum is of the form given in C. It follows that if we add N to each side of D and restore $w(n, C(n))$ to its original value, every partial sum is less than N .

Example 3.1 We now give an example of how to obtain $i = i'$.

Let $C = (a_1, a_2, \dots, a_n)$

$$a_1 = -7, a_2 = -10, a_3 = +1, a_4 = +2, a_5 = -7, a_6 = +4, a_7 = -9,$$

$$a_8 = +11, a_9 = -2, a_{10} = -1, a_{11} = -4, a_{12} = -4, a_{13} = -8, a_{14} = +9,$$

$$a_{15} = +9, a_{16} = +21, a_{17} = +1, a_{18} = -2, a_{19} = -1, a_{20} = -3,$$

$$a_{21} = -3, a_{22} = -12, a_{23} = +6, a_{24} = +2, a_{25} = +3$$

$$-7 \ -10 \ +1 \ +2 \ -7 \ +4 \ -9 \ +11 \ -2 \ -1 \ -4 \ -4 \ -8 \ +9 \ +9 \ +21 \ +1$$

$$-2 \ -1 \ -3 \ -3 \ -12 \ +6 \ +2 \ +3$$

We now add terms with like signs going from left to right. We place the ordinal number of the first number in each sum above it.

$$1 \quad 3 \quad 5 \quad 6 \quad 7 \quad 8 \quad 9 \quad 14 \quad 18 \quad 23$$

$$-17 \ +3 \ -7 \ +4 \ -9 \ +11 \ -19 \ +40 \ -21 \ +11$$

We next add the positive number to the right of each negative number to the negative number.

$$1 \quad 5 \quad 18$$

-14 -3 +2 +21 -10

We now add terms with like signs going from left to right.

1 18

-17 +23 -10

Finally, assuming that all points lie on a circle, we add like terms going from left to right. We thus obtain

18 18

-27 +23 = -4

This tells us that i' is the eighteenth ordinal number, namely, -2 .

Thus, the partial sums are: -2, -3, -6, -9, -21, -15, -13, -10, -17, -27,

-26, -24, -31, -27, -36, -25, -27, -28, -32 -36, -44, -35, -26, -5, -4

The following theorem is essential for the use of MIN(M) throughout the algorithm.

Theorem 3.2 The Floyd-Warshall Algorithm

If we perform a triangle operation for successive values $j = 1, 2, \dots, n$, each entry d_{ik} of an $n \times n$ cost matrix M becomes equal to the value of the shortest path from i to k provided that M contains no negative cycles.

The version given here is modeled on theorem 6.4 in [4].

Proof. We shall show by induction that that after the triangle operation for $j = j_0$ is executed,

d_{ik} is the value of the shortest path with intermediate vertices $v \leq j_0$, for all i and k . The theorem

holds for $j_0 = 1$ since $v = 0$. Assume that the inductive hypothesis is true for $j = j_0 - 1$ and

consider the triangle operation for $j = j_0$:

$$d_{ik} = \min\{d_{ik}, d_{ij_0} + d_{j_0k}\}.$$

If the shortest path from i through k with $v \leq j_0$ doesn't pass through j_0 , d_{ik} will be unchanged

by this operation, the first argument in the min-operation will be selected, and d_{ik} will still satisfy

the inductive hypothesis. On the other hand if the shortest path from i to k with intermediate vertices $v \leq j_0$ does pass through j_0 , d_{ik} will be replaced by $d_{ij_0} + d_{j_0k}$. By the inductive hypothesis, d_{ij_0} and d_{j_0k} are both optimal values with intermediate vertices $v \leq j_0 - 1$.

Therefore, $d_{ij_0} + d_{j_0k}$ is optimal with intermediate vertices $v \leq j_0$.

We now give an example of how F-W works.

Example 3.2. Let $d(1, 3) = 5$, $d(3, 7) = -2$, $d(1, 7) = 25$. Then

$d(1, 3) + d(3, 7) < d(1, 7)$. Note however, that the intermediate vertex 3 comes from the fact that we have reached column $j = 3$ in the algorithm. We now substitute $d(1, 3) + d(3, 7) = 3$ for the entry in $(1, 7)$. Suppose now that $d(1, 10) = 7$ while $d(7, 10) = -5$.

$$d(1, 3) + d(3, 7) + d(7, 10) = -2 < d(1, 10) = 7.$$

The intermediate vertices are now 3 and 7. Thus, -2 can be substituted for 7 in entry $(1, 10)$.

Again, we must have reached column 7 before this substitution could be made. Now suppose that $d(10, 1) = 1$. Then our negative path implies that the negative cycle $(1 \ 3 \ 7 \ 10)$ exists.

Section 3.3 The Algorithm

PHASE 1

Let M be an $n \times n$ distance matrix. An arbitrary entry of M is $d(i, j)$.

$$V = \{1, 2, \dots, n\}$$

Step 1. Sort each row of M in ascending order of numerical value. Call the matrix obtained $\text{MIN}(M)$. An arbitrary entry (i, j) of $\text{MIN}(M)$ is $\text{ORDINAL}((i, j))$ which stands for the ordinal value of entry (i, j) in row i .

Step 2. Construct a random n -cycle, $D = (a_1 \ a_2 \ \dots \ a_n)$ whose arcs are assigned the respective values of its arcs in M . Let D^{-1} be the inverse of D .

Step 3. Let the function ORD with respect to D be $\text{ORD}(i) = a_i$. Construct $\text{ORD}^{-1}(a_i) = i$, $i = 1, 2, \dots, n$.

Step 4. Given each arc $(a_i, D(a_i))$ of D , obtain

$$DIFF(a_i) = d(a_i, D(a_i)) - d(a_i, \text{MIN}(M)(a_i, 1)).$$

(If $\text{MIN}(M)(a_i, 1)$ exists in D , we choose the second smallest value of row a_i ,

$\text{MIN}(M)(a_i, 2)$, etc. .

$MIN_1 = \min\{DIFF(a_i) \in V \mid d(a_i, D(a_i)) - d(a_i, \text{MIN}(M)(a_i, j)), j \in V\}$. We then choose

$MIN_2, \dots, MIN_{\lfloor \log n \rfloor}$, corresponding to the second, third, ..., $(\lfloor \log n \rfloor + 1)$ -th smallest values of

$DIFF(a_i)$ ($i=1, 2, \dots, n$). Generally, MIN_1 is negative.

Step 5. As we pointed out in the introduction, suppose that

$D = (1 \ 7 \ 12 \ 17 \ 20 \ 15)$ is considered to be a permutation on the points in

$\{1, 2, \dots, 20\}$. Then, given arcs $(17, D(15)), (15, D(7)), (7, D(17))$ of a graph G , we obtain

the permutation $(17 \ 15 \ 7)$ which, when applied to D , yields

$D^* = (1 \ 7 \ 20 \ 15 \ 12 \ 17)$. Since $D(15) = 1$, $D(7) = 12$, $D(17) = 20$, each of the given vertices appears in D^* . Continuing, we first work with the value $d(a_i, \text{MIN}(M)(a_i, 1))$ obtained in

MIN_1 . We assume that it is negative. We now use theorem 3.1 to obtain a cycle having a negative value such that each partial sum is negative. We note that we can never choose an arc of form

$(a, D^{-1}(a))$: When we apply D to the terminal vertex of such an arc, we obtain (a, a) , a loop.

Thus, the new permutation being constructed would not be a derangement. Continuing, we obtain

$(a_i D^{-1}(\text{MIN}(M)(a_i, 1)))$ which is given the value $d(a_i, D^{-1}(j_k))$. Letting $D^{-1}(j^k) = a_k$, we next

obtain $(a_k, D^{-1}(\text{MIN}(M)(a_k + 1))) = (a_k, D^{-1}(j_m)) = (a_k, a_m)$. If the sum of the first two terms is

negative, we continue. Otherwise, we stop. If we stop, we check to see if either the sum

$(a_i a_k) + (a_k a_i)$ or $(a_i a_k) + (a_k a_m) + (a_m a_i)$ is negative. If at least one of them is negative, we

choose the one with the smaller value and save it for use later on. If the path $[a_i, a_k, a_m]$ has a

negative value, we continue with this procedure. Checking the value of $(a_i + a_k + a_m)$, if its value

is smaller than that of the first two cycles, we substitute it for the 2-cycle saved earlier. At some point in this procedure, one of the following occurs:

- (1) The partial sum of the arcs of the path becomes positive.
- (2) The partial sum of the arcs remains negative but the path repeats a vertex therefore forming a cycle out of a subpath of arcs.

Assume (1) occurs with the path $[a_i a_k a_m \dots a_z]$. In this case, we save the cycle, C_1 , with smallest negative value. If (2) occurs, let our path be $[a_i a_k a_m \dots a_p a_q a_r \dots a_y a_m]$.

As in the first case, we have obtained the cycle, C_2 , of smallest value from among the elements of the set

$$\{(a_i a_k), (a_i a_k a_m), \dots, (a_i a_k a_m \dots a_p), \dots, (a_i a_k a_m \dots a_p \dots a_y)\}.$$

Without loss of generality, suppose the two cycles $(a_i a_k)$ and $(a_m a_p a_y)$ both have negative values. Call the product of the two cycles, P_3 . Whichever of the three permutations, C_1, C_2, P_3 has the smallest (negative) value, gives us our best negative permutation to be applied to D to obtain D_1 . On the other hand, suppose that only one of the cycles, C_1 , has a negative value. Then we choose whichever of C_1 and P_3 has the smaller negative value. We follow the same procedure starting off with each value i of $\text{MIN}(M)(i, j)$ for $j = 2, 3, \dots, [\log n] + 1$. Once we have obtained the smallest (negative) cycle, say C^* , from among all cycles tested, we multiply it by D to obtain $DC^* = D_1$, a *derangement*, i.e., a permutation which moves every point of $V = \{1, 2, \dots, n\}$. We continue with this procedure until we no longer can obtain a negative cycle. At this stage, we obtain all negative entries from $\text{MIN}(M)(i, j)$ ($i = 1, 2, \dots, [\log n] + 1; j = 1, 2, \dots, [\log n] + 1$). We then apply the algorithm given above to each entry. If we obtain no further negative cycle, we go on to Phase 2.

PHASE 2.

Before going on, a *determining vertex* of a cycle of at most value L is an initial vertex, d , of a path P that traverses C such that every subpath is of value at most L . From the corollary to theorem 3.1, at least one such vertex, d , always exists in a cycle C of value M . Phase 2 uses a modified form of the Floyd-Warshall Algorithm to obtain an optimal solution to the Assignment Problem. For short, we call this algorithm the F-W n-vs algorithm, i.e., the Floyd-Warshall negatively valued subpath algorithm. Given D_i , we construct $D_i^{-1}M^-$ where M is the value matrix. D_i^{-1} permutes the columns of M . Thus, if $D_i = (a_1 \ a_2 \ \dots \ a_n)$, $a_2 = D_i(a_1)$, etc. . Thus, if $(u \ D_i(u))$ is an arc of D_i , it is mapped onto $(u \ u)$ in $D_i^{-1}M^-$. It follows that the arcs of D_i are mapped onto the diagonal elements of $D_i^{-1}M^-$. Therefore, the initial values of the diagonal elements of $D_i^{-1}M^-$ are all zero. Furthermore, if - using theorem 3.1 – we obtain a negative cycle, C_i , in $D_i^{-1}M^-$, $D_i C_i$ has a smaller value than D_i . The value of C_i is represented by an underlined negative value lying on a diagonal entry. By permuting the columns of D_i , we obtain $C_i^{-1}D_i^{-1}M^-$. We continue this process until we no longer can obtain a negative cycle. This indicates that the last permutation obtained is σ_{APOPT} . We now discuss the details of the algorithm. We first obtain $\text{MIN}(D_i^{-1}M^-)$. It consists of sorting the row numbers of each row of M from entries of smallest value to those of largest value. Using $\text{MIN}(D_i^{-1}M^-)$, we apply the F-W algorithm to each negative entry of $D_i^{-1}M^-$. As we leave a negative entry, we write its value in italics. If we are able to extend it to a negative path, we underline the value of the new path. We extend an underlined path by using $\text{MIN}(D_i^{-1}M^-)$ to obtain negatively-valued entries from $D_i^{-1}M^-$. By the nature of the F-W algorithm, we go from column 1 to column 2, column 3, \dots , column n . An *iteration* of the algorithm is completed if we have gone through each column from 1 through n once. As an iteration proceeds, we place each of the underlined paths in a balanced, binary search tree, *NEGPATHS*. We delete underlined paths that become italicized; we add paths that have just become underlined. If an extension of a negative path has a smaller negative value than an

italicized entry, we replace the italicized entry by a new underlined path. On the other hand, if we choose an element of *NEGPATHS* that isn't smaller than the current entry, we delete it from *NEGPATHS*. In general, if a new underlined path is at an entry $(a\ b)$ where b is less than the column we are currently using in the F-W algorithm, it will still be underlined at the end of an iteration. As we go through the j -th iteration, we place the row and column numbers of each new underlined path entry in a matrix, P_{in} . If we were using the normal F-W algorithm and a negative cycle existed, we would require only one iteration to obtain the cycle, because we don't impose the conditions of theorem 3.1 upon it. Hopefully, using the modified algorithm, we will be able to lower the running time, $O(n^3)$, of the F-W algorithm. During an iteration, we stop at the moment we've obtained a negative cycle, C_l . We then construct $\text{MIN}(C_l^{-1}D_i^{-1}M^-)$. We continue the algorithm until we've either obtained a negative cycle or we have not obtained any underlined paths during an entire iteration. We use $\text{MIN}(D_i^{-1}M^-)$ to extend an underlined path as well as path matrices, P_{in} ($i = 1, 2, \dots$) to keep track of the path corresponding to each negative entry (i, j) . We use theorem 3.1, starting with a (negative) determining vertex, to construct negative paths. Unlike the F-W algorithm, we don't piece together paths to construct new smaller paths. We add only one arc from $D_i^{-1}M^-$ at a time to extend a negative path. When we have gone through columns 1, 2, ..., n , we have completed one *iteration* of this portion of the algorithm. Our goal is to obtain a negative cycle if one exists. If we are able to extend a path $(a\ b)$ to $(a\ b)(b\ c) = (a\ c)$, we underline the new entry $(a\ c)$. Once we have gone through the entry – either extending the path whose value is given at $(a\ b)$ or not doing so - we write the entry at $(a\ b)$ in italics. If no negative cycle is obtained using only n columns (as is the case in the normal F-W algorithm), we continue the algorithm into a second iteration using $D_i^{-1}(n)$. As we proceed with the iteration, after underlining the value of an extended negative path in $D_i^{-1}M^-(n)$, we place the value in *NEGPATHS*. The algorithm then proceeds as outlined earlier. As we proceed, we use the entries

in P_0 to construct P_1 . If we cannot obtain a negative cycle after using $2n$ columns, we construct $D_i^{-1}M^-(2n)$ and P_2 . We continue this procedure of iteration until either we obtain a negative cycle or we are unable to extend any negative path further. Once a negative cycle, s , is obtained, we use it to construct $D_{i+1}^{-1}M^-$.

PHASE 3.

Before going on, we make a couple of definitions. Let $M > 0$, $\varepsilon > 0$, be real positive numbers.

Assume that the smallest value that any of the negatively-valued subpaths N_i has is $-M - \varepsilon$. An F-W M -valued algorithm is one in which we may initiate an iteration using an entry whose value is no greater than M . Further assume that no positive arc has a smaller value than ε .

Step 1. We use the last matrix obtained - namely, the one which contains no negative cycles - at the start of the algorithm. Our goal is to obtain an optimal tour of M , σ_{TSPOPT} .

Step 2. A *tour* is an n cycle each of whose arcs has a value. Starting with the derangements obtained during Phase 2, we check to see if any of them are tours. We also check to see if any of the permutations generated in obtaining D_i ($i=r, r-1, r-2, \dots, 1$) yields a tour when multiplied by D_i . Here

$D_r = \sigma_{APOPT}$. If none does, we repeat Phase 1 $n(\log n)$ times and choose the smallest tour (if any) obtained. There always must be at least one smallest tour, τ_0 , since we start Phase 1 with a randomly chosen tour. In the worked-out example 3.3, we've used the difference of the total distances of σ_{APOPT} and σ_1 , the tour of smallest distance obtained using the methods just discussed. $|\sigma_1| = 161$. Thus,

$m_0 = |\sigma_1| - |\sigma_{APOPT}| = 6$. Using the corollary to theorem 3.1, we can assume that each value in

$D_3^{-1}M^-$ that belongs to a cycle is no larger than $m_0 - 1$.

Step 3a. Given a tour with an initial upper bound, m_0 , for $|\sigma_{TSPOPT}|$, we use the F-W n -nvs (*Floyd-Warshall non-negatively-valued subpath*) algorithm to obtain all lowest-valued non-negative cycles obtainable. Since we no longer have to worry about negative cycles, non-negative cycles are

obtainable provided $|\sigma_{\tau_0}| \neq |\sigma_{TSPOPT}|$. We continually test for tours using sets of disjoint cycles obtained, to see if we can find a permutation, s , such that $\tau_0 s = \tau_l$ where the value of τ_l , is of value less than m_0 , say m_l . We then assume that the upper bound for each arc is no greater than m_l as we continue the algorithm. We thus discard cycles of value not less than m_l ($i = 1, 2, \dots$). We continue the algorithm through at most n iterations.

Step 3b. Let m^* be the lowest bound obtained in Step 3a for the value of a tour. The best approach to an exhaustive search for all possible non-negative cycles is to construct a set of tree-like structures, denoted by c -trees, each of which has as its root an initial vertex of arc of value less than m^* .

Furthermore, we have a strict upper bound for the value of any subpath obtained along a branch. Let the root of one of the c -trees we are constructing be v_i . After each addition of an arc, (a, b) , to a branch, we check to see if adding the arc (b, v_i) yields a cycle of value less than m^* . If not, we continue the process. From theorem 2.1, for large n , the number of cycles in an arbitrary permutation approaches $\log n$. It follows that the average number of points in an arbitrary cycle of a permutation in

S_n is approximately $\lceil \frac{n}{\log n} \rceil$. Beginning with the $\lceil \frac{n}{\log n} \rceil$ -th node, we backtrack along any path that

hasn't been ended to see if it contains a repeated vertex. If it does, we end the path. A branch also ends when there are no arcs of value less than m^* that can extend the branch to one of value less than m^* . In general, we concentrate on the last value of each path $[a, \dots, b]$ and check if the value of $[a, \dots, b, a]$ is less than m^* . If a cycle exists, it will generally occur at an earlier node of the c -tree as a simple path rather than as a non-simple path containing a subpath that is a cycle. If we obtain a cycle, we test to see if it yields a smaller tour. If not, we continue. We collect all cycles and each time a new one is obtained, we check all sets of disjoint cycles containing it to see if the product is a permutation which when multiplied by σ_{APOPT} yields a tour of value less than $|\sigma_{FWTSPOPT}|$. It may require as many as $n-1$ iterations to obtain all cycles.

Theorem 3.3 *Using the modified F-W algorithm, we can obtain a negative cycle, C , of length L in $D_i^{-1}M$ in at most L iterations.*

Proof. We can obtain at least one arc in each iteration. Thus, we can obtain C in at most L iterations.

In general, when using our specialized version of the F-W algorithm, we choose the first negative cycle we obtain. The reason for this becomes evident when we consider the following theorem:

Theorem 3.4 *Let M be a value matrix containing both positive and negative values. Suppose that M contains one or more negative cycles. Then if a negative path P becomes a non-simple path containing a negative cycle, C , as a subpath, C is obtainable as an independent cycle in the F-W negatively-valued subpaths algorithm (F-W n-vs algorithm as used in Phase 2) using fewer columns than the number used by P to construct C .*

Proof. Let $COLUMN(i)$ be the ordered list $[1, 2, \dots, n]$ for $i = 1, 2, \dots, n$.

$COLUMNS(ALL) = \{COLUMN(i) \mid i = 1, 2, \dots, n\}$. In using any of the modified F-W algorithms, the following rules hold:

(1) If (a, b) is an element of a matrix, and $a < b$, then a and b both lie in the same

$COLUMN(i)$. Define $d(ab) = b - a$.

(2) If $b > a$ and a lies in $COLUMN(i)$, then b lies in $COLUMN(i+1)$. Define

$d(ab) = (n - a) + b$.

Let $P = [a_1 a_2 \dots a_i \dots a_{det} \dots a_{j-1} a_j]$ where $j = i$, $P' = (a_1 a_2 \dots a_i \dots a_{det} \dots a_{j-1})$ is a simple path

and $C = (a_i a_{i+1} \dots a_{det} \dots a_{j-1})$ is a negative cycle. Define $d(a_k a_{k+1}) = r_k$ for $k = 1, 2, \dots, i-1$, s_k for

$k = i, i+1, \dots, det-1$, t_k for $k = det, det+1, \dots, j-1$. Here a_{det} is a determining vertex of C . Thus,

the number of columns traversed by P is $\sum_{k=1}^{k=i-1} r_k + \sum_{k=i}^{k=det-1} s_k + \sum_{k=det}^{k=j-1} t_k$. On the other hand, the number of

columns traversed around C starting at a_{det} is $\sum_{k=det}^{k=j-1} t_k + \sum_{k=i}^{k=det-1} s_k$. Since

Theorem 3.5 Let c be any real number, $S_a = \{a_i \mid i = 1, 2, \dots, n\}$, a set of real numbers in increasing order of value. For $i = 1, 2, \dots, n$, let $b_i = a_i + c$. Then $S_b = \{b_i \mid i = 1, 2, \dots, n\}$ preserves the ordering of S_a .

Proof. The theorem merely states that adding a fixed number to a set ordered according to the value of its elements retains the same ordering as that of the original set.

Comment. This is very useful when we are dealing with entries of the value matrix which have not been changed during the algorithm. However, as the algorithm goes on, if we have entry (i, j) and $d(i, j)$, we must go through all values of $j = j'$ where the (i, j') -th entry's value in the current $\sigma_a^{-l} M^-(k)$ is less than the value of the (i, j') -th entry in $D^{-l} M^-$ (The terminology will become clearer in the examples given.)

Theorem 3.6 Suppose that m_j is the last upper bound obtained in Step 3a. If $\sigma_{TSPOPT} = \sigma_{APOPT} s$, let $C = (a \ a_1 \ a_2 \ \dots \ a_s \ b)$ be an arbitrary disjoint cycle of s where a is a determining vertex of C . Then there always exists a cycle of value less than m_j obtained in Step 3a that is of the form

$$C' = (a' b_1 \ \dots \ b_j b a b_{j+1} b_{j+2} \ \dots \ b') \ .$$

Corollary 3.6 If we can obtain no cycle in Step 3a, then $\sigma_{TSPOPT} = \sigma_{FWTSPOPT}$.

Proof. $\sigma_{TSPOPT} = \sigma_{APOPT} s$. Here s is a permutation in S_n . The value of s is less than m_j since we're

assuming that $|\sigma_{TSP}| \leq |\sigma_{FWTSPOPT}|$. Let $C = (a \ a_1 \ a_2 \ \dots \ a_s \ b)$ be a disjoint cycle of s . Let

$d(a, a_1, a_2, \dots, a_s, b) = m$ and $m \cup d(b, a) = m' < m_j$. Assume that a is a determining vertex of C .

Then there must exist a shortest path, P , in $D_i^{-l} M^-$ from a to b of value no greater than m obtainable using the normal F-W algorithm. Since the value of P is no greater than m , the value of the cycle $P \cup (b \ a) = C'$ is no greater than $m' < m_j$. Now let a' be a determining vertex of C' . Then the cycle can be obtained using the F-W non-neg algorithm in Step 3a. The corollary follows directly

from the fact that if we could obtain a cycle $[a \dots b]$ in Step 3b, then a corresponding cycle is obtainable in Step 3a.

Comment. Theorem 3.6 *does not* prove that a is a determining vertex of C' . However, by construction, a is a vertex of C' such that there exists some arc $(a \ a_l)$ in $\sigma_{FWTSPOPT}^{-1} M^-$ whose value is less than m_j . Furthermore, in constructing C , the path beginning with $(a \ a_l)$ ends when we reach the predecessor of a on C' : b . Thus, we must check the vertices of each cycle in $\sigma_{FWTSPOPT}^{-1} M^-$ to obtain those that satisfy the above condition. It follows that theorem 3.6 narrows the search for determining vertices of cycles in Step 3b to vertices in cycles obtained in Step 3a that are initial vertices of arcs whose values are smaller than m_j .

One of the advantages of our procedure for finding σ_{APOPT} is that one or more of the derangements obtained may be an n -cycle. In particular, if we repeated the first part of Phase 1 $n(\log n)$ times, using a random n -cycle each time, the probability of obtaining at least one n -cycle is

$$(1 - \frac{e}{n}) \rightarrow 1 - \frac{1}{n^e} \rightarrow 1$$

for large n . Of course, there is no guarantee that the n -cycle obtained would be close in value to $|\sigma_{TSPOPT}|$. The method given always obtains σ_{TSPOPT} in (at most) Step 3b because there always exists some permutation, p , such that $(\sigma_{APOPT}) p = \sigma_{TSPOPT}$. Whether this procedure can be done in every case in a finite amount of time is an open question.

Example 3.4

PHASE 1.

Let $D = (1 \ 2 \ 3 \ 4 \ 5 \ 6 \ 7 \ 8)$, while M is the following distance matrix:

M

	1	2	3	4	5	6	7	8	
1	∞	23	99	17	12	99	18	24	1
2	43	∞	2	73	15	100	53	28	2
3	1	84	∞	19	53	68	44	34	3
4	89	41	45	∞	40	71	79	51	4
5	83	62	94	88	∞	36	6	50	5
6	61	62	98	50	29	∞	52	40	6
7	50	21	53	68	39	26	∞	25	7
8	16	42	61	54	81	34	92	∞	8
	1	2	3	4	5	6	7	8	

In the next table, MIN(M), we give the column numbers of the entries in each row of M arranged in increasing order of value.

MIN(M)

	1	2	3	4	5	6	7	8	
1	5	4	7	2	8	3	6	1	1
2	3	5	8	1	7	4	6	2	2
3	1	4	8	7	5	6	2	3	3
4	5	2	3	8	6	7	1	4	4
5	7	6	8	2	1	4	3	5	5
6	5	8	4	7	1	2	3	6	6
7	2	8	6	5	1	3	4	7	7
8	1	6	2	4	3	5	7	8	8
	1	2	3	4	5	6	7	8	

$\text{MIN}(M)(1, 1) = (1, 5): 12$; $\text{MIN}(M)(2, 1) = (2, 3): 2$;

$\text{MIN}(M)(3, 1) = (3, 1): 1$; $\text{MIN}(M)(4, 1) = (4, 5): 40$;

$\text{MIN}(M)(5, 1) = (5, 7): 6$; $\text{MIN}(M)(6, 1) = (6, 5): 29$;

$\text{MIN}(M)(7, 1) = (7, 2): 31$; $\text{MIN}(M)(8, 1) = (8, 1) = 16$.

We now obtain the values of $\text{DIFF}(i)$ ($i = 1, 2, \dots, 8$).

$$\text{DIFF}(1) = d(1, 5) - d(1, 2) = -11.$$

$$\text{DIFF}(2) = d(2, 3) - d(2, 3) = 0.$$

$$\text{DIFF}(3) = d(3, 1) - d(3, 4) = -18.$$

$$\text{DIFF}(4) = d(4, 5) - d(4, 5) = 0.$$

$$\text{DIFF}(5) = d(5, 7) - d(5, 6) = -30.$$

$$\text{DIFF}(6) = d(6, 5) - d(6, 7) = -23.$$

$$DIFF(7) = d(7, 2) - d(7, 8) = -4.$$

$$DIFF(8) = d(8, 1) - d(8, 1) = 0.$$

$$\min \{ DIFF(i) \mid i = 1, 2, \dots, 8 \} = DIFF(5) = -30.$$

We will start Phase 1 with 5.

Before doing so, we write D and D^{-1} in ROW FORM.

$$D = \begin{array}{cccccccc} -11 & 0 & -18 & 0 & -30 & -23 & -4 & 0 \\ 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \end{array}$$

$$D^{-1} = \begin{array}{cccccccc} 2 & 3 & 4 & 5 & 6 & 7 & 8 & 1 \\ 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \end{array}$$

$$D^{-1} = \begin{array}{cccccccc} 8 & 1 & 2 & 3 & 4 & 5 & 6 & 7 \end{array}$$

TRIAL 1.

$$(5, 7) \rightarrow (5, 6) \quad -30$$

$$(6, 5) \rightarrow (6, 4) \quad -23$$

$$(4, 5) \rightarrow (4, 4)$$

$(4, 5)$ is an arc of D .

$$(4, \text{MIN}(M)(4, 2)) = (4, 2) \quad d(4, 2) - d(4, 5) = 41 - 40 = 1$$

$$(4, 2) \rightarrow (4, 1)$$

$$(1, 5) \rightarrow (1, 4) \quad -11$$

$$P = [5, 6, 4, 1, 4].$$

(1, 5) is obtained from the arc (1, 6) in M .

$$d(1, 6) - d(1, 2) = 99 - 23 = 76.$$

(4, 5) is obtained from (4, 6).

$$d(4, 6) - d(4, 5) = 71 - 40 = 31$$

(6, 5) is obtained from the loop (6, 6).

$$s_{11} = (5^{-30}6^{-23}4^11^{76}), s_{112} = (5^{-30}6^{-23}4^{31}).$$

$s_{113} = (5 \ 6)$ is not admissible since Ds_{113} is not a derangement.

The value of s_{11} is -22.

$s_{12} = (5 \ 6)(4 \ 1)$. As shown earlier, (5 6) is not admissible.

(1, 4) is obtained from the arc (1, 5) of M .

$$d(1, 5) - d(1, 2) = 12 - 23 = -11.$$

Thus, since the second cycle has a negative value, we rename it

$s_{12} = (4^11^{-11})$. A reasonable question is why we go to the bother of

saving it: s_{11} has a smaller value than s_{12} . Our reason for doing so is that

in Phase 3, we want to obtain an 8-cycle with as small as possible value.

We may thus want to check to see if Ds_{11} is an 8-cycle. In any event,

$s_{112} = (5 \ 6 \ 4)$ has the smallest negative value so far: -22.

TRIAL 2. $\text{MIN}(M)(5, 2) = (5, 6)$.

$$(5, 6) \rightarrow (5, 5)$$

(5, 6) is an arc of D . We thus can go no farther with 5.

Therefore, $s_i = (5 \ 6 \ 4)$. $Ds_i = D_i$. We obtained D_i by exchanging those rows of D that have as initial arcs 5, 6, 4, with rows (5, 7), (6, 5), (4, 6), respectively. We must construct *DIFF* values for these arcs. All other *DIFF* values remain the same.

$$\begin{array}{cccccccc} -11 & 0 & -18 & -31 & 0 & 0 & -4 & 0 \\ 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \end{array}$$

$$D_I =$$

$$2 \quad 3 \quad 4 \quad 6 \quad 7 \quad 5 \quad 8 \quad 1$$

$$1 \quad 2 \quad 3 \quad 4 \quad 5 \quad 6 \quad 7 \quad 8$$

$$D_I^{-I} =$$

$$8 \quad 1 \quad 2 \quad 3 \quad 6 \quad 4 \quad 5 \quad 7$$

$$d(5, 7) - d(5, 7) = 0, \quad d(6, 5) - d(6, 5) = 0, \quad d(4, 5) - d(4, 6) = -31.$$

We start with 4.

TRIAL 1. $\text{MIN}(M)(4, 1) = (4, 5)$.

$$(4, 5) \rightarrow (4, 6) \quad -31$$

$$(6, 5) \rightarrow (6, 6)$$

$(6, 5)$ is an arc of D_I .

$$(6, \text{MIN}(6, 2)) = (6, 8) \quad d(6, 8) - d(6, 5) = 11.$$

$$(6, 8) \rightarrow (6, 7) \quad 11$$

$$(7, 2) \rightarrow (7, 1) \quad -4$$

$$(1, 5) \rightarrow (1, 6) \quad -11$$

$$P = [4, 6, 7, 1, 6].$$

The arc $(1, 4)$ is derived from the arc $(1, 6)$ of M .

$$d(1, 6) - d(1, 2) = 99 - 23 = 76.$$

$$d(7, 6) - d(7, 8) = 26 - 25 = 1.$$

$(6, 6)$ is a loop. Thus, the cycle $(4 \ 6)$ is not an admissible cycle since

$D_I(I, 6)$ is not a derangement.

$$s_{II} = (4^{-31}6^{11}7^{-4}1^{76}), \quad s_{III} = (4^{-31}6^{11}7^1).$$

The value of s_{II} is non-negative. The value of s_{III} is -19.

The only possibility for s_{2I} is $(6^{11}7^{-4}I^{-11})$. The value of the latter cycle is

-4. We keep $s_{2l} = (6 \ 7 \ 1)$ in our bag of negative cycles.

TRIAL 2. $\text{MIN}(M)(4, 2) = (4, 2)$. $d(4, 2) - d(4, 6) = 41 - 71 = -30$.

$$(4, 2) \rightarrow (4, 1) \quad -30$$

$$(1, 5) \rightarrow (1, 6) \quad -11$$

$$(6, 5) \rightarrow (6, 6)$$

$(6, 5)$ is an arc of D_l .

$$(6, \text{MIN}(M)(6, 2)) = (6, 8) \quad d(6, 8) - d(6, 5) = 40 - 29 = 11.$$

$$(6, 8) \rightarrow (6, 7) \quad 11$$

$$(7, 2) \rightarrow (7, 1) \quad -4$$

$$P = [4, 1, 6, 7, 1].$$

$(7, 4)$ is derived from $(7, 6)$ in M .

$$d(7, 6) - d(7, 8) = 26 - 25 = 1.$$

$(6, 4)$ is derived from the loop $(6, 6)$.

$(1, 4)$ is derived from $(1, 6)$.

$$d(1, 6) - d(1, 2) = 99 - 23 = 76.$$

$$s_{2l} = (4^{-30}1^{-11}6^{11}7^1), \quad s_{2ll} = (4^{-30}1^{76}).$$

s_{2l} has a negative value: -29. s_{2ll} has a non-negative value.

s_{ll} has the smallest negative value thus far.

$(7, 1)$ is derived from the arc $(7, 2)$ of M .

$$d(7, 2) - d(7, 8) = 21 - 25 = -4.$$

Then $s_{22} = (1^{-11}6^{11}7^{-4})$. The value of s_{22} is -4. We therefore keep it in our bag of negative cycles.

TRIAL 3. $\text{MIN}(M)(4, 3) = (4, 3)$. $d(4, 3) - d(4, 6) = 45 - 71 = -26$.

$$(4, 3) \rightarrow (4, 2) \quad -26$$

$$(2, 3) \rightarrow (2, 2).$$

$(2, 3)$ is an arc of D_I .

$$(2, \text{MIN}(M)(2, 2)) = (2, 5). \quad d(2, 5) - d(2, 3) = 15 - 2 = 13.$$

$$(2, 5) \rightarrow (2, 6) \quad 13$$

$$(6, 5) \rightarrow (6, 6).$$

$(6, 5)$ is an arc of D_I .

$$(6, \text{MIN}(M)(6, 2)) = (6, 8). \quad d(6, 8) - d(6, 5) = 11$$

$$(6, 8) \rightarrow (6, 7) \quad 11$$

$$(7, 2) \rightarrow (7, 1) \quad -4$$

$$(1, 5) \rightarrow (1, 6)$$

$$P = [4, 2, 6, 7, 1, 6].$$

$$(1, 4) \text{ is derived from } (1, 6). \quad d(1, 6) - d(1, 2) = 76.$$

$$d(7, 6) - d(7, 8) = 1$$

$(6, 6)$ is a loop.

$$d(2, 6) - d(2, 3) = 100 - 2 = 98.$$

$$P = [4^{-26} 2^{13} 6^{11} 7^{-4} 1^{76}].$$

Perusing the values we obtained above, the only possible negative cycle

obtainable is $s_{313} = (4 \ 2 \ 6 \ 7)$ which has a value of -1.

$$s_{32} = (6 \ 7 \ 1) = s_{22}; \text{ its value is -1.}$$

$$\text{Thus, } s_2 = s_{21} = (4 \ 1 \ 6 \ 7).$$

In order to obtain D_2 , we respectively exchange rows $(4, 6)$, $(1, 2)$,

$(6, 5)$, $(7, 8)$ of D_I with rows $(4, 2)$, $(1, 5)$, $(6, 8)$, $(7, 6)$.

Our new rows have the following *DIFF* values:

$$d(4, 5) - d(4, 2) = -1, \quad d(1, 5) - d(1, 5) = 0, \quad d(6, 5) - d(6, 8) = -11,$$

$$d(7, 2) - d(7, 6) = -5.$$

$$0 \quad 0 \quad -18 \quad -1 \quad 0 \quad -11 \quad -5 \quad 0$$

1 2 3 4 5 6 7 8

$D_2 =$

5 3 4 2 7 8 6 1

1 2 3 4 5 6 7 8

$D_2^{-1} =$

8 4 2 3 1 7 5 6

We start with 3.

TRIAL 1.

$(3, 1) \rightarrow (3, 18) \quad -18$

$(8, 1) \rightarrow (8, 8)$

$(8, 1)$ is an arc of D_2 .

$(8, \text{MIN}(M)(8, 2)) = (8, 6). \quad d(8, 6) - d(8, 1) = 18.$

The path is non-negative. If we can go no further using the smallest negative *DIFF*, we are allowed to apply the procedure to at most the next $\lceil \log n \rceil$ smallest negative *DIFF* values.

Thus, we start with 6.

TRIAL 1.

$(6, 5) \rightarrow (6, 1) \quad -11$

$(1, 5) \rightarrow (1, 1)$

$(1, 5)$ is an arc of D_2 .

$(1, \text{MIN}(M)(1, 2)) = (1, 4). \quad d(1, 4) - d(1, 5) = 5.$

$(1, 4) \rightarrow (1, 3) \quad 5$

$$(3, 1) \rightarrow (3, 8) \quad -18$$

$$(8, 1) \rightarrow (8, 8)$$

$(8, 1)$ is an arc of D_2 .

$$(8, \text{MIN}(M)(8, 2)) = (8, 6). \quad d(8, 6) - d(8, 1) = 18.$$

$$(8, 6) \rightarrow (8, 7) \quad 18$$

$$(7, 2) \rightarrow (7, 4) \quad -5$$

$$(4, 5) \rightarrow (4, 1)$$

$$P = [6 \ 1 \ 3 \ 8 \ 7 \ 4 \ 1].$$

$(4, 6)$ is derived from $(4, 8)$.

$$d(4, 8) - d(4, 2) = 51 - 41 = 10.$$

$$d(7, 8) - d(7, 6) = 25 - 26 = -1.$$

$(8, 8)$ is a loop.

$$d(3, 8) - d(3, 4) = 34 - 19 = 15.$$

$$d(1, 8) - d(1, 5) = 24 - 12 = 12.$$

We thus obtain the following cycles from P :

$$s_{II} = (6^{-11}1^53^{-18}8^{18}7^{-5}4^{10}), \quad s_{III} = (6^{-11}1^53^{-18}8^{18}7^{-1}), \quad s_{III} = (6^{-11}1^53^{15}),$$

$$s_{II4} = (6^{-11}1^{12}).$$

s_{II} has a value of -1. s_{III} has a value of -7. The other two cycles have non-negative values.

$(4, 1)$ is derived from the arc $(4, 5)$ in M .

$$d(4, 5) - d(4, 2) = -1.$$

$$s_{I2} = (1^53^{-18}8^{18}7^{-5}4^{-1}). \quad \text{The value of } s_{I2} \text{ is } -1.$$

$$\text{TRIAL 2. } \text{MIN}(M)(6, 2)) = (6, 8). \quad d(6, 8) - d(6, 8) = 0.$$

We can proceed no further using 6.

Since we have obtained a negative cycle using 6, we need not go to 7.

Thus, let $s_3 = s_{III} = (6 \ 1 \ 3 \ 8 \ 7)$. $D_3 = D_2 s_3$.

Rows (6, 8), (1, 5), (3, 4), *, 1), (7, 6) are replaced, respectively, by rows (6, 5), (1, 4), (3, 1), (8, 6), 7, 8) to obtain D_3 .

The following are the new *DIFF* values:

$$d(6, 5) - d(6, 5) = 0.$$

$$d(1, 5) - d(1, 4) = -5.$$

$$d(3, 1) - d(3, 1) = 0.$$

$$d(8, 1) - d(8, 6) = -18.$$

$$d(7, 2) - d(7, 8) = -4.$$

$$\begin{array}{cccccccc} -5 & 0 & 0 & -1 & 0 & 0 & -4 & -15 \end{array}$$

$$\begin{array}{cccccccc} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \end{array}$$

$$D_3 =$$

$$\begin{array}{cccccccc} 4 & 3 & 1 & 2 & 7 & 5 & 8 & 6 \end{array}$$

$$\begin{array}{cccccccc} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \end{array}$$

$$D_3^{-1} =$$

$$\begin{array}{cccccccc} 3 & 4 & 2 & 1 & 6 & 8 & 5 & 7 \end{array}$$

We now start with 8.

TRIAL 1.

$$(8, 1) \rightarrow (8, 3) \quad -18$$

$$(3, 1) \rightarrow (3, 3).$$

(3, 1) is an arc of D_3 .

$$(3, \text{MIN}(M)(3, 1)) = (3, 4).$$

$$(3, 4) \rightarrow (3, 1) \quad 18$$

The path $P = [8, 3, 1]$ is non-negative.

TRIAL 2. $\text{MIN}(M)(8, 2) = (8, 6)$. $d(8, 6) - d(8, 6) = 0$. Thus, $(8, 6)$ is an arc of D_3 . We can go no further with 8.

We now start with 1.

TRIAL 1.

$$(1, 5) \rightarrow (1, 6) \quad -5$$

$$(6, 5) \rightarrow (6, 6).$$

$(6, 5)$ is an arc of D_3 .

$$(6, \text{MIN}(M)(6, 2)) = (6, 8). \quad d(6, 8) - d(6, 5) = 11.$$

$P = [1, 6, 8]$ is non-negative.

$(6, 1)$ is derived from $(6, 4)$ of M . $d(6, 4) - d(6, 5) = 50 - 29 = 21$.

The cycle $(1 \ 6)$ has a non-negative value.

TRIAL 2. $\text{MIN}(M)(1, 2) = (1, 4)$. $d(1, 4) - d(1, 4) = 0$.

$(1, 4)$ is an arc of D_3 . We can go no farther with 1.

The third smallest negative *DIFF* value is -4. It occurs at 7.

We start with 7.

TRIAL 1.

$$(7, 2) \rightarrow (7, 4) \quad -4$$

$$(4, 5) \rightarrow (4, 6) \quad -1$$

$$(6, 5) \rightarrow (6, 6).$$

$(6, 5)$ is an arc of D_3 .

$$(6, \text{MIN}(M)(6, 2)) = (6, 8). \quad d(6, 8) - d(6, 5) = 11.$$

We can go no further since $P = [7, 4, 6, 8]$ is non-negative.

$(6, 7)$ is derived from $(6, 8)$ of M .

$$d(6, 8) - d(6, 5) = 11.$$

$$d(4, 8) - d(4, 5) = 51 - 40 = 11.$$

$$s_{II} = (7^4 4^{-1} 6^{11}), \quad s_{III} = (7^4 4^{11}).$$

Both cycles have non-negative values.

$$\text{TRIAL 2. } \text{MIN}(M)(7, 2)) = (7, 8). \quad d(7, 8) - d(7, 8) = 0.$$

$(7, 8)$ is an arc of D_3 . We thus conclude Phase 1 with

$$D_3 = (1^{17} 4^{41} 2^{23} 3^1)(5^6 7^{25} 8^{34} 6^{29}).$$
 We next go to Phase 2 to try to obtain a derangement with a

smaller value than D_3 .

PHASE 2.

$$D_3^{-I} M$$

	4	3	1	2	7	5	8	6	
	1	2	3	4	5	6	7	8	
1	17	99	∞	23	16	12	24	99	1
2	73	2	43	∞	53	15	28	100	2
3	19	∞	1	84	44	53	34	68	3
4	∞	45	89	41	79	40	51	71	4
5	88	94	83	62	6	∞	50	36	5
6	50	98	61	62	52	29	40	∞	6
7	68	53	50	21	∞	39	25	26	7
8	54	61	16	42	92	81	∞	34	8
	1	2	3	4	5	6	7	8	

$$D_3^{-1}M^{-}$$

	4	3	1	2	7	5	8	6	
	1	2	3	4	5	6	7	8	
1	0	82	∞	6	1	-5	7	82	1
2	71	0	41	∞	51	13	26	98	2
3	18	∞	0	83	43	52	33	67	3
4	∞	4	48	0	38	-1	10	30	4
5	82	88	77	56	0	∞	44	30	5
6	21	69	32	33	23	0	11	∞	6
7	43	28	25	-4	∞	14	0	1	7
8	20	27	-18	8	58	47	∞	0	8
	1	2	3	4	5	6	7	8	

$j = 1, 2.$

There are no negative entries in the first and second columns.

$j = 3.$

The only negative entry in the third column is in row 8.

$d(8, 3) = -18$. $\text{MIN}(M)(3, 1) = 1$. $\sigma_4^{-1}(1) = 3$. $(3, 3)$ is a loop.

$\text{MIN}(M)(3, 2) = 4$. $\sigma_4^{-1}(4) = 1$. $d(3, 1) = 18$. $d(8, 3) + d(3, 1) \geq 0$.

$\text{MIN}(M)(3, i)$, $i = 3, 4, 5, 6, 7, 8$ are each greater than 18. Thus,

$d(8, 3) + d(3, i) > 0$, $i = 3, 4, 5, 6, 7, 8$.

$j = 4.$

The only negative entry in column four occurs in row 7.

$d(7, 4) = -4$. $\text{MIN}(M)(4, 1) = 5$. $\sigma_4^{-1}(5) = 6$. $d(4, 6) = -1$.

$d(7, 4) + d(4, 6) = -5 < d(7, 6)$. Thus, we place -5 in $(7, 6)$.

$\text{MIN}(M)(4, 2) = 2$. $\sigma_4^{-1}(2) = 4$. $(4, 4)$ is a loop.

$\text{MIN}(M)(4, 3) = 3$. $\sigma_4^{-1}(3) = 2$. $d(4, 2) = 4$. $d(7, 4) + d(4, 6) \geq 0$. The $\text{MIN}(M)(4, i)$ for $i = 4, 5, 6, 7, 8$

yield values in $\sigma_4^{-1}M^-$ greater than 4. Thus, $d(7, 4) + d(4, i) > 0$ for $i = 4, 5, 6, 7, 8$.

$P_4(4)$

	1	2	3	4	5	6	7	8	
1									1
2									2
3									3
4									4
5									5
6									6
7						1			7
8									8
	1	2	3	4	5	6	7	8	

$j = 5$.

There are no negative entries in column 5.

$j = 6$.

There are three negative entries in column 6: rows 1, 4 and 7.

$d(1, 6) = -5$. $\text{MIN}(M)(6, 1) = 5$. $\sigma_6^{-1}(5) = 6$. $(6, 6)$ is a loop.

$\text{MIN}(M)(6, 2) = 8$. $\sigma_6^{-1}(8) = 7$. $d(6, 7) = 11$. $d(1, 6) + d(6, 7) > 0$.

For $i = 3, 4, 5, 6, 7, 8$, $\text{MIN}(6, i) > \text{MIN}(M)(6, 2)$. Thus,

$$d(1, 6) + d(6, i) > 0.$$

$$d(4, 6) = -1. \text{ MIN}(M)(6, 2) = 8. \sigma_4^{-1}(8) = 7. d(6, 7) = 11.$$

$$d(4, 6) + d(6, 7) > 0. \text{ The same is true with}$$

$$\text{MIN}(M)(6, i), i = 3, 4, 5, 6, 7, 8.$$

$$d(7, 6) = -5. \text{ MIN}(M)(6, 2) = 8. \sigma_4^{-1}(8) = 7. d(6, 7) = 11.$$

$$d(4, 6) + d(6, 7) > 0. \text{ The same is true with}$$

$$\text{MIN}(M)(6, i), i = 3, 4, 5, 6, 7, 8.$$

$$j = 7, 8.$$

There are no negative entries in columns 7 and 8.

We have only one negative path in $D_3^{-1}M^-(8)$: $[7, 4, 6]$ with a value of -5. This path can't be extended in $D_3^{-1}M^-(16)$. It follows that $\sigma_{APOPT} = D_3$. The total sum of its values is 155.

PHASE 3.

We now apply Step 3a to obtain an approximation to σ_{TSPOPT} .

In order to obtain the smallest possible initial tour, we go back to D_2 to check to see if any of the permutations we obtained yields an 8-cycle when multiplied by D_2 , i.e., a tour. We first check those cycles that have negative values: s_{11} and s_{21} both have values of -1.

$$D_2 s_{11} = (1^{17} 4^{51} 8^{34} 6^{29} 5^6 7^{21} 2^2 3^1). \text{ Its value is } 161 - 6 \text{ greater than } \sigma_{APOPT}.$$

Since 6 is so small, we will let $m_0 = 6$. Furthermore, define $\sigma_l = D_2 s_{11}$.

We thus must be able to obtain a positive cycle of value no greater than 5 using only entries of $D_3^{-1}M^-$ no greater than 5. We use the F-W non- n -vs algorithm.

Comment. If we hadn't been able to obtain a tour using permutations obtained while working to obtain D_3 , we would use those obtained trying to obtain D_2 , etc. . As we go along, we may obtain cycles

that lower the upper bound from $m_0 = 6$ to m_i . From that moment onward, we only consider paths

whose value is less than m_i . This may occur a number of times during one iteration

($j = 1$ through $j = 8$). Therefore, the best approach is to complete the iteration and underline in

$D_3^{-j}M^{-}(8)$ and P_8 only those entries whose values are less than the last m_i obtained. Henceforth, we

will only give each sum of two arcs and its respective value in obtaining new entries. We underline all

paths that we are currently trying to extend. We place in italics all entries that we've passed through

always using the smallest value obtained in a path.

$$D_3^{-j}M^{-}$$

	4	3	1	2	7	5	8	6	
	1	2	3	4	5	6	7	8	
1	0	82	∞	6	1	-5	7	82	1
2	71	0	41	∞	51	13	26	98	2
3	18	∞	0	83	43	52	33	67	3
4	∞	4	48	0	38	-1	10	30	4
5	82	88	77	56	0	∞	44	30	5
6	21	69	32	33	23	0	11	∞	6
7	43	28	25	-4	∞	14	0	1	7
8	20	27	-18	8	58	47	∞	0	8
	1	2	3	4	5	6	7	8	

$j=3$.

(8, 3)(3, 1): 0.

$j=4$.

(7, 4)(4, 2): 0; (7, 4)(4, 6): -5.

$j=8$.

$(7, 8)(8, 3): -17.$

$$D_3^{-1}M^-(8)$$

	4	3	1	2	7	5	8	6	
	1	2	3	4	5	6	7	8	
1	0	82	∞	6	1	-5	7	82	1
2	71	0	41	∞	51	13	26	98	2
3	18	∞	0	83	43	52	33	67	3
4	∞	4	48	0	38	-1	10	30	4
5	82	88	77	56	0	∞	44	30	5
6	21	69	32	33	23	0	11	∞	6
7	43	<u>0</u>	<u>-17</u>	-4	∞	<u>-5</u>	0	1	7
8	<u>0</u>	27	-18	8	58	47	∞	0	8
	1	2	3	4	5	6	7	8	

$j=1.$

$(8, 1)(1, 6): -5; (8, 1)(1, 5): 1.$

$j=3.$

$(7, 3)(3, 1): 1.$

$$D_3^{-I}M^-(16)$$

	4	3	1	2	7	5	8	6	
	1	2	3	4	5	6	7	8	
1	0	82	∞	6	1	-5	7	82	1
2	71	0	41	∞	51	13	26	98	2
3	18	∞	0	83	43	52	33	67	3
4	∞	4	48	0	38	-1	10	30	4
5	82	88	77	56	0	∞	44	30	5
6	21	69	32	33	23	0	11	∞	6
7	<u>1</u>	0	-17	-4	∞	-5	0	1	7
8	0	27	-18	8	<u>1</u>	<u>-5</u>	∞	0	8
	1	2	3	4	5	6	7	8	

$j=1$.

(7, 1)(1, 5): 2

(7, 5) cannot be extended to a path of value less than 6.

It follows that $\sigma_{FWTSP_{OPT}} = \sigma_I = (1^{17} 4^{51} 8^{34} 6^{29} 7^{21} 2^2 3^1)$.

Furthermore, from the corollary to theorem 3.6, since we are unable to obtain a positive cycle of value

less than 6, $\sigma_{TSP_{OPT}} = \sigma_I$.

Example 3.5

	M																				
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
1	∞	88	72	97	14	38	9	59	39	46	52	50	29	17	48	65	2	72	86	65	1
2	80	∞	28	72	67	18	99	20	58	66	24	76	32	45	11	62	54	62	25	45	2
3	55	40	∞	82	34	26	73	76	97	17	13	33	23	94	76	87	56	32	74	81	3
4	29	76	77	∞	92	63	94	88	87	18	38	59	94	62	33	18	9	67	93	31	4
5	17	32	60	80	∞	49	21	64	77	54	41	18	91	4	35	29	10	19	99	35	5
6	21	19	31	75	19	∞	98	94	72	40	30	43	48	18	94	82	69	70	22	71	6
7	1	99	82	68	4	35	∞	74	28	44	81	59	24	33	9	10	91	95	58	89	7
8	1	49	71	49	52	30	52	∞	59	92	25	48	56	65	34	59	24	78	67	70	8
9	92	99	49	6	63	79	66	19	∞	63	90	24	92	21	4	43	77	68	84	66	9
10	46	65	2	4	80	54	92	92	72	∞	34	10	86	63	63	40	73	99	85	5	10
11	11	14	81	18	91	37	46	51	32	23	∞	90	30	85	18	66	54	85	31	19	11
12	43	1	52	64	6	39	79	89	39	44	77	∞	78	42	47	62	68	65	25	7	12
13	19	82	93	76	20	80	15	81	15	87	45	67	∞	54	9	92	16	67	7	4	13
14	84	73	89	90	56	96	31	52	28	26	23	54	19	∞	91	37	6	95	59	26	14
15	84	53	65	42	65	54	62	81	90	80	98	52	59	44	∞	18	79	39	50	91	15
16	13	5	77	60	81	5	88	17	58	48	62	12	59	20	48	∞	69	61	20	57	16
17	46	48	25	59	8	83	83	24	28	1	19	75	17	28	82	75	∞	71	31	6	17
18	31	50	84	98	26	80	67	51	83	80	82	90	42	9	3	26	68	∞	51	41	18
19	53	89	6	44	58	48	26	17	64	88	63	63	87	61	42	57	32	59	∞	68	19
20	43	6	73	51	49	52	14	56	35	18	24	80	65	47	6	55	91	85	84	∞	20
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	

MIN(M)

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
1	17	7	5	14	13	6	9	10	15	12	11	8	16	20	3	18	19	2	4	1	1
2	15	6	8	11	19	3	13	14	20	17	9	16	18	10	5	4	12	1	7	2	2
3	11	10	13	6	18	12	5	2	1	17	7	19	8	15	20	4	16	14	9	3	3
4	17	10	16	1	20	15	11	12	14	6	18	2	3	9	8	5	7	13	19	4	4
5	14	17	1	12	18	7	16	2	15	20	11	6	10	3	8	9	4	13	19	5	5
6	14	2	5	1	19	11	3	10	12	13	17	18	20	9	4	16	8	15	7	6	6
7	1	5	9	16	13	9	6	14	10	19	12	4	8	11	3	20	17	18	2	7	7
8	1	17	11	6	15	12	2	4	5	7	13	9	16	14	19	20	3	18	10	8	8
9	15	4	8	14	12	16	3	5	10	7	20	18	17	6	19	11	1	13	2	9	9
10	3	4	20	12	11	16	1	6	14	15	2	9	17	5	19	13	7	8	18	10	10
11	1	2	4	15	20	10	13	19	9	6	7	8	17	16	3	14	18	12	5	11	11
12	2	5	20	19	6	9	14	1	10	15	3	16	4	17	18	11	13	7	8	12	12
13	20	19	15	7	9	17	1	5	11	14	12	18	4	6	8	2	10	16	3	13	13
14	17	13	11	10	20	9	7	16	8	12	5	19	2	1	3	4	15	18	6	14	14
15	16	18	4	14	12	2	6	13	19	7	3	5	17	10	8	1	9	20	11	15	15
16	2	6	12	1	8	14	19	10	15	20	9	13	4	18	11	17	3	5	7	16	16
17	10	5	20	13	11	8	3	9	14	19	1	2	4	18	12	16	15	6	7	17	17
18	15	14	5	16	1	20	13	2	8	19	7	17	6	10	11	9	3	12	4	18	18
19	3	8	7	17	15	4	6	1	16	5	18	14	11	12	9	20	13	10	2	19	19
20	2	15	7	10	11	9	1	14	5	4	6	16	8	13	3	12	19	18	17	20	20
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	

$$D = (1\ 2\ 3\ 4\ 5\ 6\ 7\ 8\ 9\ 10\ 11\ 12\ 13\ 14\ 15\ 16\ 17\ 18\ 19\ 20).$$

From M , the values of the arcs of D are

$$\begin{array}{cccccccccc} 88 & 28 & 82 & 92 & 49 & 98 & 74 & 59 & 63 & 34 \\ (1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 \\ \\ 90 & 78 & 54 & 91 & 18 & 69 & 71 & 51 & 68 & 43 \\ 11 & 12 & 13 & 14 & 15 & 16 & 17 & 18 & 19 & 20). \end{array}$$

We now use $\text{MIN}(M)$ to obtain the smallest value in each row of M .

$$\text{MIN}(M)(1, 1) = 17: d(1, 17) = 2. \text{MIN}(M)(2, 1) = 15: d(2, 15) = 11.$$

$$\text{MIN}(M)(3, 1) = 11: d(3, 11) = 13. \text{MIN}(M)(4, 1) = 17: d(4, 17) = 9. \text{MIN}(M)(5, 1) = 14: d(5, 14) = 18.$$

$$\text{MIN}(M)(6, 1) = 14: d(6, 14) = 18.$$

$$\text{MIN}(M)(7, 1) = 1: d(7, 1) = 1. \text{MIN}(M)(8, 1) = 1: d(8, 1) = 1.$$

$$\text{MIN}(M)(9, 1) = 15: d(9, 15) = 4. \text{MIN}(M)(10, 1) = 3: d(10, 3) = 2.$$

$$\text{MIN}(M)(11, 1) = 1: d(11, 1) = 11. \text{MIN}(M)(12, 1) = 2: d(12, 2) = 1.$$

$$\text{MIN}(M)(13, 1) = 20: d(13, 20) = 4. \text{MIN}(M)(14, 1) = 17: d(14, 17) = 6.$$

$$\text{MIN}(M)(15, 1) = 16: d(15, 16) = 18. \text{MIN}(M)(16, 1) = 2: d(16, 2) = 5.$$

$$\text{MIN}(M)(17, 1) = 10: d(17, 10) = 1. \text{MIN}(M)(18, 1) = 15: d(18, 15) = 3.$$

$$\text{MIN}(M)(19, 1) = 3: d(19, 3) = 6. \text{MIN}(M)(20, 1) = 2: d(20, 2) = 6.$$

For $j = 1, 2, \dots, 19$, we now obtain $d(j, \text{MIN}(M)(j, 1)) - d(j, j+1)$;

also, $d(20, 2) - d(20, 1)$. We then replace the values given the arcs of D

by the corresponding differences we've just constructed. We now have

$$\begin{array}{cccccccccc}
 & -86 & -17 & -69 & -83 & -45 & -80 & -73 & -58 & -59 & -32 \\
 D = (& 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 \\
 & -79 & -77 & -50 & -85 & 0 & -64 & -70 & -48 & -62 & -37 \\
 & 11 & 12 & 13 & 14 & 15 & 16 & 17 & 18 & 19 & 20)
 \end{array}$$

We next construct a permutation consisting of at most two cycles which when applied to D will reduce the total sum of its arcs in M . To start the construction, we choose an arc where the absolute value of its difference is greatest. We therefore start our first cycle with 1. Before going on, we introduce an economical way to write permutations, their inverses and their differences as we proceed from derangement to derangement.

We first write two ordered lists of the numbers from 1 through n .

We then first enter the number corresponding to $D_i(a)$ underneath a in the first list. We then write the number preceding a underneath a in the second list. Simply speaking, if have

a

b

occurring in a column in the first list, we have

b

a

occurring in a column in the second list. If we don't know the predecessor, p , of the first point a in a cycle of a permutation, it will always occur in the second row of the permutation when we reach the arc $(p, D_i(p))$ where $D_i(p) = a$. Finally, we place the value of $DIFF(a)$ above a in the second list.

When we follow this practice for D , we must write all points of both D and D^{-1} because the permutation D is itself a derangement. However, when constructing further permutations, we need only change those columns in which the point a is contained in a permutation.

We now rewrite D and D^{-1} in the form mentioned earlier. For simplicity, we call it the *row form* of D .

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20

2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 1

1⁻⁸⁶ 2⁻¹⁷ 3⁻⁶⁹ 4⁻⁸³ 5⁻⁴⁵ 6⁻⁸⁰ 7⁻⁷³ 8⁻⁵⁸ 9⁻⁵⁹ 10⁻³² 11⁻⁷⁹ 12⁻⁷⁷ 13⁻⁵⁰ 14⁻⁸⁵

20 1 2 3 4 5 6 7 8 9 10 11 12 13

15⁰ 16⁻⁶⁴ 17⁻⁷⁰ 18⁻⁴⁸ 19⁻⁶² 20⁻³⁷

14 15 16 17 18 19

As we construct our lists, we keep track of the size of the values of *DIFF* obtained. We start off with *DIFF*(1). If and when we find a smaller value of *DIFF*, say at point *a*, we replace *DIFF*(1) by *DIFF*(*a*), replacing the preceding value of *DIFF*. Thus, when we are finished we have obtained both the point at which the smallest value of *DIFF* occurs as well as the value of *DIFF* there.

Using this procedure, we see that the smallest value of *DIFF* occurs when $a = 1$. $DIFF(1) = -86$.

The rule that we adopted in Phase 1 was that we try the first $\lfloor \log n \rfloor$ entries $\text{MIN}(M)(i, j)$ in row *i* where $\lfloor \log n \rfloor$ represents the positive integer closest to $\log n$ ($j = 1, 2, \dots, \lfloor \log n \rfloor$). $\log(20) = 2.996$.

Thus, in this case, we will pick the first three possibilities $\{\text{MIN}(M)(i, j) \mid j = 1, 2, 3\}$. There are two

more things of which we must be aware. The first is that if an arc (*a*, *b*) of D_i is chosen when

constructing a permutation, we will obtain (*a*, *a*) for the corresponding arc of the permutation. If (*a*, *b*)

comes from $\text{MIN}(M)(a, 1)$, then we must choose $\text{MIN}(M)(a, 2)$. Otherwise, we will choose

$\text{MIN}(M)(a, 1)$. Secondly, in testing for cycles, the last arc of a permutation cycle is often chosen

without knowing the arc of the value matrix from which it is obtained. Occasionally, it may occur that

the arc chosen actually comes from a loop. i.e., an arc of the form (*a*, *a*). Thus, the current

derangement would lead to a permutation which is not a derangement. To avoid this occurrence, we

must check the terminal arc of any possible which we have not obtained directly from an arc obtained from $\text{MIN}(M)$. To do this, given the permutation arc, say (a, b) , we obtain from the first two rows of the row form, $(a, D_i(b))$. If $D_i(b) \neq a$, then we have no difficulty. Otherwise, we go on to test the next possible terminal arc of a cycle.

TRIAL 1.

$\text{MIN}(M)(1,1) = 17$. $d(1, 17) = 2$. In order to construct a corresponding arc of the cycle, we must obtain

$D^{-1}(17) = 16$. Thus, the first arc of the first cycle is $(1, 16)$. We now continue with the vertex 16.

$\text{MIN}(M)(16, 1) = 2$. Thus, the second arc of M chosen is $(16, 2)$ where $d(16, 2) = 5$. $D^{-1}(2) = 1$. Thus,

the first cycle we've obtained is $(1\ 16)$. Adding up the values corresponding to 1 and 16, respectively,

if we applied the 2-cycle $(1\ 16)$ to D , we would reduce the total sum of arcs of D in M by

$$86 + 64 = 150.$$

TRIAL 2.

$\text{MIN}(M)(1,2) = 7$. $d(1, 7) = 9$. Thus, instead of - 86, we now have - 79. $D^{-1}(7) = 6$. Thus, the first arc

of the first cycle of a permutation is $(1\ 6)$. We will now proceed with the construction of the first

cycle. We must always keep in mind that the path we construct must always have a negative sum of

values. Otherwise, the permutation won't reduce the total value of the derangement to which it will be

applied.

$$(1\ 7) \quad (1\ 6) \quad - 79$$

$$(6\ 14) \quad (6\ 13) \quad - 80$$

$$(13\ 20) \quad (13\ 19) \quad - 50$$

$$(19\ 3) \quad (19\ 2) \quad - 62$$

$$(2\ 15) \quad (2\ 14) \quad - 17$$

$$(14\ 17) \quad (14\ 16) \quad - 85$$

$$(16\ 2) \quad (16\ 1) \quad - 64.$$

The cycle $(1\ 6\ 13\ 19\ 2\ 14\ 16)$ reduces the sum of the values of the arcs of D in M by 437.

TRIAL 3.

$\text{MIN}(M)(1, 3) = 5$. $d(1, 5) = 14$. Thus, the value of the arc of D out of 1 is reduced by 74.

$$(1 \ 5) \ (1 \ 4) - 74$$

$$(4 \ 17) \ (4 \ 16) - 83$$

$$(16 \ 2) \ (16 \ 1) - 64.$$

The cycle $(1 \ 4 \ 16)$ reduces the sum of the values of the arcs of D by 221.

It follows that our best choice is the second one. Let

$s_l = (1 \ 6 \ 13 \ 19 \ 2 \ 14 \ 16)$. Next, we give the row form of the new permutation, D_l , obtained by applying s_l to D . Asterisks have been placed above each point of D which has been moved.

$$\begin{array}{cccccccccccccccccccc}
 * & * & & & * & & & & * & * & & * & & * & & & & & & \\
 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 11 & 12 & 13 & 14 & 15 & 16 & 17 & 18 & 19 & 20 \\
 7 & 15 & 4 & 5 & 6 & 14 & 8 & 9 & 10 & 11 & 12 & 13 & 20 & 17 & 16 & 2 & 18 & 19 & 3 & 1 \\
 \\
 1 & 2 & 3^{-69} & 4^{-83} & 5^{-45} & 6 & 7^{-73} & 8^{-58} & 9^{-59} & 10^{-32} & 11^{-79} & 12^{-77} & 13 & 14 & 15^0 & 16 \\
 20 & 16 & 19 & 3 & 4 & 5 & 1 & 7 & 8 & 9 & 10 & 11 & 12 & 6 & 2 & 15 \\
 \\
 17^{-70} & 18^{-48} & 19 & 20^{-37} \\
 14 & 17 & 18 & 13
 \end{array}$$

We now must obtain the *DIFF* values for points under asterisks.

$$\text{MIN}(M)(1, 1) = 17 \Rightarrow \text{DIFF}(1) = d(1, 17) - d(1, 17) = 0.$$

$$\text{MIN}(M)(2, 1) = 15 \Rightarrow \text{DIFF}(2) = d(2, 15) - d(2, 15) = 0.$$

$$\text{MIN}(M)(6, 1) = 14 \Rightarrow \text{DIFF}(6) = d(6, 14) - d(6, 14) = 0.$$

$$\text{MIN}(M)(13, 1) = 20 \Rightarrow \text{DIFF}(13) = d(13, 20) - d(13, 20) = 0.$$

$$\text{MIN}(M)(14, 1) = 17 \Rightarrow \text{DIFF}(14) = d(14, 17) - d(14, 17) = 0.$$

$$\text{MIN}(M)(16, 1) = 2 \Rightarrow \text{DIFF}(16) = d(16, 2) - d(16, 2) = 0.$$

$$\text{MIN}(M)(19, 1) = 3 \Rightarrow \text{DIFF}(19) = d(19, 3) - d(19, 3) = 0.$$

Thus, we can complete the row form above.

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20

D_I

7 15 4 5 6 14 8 9 10 11 12 13 20 17 16 2 18 19 3 1

$1^0 2^0 3^{-69} 4^{-83} 5^{-45} 6^0 7^{-73} 8^{-58} 9^{-59} 10^{-32} 11^{-79} 12^{-77} 13^0 14^0$

D_I^{-1}

20 16 19 3 4 5 1 7 8 9 10 11 12 6

$15^0 16^0 17^{-70} 18^{-48} 19^0 20^{-37}$

2 15 14 17 18 13

The smallest value of *DIFF* occurs when $a = 4$.

TRIAL 1.

$$\text{MIN}(M)(4, 1) = 17.$$

$$(4, 17) \rightarrow (4, 14)$$

$$(14, 17) \rightarrow (14, 14)$$

(14, 17) is an arc of D_I . We next try to obtain $\text{MIN}(M)(14, 2) = 13$.

$$(14, 13) \rightarrow (14, 12)$$

$$(12, 2) \rightarrow (12, 16)$$

$$(16, 2) \rightarrow (16, 16)$$

(16, 2) is an arc of D_I . $\text{MIN}(M)(16, 2) = 6$.

$$(16, 6) \rightarrow (16, 5)$$

$$(5, 14) \rightarrow (5, 6)$$

$$(6, 14) \rightarrow (6, 6)$$

$(6, 14)$ is an arc of D_I .

$$(6, 2) \rightarrow (6, 16).$$

We thus have a path $P = [4, 14, 12, 16, 5, 6, 16]$.

Let $s_{II} = (4 \ 14 \ 12 \ 16 \ 5 \ 6)$.

We check to see the arc of D_I associated with $(6 \ 4)$. Going to the first two rows of *ROW FORM*, we obtain $(6 \ 5)$. In those cases in which we didn't encounter an arc of D_I in constructing an arc of s_{II} , we can use the values associated with the points of the cycle in *ROW FORM*.

Thus, $s_{II} = (4^{-83} \ 14 \ 12^{-77} \ 16 \ 5^{-45})$.

We now find the values of 14, 16 and 6. For each vertex a , we have the arc in M , obtained from $\text{MIN}(M)(a, 2)$. We then subtract the arc in D_I with the same initial vertex.

$$d(14, 13) - d(14, 17) = 19 - 6 = 13.$$

$$d(16, 6) - d(16, 2) = 5 - 5 = 0.$$

$$d(6, 5) - d(6, 14) = 19 - 18 = 1$$

$$\text{Thus, } s_{II} = (4^{-83} \ 14^{13} \ 12^{-77} \ 16^0 \ 5^{-45} \ 6^1).$$

s_{II} has the value -191. Thus, when applied to D_I , it reduces the value of D_I by 191.

We now consider the permutations obtained by including all partial paths of s_{II} which begin with 4.

Let

$$s_{III} = (4 \ 14), \ s_{II2} = (4 \ 14 \ 12), \ s_{II3} = (4 \ 14 \ 12 \ 16),$$

$$s_{II4} = (4 \ 14 \ 12 \ 16 \ 5).$$

Since 4 is mapped into 5 by D_I , in order to obtain the value of each of these permutations, we have to obtain the respective values of (14, 5), (12, 5), (16, 5). Since (5,5) is a loop, we need not consider

$$s_{114}.$$

$$d(14, 5) = 56, d(12, 5) = 6, d(16, 5) = 81.$$

We next subtract the value of each arc of form $(a, D_I(a))$

$(a = 14, 12, 5)$ from the respective values obtained above to obtain the value of each arc in its respective permutation.

$$d(14, 5) - d(14, 17) = 56 - 6 = 50.$$

$$d(12, 5) - d(14, 17) = 6 - 6 = 0.$$

$$d(16, 5) - d(16, 2) = 81 - 5 = 76.$$

$$\text{Thus, } s_{111} = (4^{-83} 14^{50}), s_{112} = (4^{-83} 14^{13} 12^0),$$

$$s_{113} = (4^{-83} 14^{13} 12^{-77} 16^{76}). \text{ Thus, the respective values of these permutations are -33, -70 and -71.}$$

We now consider the permutation obtained by using the path P to construct a permutation of two cycles.

$$s_{12} = (4 \ 14 \ 12)(16 \ 5 \ 6).$$

We already have the values of all of the points of s_{12} except for the arc (6 16). We now obtain the value associated with 6 in s_{12} .

$$d(6, 16) \rightarrow d(6, 15) = 94$$

$$d(6, 15) - d(6, 14) = 94 - 18 = 76.$$

$$s_{12} = (4^{-83} 14^{13} 12^0)(16^0 5^{-45} 6^{76}).$$

The first cycle of s_{12} is s_{112} . The value of the second cycle is non-negative so we needn't include it.

TRIAL 2. $\text{MIN}(M)(4, 2) = 10.$

$$(4, 10) \rightarrow (4, 9)$$

$$(9, 15) \rightarrow (9, 2)$$

$$(2, 15) \rightarrow (2, 2)$$

Thus, $(2, 15)$ is an arc of D_I . $\text{MIN}(M)(2, 2) = 6$

$$(2, 6) \rightarrow (2, 5)$$

$$(5, 14) \rightarrow (5, 6)$$

$$(6, 14) \rightarrow (6, 6).$$

$(6, 14)$ is an arc of D_I . $\text{MIN}(M)(6, 2) = 2$.

$$(6, 2) \rightarrow (6, 16)$$

$$(16, 2) \rightarrow (16, 16).$$

$(16, 2)$ is an arc of D_I .

$$(16, 6) \rightarrow (16, 5).$$

Let $P = [4, 9, 2, 5, 6, 16, 5]$.

$$s_{2I} = (4 \ 9 \ 2 \ 5 \ 6 \ 16).$$

We must obtain the values associated with the initial vertices 2, 6, 16. The other arcs were all obtained using terminal vertices in the first column of $\text{MIN}(M)$. Thus, we can read them off directly from *ROW FORM*. We must compute $d(16, D_I(4)) = d(16, 5)$ rather than using $d(16, 6)$.

$$d(2, 6) - d(2, 15) = 18 - 11 = 7.$$

$$d(6, 2) - d(6, 14) = 19 - 18 = 1.$$

$$d(16, 5) - d(16, 2) = 81 - 5 = 76.$$

Thus, $s_{2I} = (4^{-83} 9^{-59} 2^7 5^{-45} 6^1 16^{76})$. The value of s_{2I} is -103.

$$s_{2I1} = (4 \ 9), s_{2I2} = (4 \ 9 \ 2), s_{2I3} = (4 \ 9 \ 2 \ 5), s_{2I4} = (4 \ 9 \ 2 \ 5 \ 6).$$

We must thus obtain $d(9, 5)$, $d(2, 5)$, $d(6, 5)$. s_{2I3} need not be considered since the arc $(5, 4)$ of the permutation maps into the loop $(5, 5)$.

$$d(9, 5) = 63, d(2, 5) = 67, d(6, 5) = 19.$$

The respective values in s_{2I1} , s_{2I2} , s_{2I4} are

$$d(9, 5) - d(9, 10) = 63 - 63 = 0, d(2, 5) - d(2, 15) = 67 - 11 = 56,$$

$$d(6, 5) - d(6, 14) = 19 - 18 = 1.$$

$$s_{211} = (4^{-83} 9^0), (4^{-83} 9^{-59} 2^{56}), s_{214} = (4^{-83} 9^{-59} 2^7 5^{-45} 6^1).$$

The respective values of these permutations are -83, -86, -179.

We now construct s_{22} from P .

$s_{22} = (4 \ 9 \ 2)(5 \ 6 \ 16)$. We have already obtained the values for all vertices except 16 previously.

$$d(16, D_I(5)) = d(16, 6). d(16, 6) - d(16, 2) = 5 - 5 = 0.$$

Thus, we have

$$s_{22} = (4^{-93} 9^{-59} 2^{56})(5^{-45} 6^1 16^0).$$

The value of s_{22} is -140.

TRIAL 3. $\text{MIN}(M)(4, 3) = 16$.

$$(4, 16) \rightarrow (4, 15)$$

$$(15, 16) \rightarrow (15, 15)$$

$(15, 16)$ is an arc of D_I .

$$(15, 18) \rightarrow (15, 17)$$

$$(17, 10) \rightarrow (17, 9)$$

$$(9, 15) \rightarrow (9, 2)$$

$$(2, 15) \rightarrow (2, 2)$$

$(2, 15)$ is an arc of D_I .

$$(2, 6) \rightarrow (2, 5)$$

$$(5, 14) \rightarrow (5, 6)$$

$$(6, 14) \rightarrow (6, 6)$$

$(6, 14)$ is an arc of D_I .

$$(6, 2) \rightarrow (6, 16)$$

$$(16, 2) \rightarrow (16, 16)$$

(16, 2) is an arc of D_I .

$$(16, 6) \rightarrow (16, 5)$$

$$P = [4, 15, 17, 9, 2, 5, 6, 16, 5].$$

15, 2, 6 and 16 were the initial vertices of arcs in D_I . Thus, we had to use arcs obtained from the second column of MIN(M). We can immediately read off the values of the remaining vertices from *ROW FORM*..

$$s_{3I} = (4^{-83} 15 17^{-70} 9^{-59} 2 5^{-45} 6 16).$$

$$d(15, 18) - d(15, 16) = 39 - 18 = 21.$$

$$d(2, 6) - d(2, 15) = 18 - 11 = 7.$$

$$d(6, 2) - d(6, 14) = 19 - 18 = 1.$$

$$d(16, 5) - d(16, 2) = 81 - 5 = 76.$$

We thus obtain $s_{3I} = (4^{-83} 15^{21} 17^{-70} 9^{-59} 2^7 5^{-45} 6^1 16^{76})$. The value of s_{3I} is -152.

$s_{3II} = (4 15)$. $s_{3I2} = (4 15 17)$. $s_{3I3} = (4 15 17 9)$. $s_{3I4} = (4 15 17 9 2)$. Since (5, 4) maps into (5, 5), we need not consider (4 15 17 9 2 5).

$$s_{3I5} = (4 15 17 9 2 5 6).$$

$$d(15, 5) - d(15, 16) = 65 - 18 = 47.$$

$$d(17, 5) - d(17, 18) = 8 - 71 = -63.$$

$$d(9, 5) - d(9, 10) = 63 - 63 = 0.$$

$$d(2, 5) - d(2, 15) = 67 - 11 = 56.$$

$$d(6, 5) - d(6, 14) = 19 - 18 = 1.$$

$$d(16, 5) - d(16, 2) = 81 - 5 = 76.$$

Thus, $s_{3II} = (4^{-83} 15^{47})$, $s_{3I2} = (4^{-73} 15^{21} 17^{-73})$, $s_{3I3} = (4^{-73} 15^{21} 17^{-70} 9^0)$,

$$s_{3I4} = (4^{-83} 15^{21} 17^{-70} 9^{-59} 2^{56}), s_{3I5} = (4^{-83} 15^{21} 17^{-70} 9^{-59} 2^7 5^{-45} 6^1).$$

The respective values of these permutations are: -36, -125, -122, -135,

From P , $s_{32} = (4\ 15\ 17\ 9\ 2)(5\ 6\ 16)$. The first cycle is s_{314} . The second was obtained in trial 2 as the second factor of s_{22} . It follows that the value of s_{32} is $(-135) + (-44) = -179$.

After checking all cycles obtained, s_{315} has the smallest value: -228.

Thus, $s_2 = (4\ 15\ 17\ 9\ 2\ 5\ 6)$. In order to obtain D_2 in row form, we need only change the points 4, 15, 17, 9, 2, 5, 6. In constructing permutations, we have obtained the changes in the arcs of D_1 . However, if we wish, we can use the row form of D_1 to obtain new columns in D_2 .

Our new arcs are: (4 16), (15 18), (17 10), (9 15), (2 6), (5 14), (6 5).

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20

D_2

7 6 4 16 14 5 8 9 15 11 12 13 20 17 18 2 10 19 3 1

0 -7 -69 -9 0 -1 -73 -58 0 -32 -79 -77 0 0 -21 0 0 -48 0 -37

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20

D_2^{-1}

20 16 19 3 6 2 1 7 8 17 10 11 12 5 9 4 14 15 18 13

$$d(2, 15) - d(2, 6) = 11 - 18 = -7.$$

$$d(4, 17) - d(4, 16) = 9 - 18 = -9.$$

$$d(5, 14) - d(5, 14) = 0.$$

$$d(6, 14) - d(6, 5) = 18 - 19 = -1$$

$$d(9, 15) - d(9, 15) = 0.$$

$$d(15, 16) - d(15, 18) = 18 - 39 = -21.$$

$$d(17, 10) - d(17, 10) = 0.$$

We choose 11 as an initial vertex for our trials.

TRIAL 1. $\text{MIN}(M)(11, 1) = 1$.

$$(11, 1) \rightarrow (11, 20)$$

$$(20, 2) \rightarrow (20, 16)$$

$$(16, 2) \rightarrow (16, 16)$$

$(16, 2)$ is an arc of D_2 .

$$(16, 6) \rightarrow (16, 2)$$

$$(2, 15) \rightarrow (2, 9)$$

$$(9, 15) \rightarrow (9, 9).$$

$(9, 15)$ is an arc of D_2 .

$$(9, 4) \rightarrow (9, 3).$$

$$(3, 11) \rightarrow (3, 10)$$

$$(10, 3) \rightarrow (10, 19)$$

$$(19, 3) \rightarrow (19, 19)$$

$(19, 3)$ is an arc of D_2 .

$$(19, 8) \rightarrow (19, 7)$$

$$(7, 1) \rightarrow (7, 20)$$

$$P = [11, 20, 16, 2, 9, 3, 10, 19, 7, 20].$$

$$s_{II} = (11^{-79} 20^{-37} 16^2 9^3 10^{-69} 19^{-32} 7)$$

$$d(16, 6) - d(16, 2) = 5 - 5 = 0.$$

$$d(2, 6) - d(2, 15) = 18 - 11 = 7.$$

$$d(9, 4) - d(9, 15) = 6 - 4 = 2.$$

$$d(19, 8) - d(19, 3) = 17 - 6 = 11.$$

$$d(7, 12) - d(7, 1) = 59 - 1 = 58$$

$$s_{II} = (11^{-79} 20^{-37} 16^0 2^7 9^2 3^{-69} 10^{-32} 19^{11} 7^{58}).$$

The value of s_{II} is -139.

$$s_{111} = (11\ 20), s_{112} = (11\ 20\ 16), s_{113} = (11\ 20\ 16\ 2), s_{114} = (11\ 20\ 16\ 2\ 9), s_{115} = (11\ 20\ 16\ 2\ 9\ 3),$$

$$s_{116} = (11\ 20\ 16\ 2\ 9\ 3\ 10), s_{117} = (11\ 20\ 16\ 2\ 9\ 3\ 10\ 19).$$

$$d(20, 12) - d(20, 1) = 80 - 43 = 37.$$

$$d(16, 12) - d(16, 2) = 12 - 5 = 7.$$

$$d(2, 12) - d(2, 6) = 76 - 18 = 58.$$

$$d(9, 12) - d(9, 15) = 24 - 4 = 20.$$

$$d(3, 12) - d(3, 4) = 33 - 82 = -49.$$

$$d(10, 12) - d(10, 11) = 10 - 34 = -24.$$

$$d(19, 12) - d(19, 3) = 63 - 6 = 57.$$

$$s_{111} = (11^{-79}20^{37}), s_{112} = (11^{-79}20^{-37}16^7), s_{113} = (11^{-79}20^{-37}16^02^{58}),$$

$$s_{114} = (11^{-79}20^{-37}16^02^79^{20}), s_{115} = (11^{-79}20^{-37}16^02^79^23^{-49}).$$

$$s_{116} = (11^{-79}20^{-37}16^02^79^23^{-69}10^{-24}), s_{117} = (11^{-79}20^{-37}16^02^79^23^{-69}10^{-32}19^{57}).$$

The respective values of the above permutations are: -42, -109, -58, -89, -156, -200, -151.

$$s_{12} = (20^{-37}16^02^79^23^{-69}10^{-32}19^{11}7^0).$$

The value of s_{12} is -118.

TRIAL 2. $\text{MIN}(M)(11, 2) = 2$.

$$(11, 2) \rightarrow (11, 16)$$

From this point on, the path is the same as that of P in trial 1 up until we reach 7. Thus, let us continue on from 7 allowing P' to be

$$P' = [11, 16, 2, 9, 3, 10, 19, 7]$$

$$(7, 1) \rightarrow (7, 20)$$

$$(20, 2) \rightarrow (20, 16).$$

Thus, $P = [11, 16, 2, 9, 3, 10, 19, 7, 20, 16]$

$$s_{2l} = (11 \ 16 \ 2 \ 9 \ 3 \ 10 \ 19 \ 7 \ 20)$$

$$d(11, 2) - d(11, 12) = 14 - 90 = -76.$$

$$d(20, 12) - d(20, 1) = 80 - 43 = 37.$$

$$s_{2l} = (11^{-76}16^02^79^23^{-69}10^{-32}19^{11}7^020^{37}).$$

The value of s_{2l} is -120.

$$s_{2l1} = (11^{-76}16), s_{2l2} = (11^{-76}16^02), s_{2l3} = (11^{-76}16^02^79),$$

$$s_{2l4} = (11^{-76}16^02^79^23), s_{2l5} = (11^{-76}16^02^79^23^{-69}), s_{2l6} = (11^{-76}16^02^79^23^{-69}10^{-32}19),$$

$$s_{2l7} = (11^{-76}16^02^79^23^{-69}10^{-32}19^{11}7)$$

$$d(16, 12) - d(16, 2) = 12 - 5 = 7.$$

$$d(2, 12) - d(2, 6) = 76 - 18 = 58.$$

$$d(9, 12) - d(9, 15) = 24 - 4 = 20.$$

$$d(3, 12) - d(3, 4) = 33 - 82 = -49.$$

$$d(10, 12) - d(10, 11) = 10 - 34 = -24.$$

$$d(19, 12) - d(19, 3) = 80 - 6 = 74.$$

$$d(7, 12) - d(7, 8) = 59 - 74 = -15.$$

$$s_{2l1} = (11^{-76}16^7), s_{2l2} = (11^{-76}16^02^{58}), s_{2l3} = (11^{-76}16^02^79^{20}),$$

$$s_{2l4} = (11^{-76}16^02^79^23^{-49}), s_{2l5} = (11^{-76}16^02^79^23^{-69}10^{-24}),$$

$$s_{2l6} = (11^{-76}16^02^79^23^{-69}10^{-32}19^{74}), s_{2l7} = (11^{-76}16^02^79^23^{-69}10^{-32}19^{11}7^{-15}).$$

The respective values of the above permutations are: -69, -18, -49,

-116, -172, -94, -172.

$$s_{22} = (16^02^79^23^{-69}10^{-32}19^{11}7^020)$$

$$d(20, 4) - d(20, 1) = 51 - 43 = 8.$$

$$s_{22} = (16^02^79^23^{-69}10^{-32}19^{11}7^020^8).$$

The value of s_{22} is -73.

TRIAL 3. $\text{MIN}(M)(11, 3) = 4$.

$$(11, 4) \rightarrow (11, 3)$$

$$(3, 11) \rightarrow (3, 10)$$

Using s_{22} , we can assume that P' , a subpath of P , is

$$P' = [11, 3, 10, 19, 7, 20].$$

Continuing from 20, we obtain

$$(20, 2) \rightarrow (20, 16)$$

$$(16, 2) \rightarrow (16, 16)$$

$(16, 2)$ is an arc of D_2 .

$$(16, 6) \rightarrow (16, 2)$$

$$(2, 15) \rightarrow (2, 9)$$

$$(9, 15) \rightarrow (9, 9)$$

$(9, 15)$ is an arc of D_2 .

$$(9, 4) \rightarrow (9, 3).$$

$$P = [11, 3, 10, 19, 7, 20, 16, 2, 9, 3].$$

$$s_{3l} = (11 \ 3 \ 10 \ 19 \ 7 \ 20 \ 16 \ 2 \ 9).$$

$$d(11, 4) - d(11, 12) = 18 - 90 = -72.$$

$$d(20, 2) - d(20, 1) = 6 - 43 = -37.$$

$$d(16, 6) - d(16, 2) = 5 - 5 = 0.$$

$$d(2, 15) - d(2, 6) = 11 - 18 = -7.$$

$$d(9, 12) - d(9, 15) = 24 - 4 = 20.$$

$$s_{3l} = ((11^{-72} 3^{-69} 10^{-32} 19^{11} 7^0 20^{-37} 16^0 2^{-7} 9^{20}).$$

The value of s_{3l} is -186.

$$d(3, 12) - d(3, 4) = 33 - 82 = -59.$$

$$d(10, 12) - d(10, 11) = 10 - 34 = -24.$$

$$d(9, 15) - d(9, 4) = 4 - 6 = -2.$$

$$d(10, 3) - d(10, 12) = 2 - 10 = -8.$$

$$d(11, 1) - d(11, 1) = 0.$$

$$d(16, 2) - d(16, 6) = 5 - 5 = 0.$$

$$d(20, 2) - d(20, 2) = 0.$$

0	0	0	-9	0	0	-73	-58	-2	-8	0	-77	0	0	-21	0	0	-48	0	0
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
11	20	19	9	6	16	1	7	8	17	3	10	12	5	2	4	14	15	18	13

The smallest value of $DIFF$ occurs out of 12.

TRIAL 1. $\text{MIN}(M)(12, 1) = 2.$

$$(12, 2) \rightarrow (12, 20)$$

$$(20, 2) \rightarrow (20, 20).$$

$(20, 2)$ is an arc of D_3 .

$$(20, 15) \rightarrow (20, 2)$$

$$(2, 15) \rightarrow (2, 2).$$

$(2, 15)$ is an arc of D_3 .

$$(2, 6) \rightarrow (2, 16)$$

$$(16, 2) \rightarrow (16, 20)$$

$P = [12, 20, 2, 16, 20].$

$$s_{II} = (12\ 20\ 2\ 16)$$

$$d(20, 2) - d(20, 15) = 0.$$

$$d(2, 6) - d(2, 15) = 18 - 11 = 7.$$

$$d(16, 2) - d(16, 13) = 59 - 5 = 54.$$

$$s_{11} = (12^{-77} 20^0 2^7 16^{54}).$$

The value of s_{11} is -16.

$$s_{12} = (20 \ 2 \ 16).$$

$$d(16, 2) - d(16, 2) = 0.$$

$$s_{12} = (20^0 \ 2^7 \ 16^0).$$

The value of s_{12} is non-negative.

TRIAL 2. $\text{MIN}(M)(12, 2) = 5$.

$$d(12, 5) - d(12, 13) = 6 - 78 = -72.$$

$$(12, 5) \rightarrow (12, 6)$$

$$(6, 14) \rightarrow (6, 5)$$

$$(5, 14) \rightarrow (5, 5)$$

$(5, 14)$ is an arc of D_3 .

$$(5, 17) \rightarrow (5, 14)$$

$$(14, 17) \rightarrow (14, 14)$$

$(14, 17)$ is an arc of D_3 .

$$(14, 13) \rightarrow (14, 12)$$

$$P = [12, 6, 5, 14, 12]$$

$$s_{21} = (12 \ 6 \ 5 \ 14)$$

$$d(6, 14) - d(6, 5) = 18 - 19 = -1.$$

$$d(5, 17) - d(5, 14) = 10 - 4 = 6.$$

$$d(14, 13) - d(14, 17) = 19 - 6 = 13.$$

$$s_{21} = (12^{-72} 6^{-1} 5^6 14^{13}).$$

The value of s_{21} is -54.

TRIAL 3. $\text{MIN}(M)(12, 3) = 20$.

$$d(12, 20) - d(12, 13) = 7 - 78 = -71.$$

$$(12, 20) \rightarrow (12, 13)$$

$$(13, 20) \rightarrow (13, 13)$$

$(13, 20)$ is an arc of D_3 .

$$(13, 19) \rightarrow (13, 18)$$

$$(18, 15) \rightarrow (18, 2)$$

$$(2, 15) \rightarrow (2, 2)$$

$(2, 15)$ is an arc of D_3 .

$$(2, 6) \rightarrow (2, 16)$$

$$(16, 2) \rightarrow (16, 20)$$

$$(20, 2) \rightarrow (20, 20)$$

$(20, 2)$ is an arc of D_3 .

$$(20, 15) \rightarrow (20, 2)$$

$$P = [12, 13, 18, 2, 16, 20, 2].$$

$$s_{3l} = (12 \ 13 \ 18 \ 2 \ 16 \ 20)$$

$$(12, 13) \rightarrow (12, 20), (13, 18) \rightarrow (13, 19), (18, 2) \rightarrow (18, 15),$$

$$(2, 16) \rightarrow (2, 6), (16, 20) \rightarrow (16, 2), (20, 12) \rightarrow (20, 13).$$

$$d(12, 20) - d(12, 13) = 7 - 78 = -71.$$

$$d(13, 19) - d(13, 20) = 7 - 4 = 3.$$

$$d(18, 15) - d(18, 19) = 3 - 51 = -48.$$

$$d(2, 6) - d(2, 15) = 18 - 11 = 7.$$

$$d(16, 2) - d(16, 6) = 5 - 5 = 0.$$

$$d(20, 13) - d(20, 2) = 65 - 6 = 59.$$

$(13, 12) \rightarrow (13, 13)$. Thus, s_{3ll} need not be considered.

$$s_{3l2} = (12^{-71}13^318^{-9}), s_{3l3} = (12^{-71}13^318^{-48}2^{21}), s_{3l4} = (12^{-71}13^318^{-48}2^716^{54}).$$

The respective values of the above permutations are: -77, -95, -55.

$$d(20, 15) - d(20, 2) = 6 - 6 = 0.$$

$$s_{32} = (12^{-71}13^318^{-9})(2^716^020^0).$$

The value of the second cycle of s_{32} is non-negative, while the first cycle is precisely s_{312} .

The smallest-valued permutation is s_{313} . Thus, $s_3 = (12\ 13\ 18\ 2)$.

$$(12, 13) \rightarrow (12, 20), (13, 18) \rightarrow (13, 19), (18, 2) \rightarrow (18, 15),$$

$$(2, 12) \rightarrow (2, 13). \text{ Thus, the new arcs of } D_4 \text{ are: } (12, 20), (13, 19),$$

$$(18, 15) \text{ and } (2, 13).$$

$$D_4 \begin{array}{cccccccccccccccccccccccc} & & * & & & & & & & & & & * & * & & & & & * & & & \\ 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 11 & 12 & 13 & 14 & 15 & 16 & 17 & 18 & 19 & 20 \\ 7 & 13 & 11 & 16 & 14 & 5 & 8 & 9 & 4 & 12 & 1 & 20 & 19 & 17 & 18 & 6 & 10 & 15 & 3 & 2 \end{array}$$

In order to create D_4^I , we need only invert the four new arcs and use all of the remaining arcs from

D_3^I . Before going on, we obtain $DIFF(a)$ for $a = 2, 12, 13$ and 18.

$$d(2, 15) - d(2, 13) = 11 - 32 = -21.$$

$$d(12, 2) - d(12, 20) = 1 - 7 = -6.$$

$$d(13, 20) - d(13, 19) = 4 - 7 = -3$$

$$d(18, 15) - d(18, 15) = 0.$$

$$\begin{array}{cccccccccccccccccccccccc} 0 & -21 & 0 & -9 & 0 & 0 & -73 & -58 & -2 & -8 & 0 & -6 & -3 & 0 & -21 & 0 & 0 & 0 & 0 & 0 \\ 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 11 & 12 & 13 & 14 & 15 & 16 & 17 & 18 & 19 & 20 \\ 11 & 20 & 19 & 9 & 6 & 16 & 1 & 7 & 8 & 17 & 3 & 10 & 2 & 5 & 18 & 4 & 14 & 15 & 13 & 12 \end{array}$$

The smallest $DIFF$ value occurs when $a = 7$.

TRIAL 1.

$$(7, 1) \rightarrow (7, 11)$$

$$(11, 1) \rightarrow (11, 11)$$

$(11, 1)$ is an arc of D_4 .

$$(11, 2) \rightarrow (11, 20)$$

$$(20, 2) \rightarrow (20, 20)$$

$(20, 2)$ is an arc of D_4 .

$$(20, 15) \rightarrow (20, 18)$$

$$(18, 15) \rightarrow (18, 18)$$

$(18, 15)$ is an arc of D_4 .

$$(18, 14) \rightarrow (18, 5)$$

$$(5, 14) \rightarrow (5, 5)$$

$(5, 14)$ is an arc of D_4 .

$$(5, 17) \rightarrow (5, 14)$$

$$(14, 17) \rightarrow (14, 14)$$

$(14, 17)$ is an arc of D_4 .

$$(14, 13) \rightarrow (14, 2)$$

$$(2, 15) \rightarrow (2, 18)$$

$$P = [7, 11, 20, 18, 5, 14, 2, 18].$$

$$s_{II} = (7 \ 11 \ 20 \ 18 \ 5 \ 14 \ 2)$$

$$d(7, 1) - d(7, 8) = 1 - 74 = -73$$

$$d(11, 2) - d(11, 1) = 14 - 11 = 3$$

$$d(20, 15) - d(20, 2) = 6 - 6 = 0.$$

$$d(18, 14) - d(18, 15) = 9 - 3 = 6.$$

$$d(5, 17) - d(5, 14) = 10 - 4 = 6.$$

$$d(14, 13) - d(14, 17) = 19 - 6 = 13.$$

$$d(2, 8) - d(2, 13) = 20 - 32 = -12.$$

$$s_{11} = (7^{-73}11^320^018^65^614^{13}2^{-12}).$$

The value of s_{11} is -57.

$$d(11, 8) - d(11, 1) = 51 - 11 = 40.$$

$$d(20, 8) - d(20, 2) = 56 - 6 = 50.$$

$$d(18, 8) - d(18, 15) = 51 - 3 = 48.$$

$$d(5, 8) - d(5, 14) = 64 - 4 = 60.$$

$$d(14, 8) - d(14, 17) = 52 - 6 = 46.$$

$$s_{114} = (7^{-73}11^320^018^65^{60}), s_{115} = (7^{-73}11^320^018^65^614^{46}).$$

The respective values of the above permutations are: -33, -20, -21, -14, -12.

$$s_{12} = (7 \ 11 \ 20)(18 \ 5 \ 14) = (7^{-73}11^320^{50})(18^65^614^{85}).$$

$$d(14, 15) - d(14, 17) = 91 - 6 = 85.$$

The first cycle is s_{112} ; the second cycle is non-negative.

TRIAL 2. $\text{MIN}(M)(7, 2) = 5$.

$$d(7, 5) - d(7, 8) = 4 - 74 = -70$$

$$(7, 5) \rightarrow (7, 6)$$

$$(6, 14) \rightarrow (6, 5)$$

In Trial 1, 5 was followed by 14, 2 and 18. We continue with 18.

$$(18, 15) \rightarrow (18, 18)$$

(18, 15) is an arc of D_4 .

$$(18, 14) \rightarrow (18, 5)$$

$$P = [7, 6, 5, 14, 2, 18, 5].$$

$$s_{21} = (7 \ 6 \ 5 \ 14 \ 2 \ 18).$$

$$d(6, 14) - d(6, 5) = 18 - 19 = -1.$$

$$d(5, 17) - d(5, 14) = 6.$$

$$d(14, 13) - d(14, 17) = 13.$$

$$d(2, 15) - d(2, 13) = 11 - 32 = -21.$$

$$d(18, 8) - d(18, 15) = 51 - 3 = 48.$$

$$s_{21} = (7^{-70} 6^{-1} 5^6 14^{13} 2^{-21} 18^{48})$$

The value of s_{21} is -25.

$$d(6, 8) - d(6, 5) = 94 - 19 = 75.$$

$$d(5, 8) - d(5, 14) = 64 - 4 = 60.$$

$$d(14, 8) - d(14, 17) = 52 - 6 = 46.$$

$$d(2, 8) - d(2, 13) = 20 - 32 = -12.$$

$$s_{211} = (7^{-70} 6^{75}), s_{212} = (7^{-70} 6^0 5^{60}), s_{213} = (7^{-70} 6^0 5^6 14^{46}),$$

$$s_{214} = (7^{-70} 6^0 5^6 14^{13} 2^{-12}).$$

The respective values of the above permutations are: 5, -10, -18, -63.

$$d(2, 15) - d(2, 13) = 11 - 32 = -21.$$

$$d(18, 14) - d(18, 15) = 9 - 3 = 6.$$

$$s_{22} = (7^{-70} 6^{76})(5^6 14^{13} 2^{-21} 18^6).$$

Each cycle has a non-negative value.

TRIAL 3. $\text{MIN}(M)(7, 3) = 9$.

$$d(7, 9) - d(7, 8) = 28 - 74 = -46.$$

$$(7, 9) \rightarrow (7, 4)$$

$$(4, 17) \rightarrow (4, 14)$$

In Trial 1, we obtained 14, 2, 18, 5.

We continue with 5.

$$(5, 14) \rightarrow (5, 5)$$

(5, 14) is an arc of D_4 .

$$(5, 17) \rightarrow (5, 14)$$

$$P = [7, 4, 14, 2, 18, 5, 14].$$

$$s_{3l} = (7 \ 4 \ 14 \ 2 \ 18 \ 5).$$

$$d(4, 17) - d(4, 16) = 9 - 18 = -9.$$

$$d(14, 13) - d(14, 17) = 19 - 6 = 13.$$

$$d(2, 15) - d(2, 13) = 11 - 32 = -21.$$

$$d(18, 14) - d(18, 15) = 9 - 3 = 6.$$

$$d(5, 8) - d(5, 14) = 64 - 4 = 60.$$

$$s_{31} = (7^{-46} 4^{-9} 14^{13} 2^{-21} 18^6 5^{60}).$$

The value of s_{3l} is 3.

$$d(4, 8) - d(4, 16) = 88 - 18 = 70.$$

$$d(14, 8) - d(14, 17) = 52 - 6 = 46.$$

$$d(2, 8) - d(2, 13) = 20 - 32 = -12.$$

$$d(18, 8) - d(18, 15) = 51 - 3 = 48.$$

$$s_{311} = (7^{-46} 4^{70}), \quad s_{312} = (7^{-46} 4^{-9} 14^{46}), \quad s_{313} = (7^{-46} 4^{-9} 14^{13} 2^{-12}),$$

$$s_{314} = (7^{-46} 4^{-9} 14^{13} 2^{-21} 18^{48}).$$

The respective values of the above permutations are: 24, 1, -54, -15.

$$d(5, 17) - d(5, 14) = 10 - 4 = 6.$$

$$s_{32} = (7^{-46} 4^{70})(14^{13} 2^{-21} 18^6 5^6)$$

Both cycles are non-negative.

The smallest value of all permutations is $s_{214} = (7^{-70} 6^0 5^6 14^{13} 2^{-12})$. We obtain the following new arcs

for D_5 : (7, 5), (6, 14), (5, 17), (14, 13), (2, 8).

* * * *

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20

$$D_5$$

7 8 11 16 17 14 5 9 4 12 1 20 19 13 18 6 10 15 3 2

$$d(2, 15) - d(2, 8) = 11 - 20 = -9.$$

$$d(5, 14) - d(5, 17) = 4 - 10 = -6.$$

$$d(6, 14) - d(6, 14) = 0.$$

$$d(7, 1) - d(7, 5) = 1 - 4 = -3.$$

$$d(14, 17) - d(14, 13) = 6 - 19 = -13.$$

$$0 \ -9 \ 0 \ -9 \ -6 \ 0 \ -3 \ -58 \ -2 \ -8 \ 0 \ -6 \ -3 \ -13 \ -21 \ 0 \ 0 \ 0 \ 0 \ 0$$

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20

$$D_5^{-l}$$

11 20 19 9 7 16 1 2 8 17 3 10 14 6 18 4 5 15 13 12

We start with $a = 8$.

TRIAL 1.

$$d(8, 1) - d(8, 9) = 1 - 59 = -58.$$

$$(8, 1) \rightarrow (8, 11)$$

$$(11, 1) \rightarrow (11, 11)$$

$(11, 1)$ is an arc of D_5 .

$$(11, 2) \rightarrow (11, 20)$$

$$(20, 2) \rightarrow (20, 20)$$

$(20, 2)$ is an arc of D_5 .

$$(20, 15) \rightarrow (20, 18)$$

$$(18, 15) \rightarrow (18, 18)$$

$(18, 15)$ is an arc of D_5 .

$$(18, 14) \rightarrow (18, 6)$$

$$(6, 14) \rightarrow (6, 6)$$

$(6, 14)$ is an arc of D_5 .

$$(6, 2) \rightarrow (6, 20)$$

$$P = [8, 11, 20, 18, 6, 20].$$

$$s_{II} = (8 \ 11 \ 20 \ 18 \ 6)$$

$$d(8, 1) - d(8, 9) = -58.$$

$$d(11, 2) - d(11, 1) = 14 - 11 = 3.$$

$$d(20, 15) - d(20, 2) = 6 - 6 = 0.$$

$$d(18, 14) - d(18, 15) = 9 - 3 = 6.$$

$$d(6, 9) - d(6, 14) = 72 - 18 = 54.$$

$$s_{II} = (8^{-58}11^320^018^66^{54}).$$

$$d(11, 9) - d(11, 1) = 32 - 11 = 21.$$

$$d(20, 9) - d(20, 2) = 35 - 6 = 29.$$

$$d(18, 9) - d(18, 15) = 83 - 3 = 80.$$

$$s_{III} = (8^{-58}11^{21}), s_{II2} = (8^{-58}11^320^{29}), s_{II3} = (8^{-58}11^320^018^{80}).$$

The respective values of the above permutations are: -37, -27, 25.

$$d(6, 2) - d(6, 14) = 19 - 18 = 1.$$

$$s_{I2} = (8^{-58}11^{21})(20^018^66^1).$$

The first cycle is s_{III} . The second cycle has a non-negative value.

$$\text{TRIAL 2. MIN(M)}(8, 2) = 17.$$

$$d(8, 17) - d(8, 9) = 24 - 58 = -34.$$

$$(8, 17) \rightarrow (8, 5)$$

$$(5, 14) \rightarrow (5, 13)$$

$$(13, 20) \rightarrow (13, 2)$$

$$(2, 15) \rightarrow (2, 18)$$

$$(18, 15) \rightarrow (18, 18)$$

$(18, 15)$ is an arc of D_5 .

$$(18, 14) \rightarrow (18, 6)$$

$$(6, 14) \rightarrow (6, 6)$$

$(6, 14)$ is an arc of D_5 .

$$(6, 2) \rightarrow (6, 20)$$

$$(20, 2) \rightarrow (20, 20)$$

$(20, 2)$ is an arc of D_5 .

$$(20, 15) \rightarrow (20, 18)$$

$$P = [8, 5, 13, 2, 18, 6, 20, 18].$$

$$s_{2l} = (8\ 5\ 13\ 2\ 18\ 6\ 20).$$

$$d(8, 17) - d(8, 9) = -34.$$

$$d(5, 14) - d(5, 17) = 4 - 10 = -6.$$

$$d(13, 20) - d(13, 19) = 4 - 7 = -3.$$

$$d(2, 15) - d(2, 8) = 11 - 20 = -9.$$

$$d(18, 14) - d(18, 15) = 9 - 3 = 6.$$

$$d(6, 2) - d(6, 14) = 19 - 18 = 1.$$

$$d(20, 15) - d(20, 2) = 6 - 6 = 0.$$

$$s_{2l} = (8^{-34}5^{-6}13^{-3}2^{-9}18^66^120^{29}).$$

The value of s_{2l} is -16 .

$$d(5, 9) - d(5, 17) = 77 - 10 = 67.$$

$$d(13, 9) - d(13, 19) = 15 - 7 = 8.$$

$$d(2, 9) - d(2, 8) = 58 - 20 = 38.$$

$$d(18, 9) - d(18, 15) = 83 - 3 = 80.$$

$$d(6, 9) - d(6, 14) = 72 - 18 = 54.$$

$$s_{211} = (8^{-34}5^{67}), s_{212} = (8^{-34}5^{-6}13^8), s_{213} = (8^{-34}5^{-6}13^{-3}2^{38}),$$

$$s_{214} = (8^{-34}5^{-6}13^{-3}2^{-9}18^{80}), s_{215} = (8^{-34}5^{-6}13^{-3}2^{-9}18^66^{54}).$$

The respective values of the above permutations are: 33, -32, -5, 28, 8.

$$s_{22} = (8\ 5\ 13\ 2)(18\ 6\ 20).$$

The first cycle is s_{213} .

$d(20, 15) - d(20, 2) = 6 - 6 = 0$. Thus, the second cycle is $(18^6 6^1 20^0)$. Thus, it has a non-negative value.

TRIAL 3. $\text{MIN}(M)(8, 3) = 11$.

$$d(8, 11) - d(8, 9) = 25 - 59 = -34.$$

$$(8, 11) \rightarrow (8, 3)$$

$$(3, 11) \rightarrow (3, 3)$$

$(3, 11)$ is an arc of D_5 .

$$(3, 10) \rightarrow (3, 17)$$

$$(17, 10) \rightarrow (17, 17)$$

$(17, 10)$ is an arc of D_5 .

$$(17, 5) \rightarrow (17, 7)$$

$$(7, 1) \rightarrow (7, 11)$$

$$(11, 1) \rightarrow (11, 11)$$

$(11, 1)$ is an arc of D_5 .

$$(11, 2) \rightarrow (11, 20)$$

$$(20, 2) \rightarrow (20, 20)$$

$(20, 2)$ is an arc of D_5 .

$$(20, 15) \rightarrow (20, 18)$$

$$(18, 15) \rightarrow (18, 18)$$

$(18, 15)$ is an arc of D_5 .

$$(18, 14) \rightarrow (18, 6)$$

$$(6, 14) \rightarrow (6, 6)$$

$(6, 14)$ is an arc of D_5 .

$$(6, 2) \rightarrow (6, 20)$$

$$(20, 2) \rightarrow (20, 20)$$

$(20, 2)$ is an arc of D_5 .

$$(20, 15) \rightarrow (20, 18)$$

$$P = [8, 3, 17, 7, 11, 20, 18, 6, 20].$$

$$d(3, 10) - d(3, 11) = 17 - 13 = 4.$$

$$d(17, 5) - d(17, 10) = 8 - 1 = 7.$$

$$d(7, 1) - d(7, 5) = 1 - 4 = -3.$$

$$d(11, 2) - d(11, 1) = 14 - 11 = 3.$$

$$d(20, 15) - d(20, 2) = 6 - 6 = 0.$$

$$d(18, 14) - d(18, 15) = 9 - 3 = 6.$$

$$d(6, 9) - d(6, 14) = 72 - 18 = 54.$$

$$s_{3l} = (8^{-34}3^417^77^{-3}11^320^018^66^{54}).$$

$$d(3, 9) - d(3, 11) = 97 - 13 = 84.$$

$$d(17, 9) - d(17, 10) = 28 - 1 = 27.$$

$$d(7, 9) - d(7, 5) = 28 - 4 = 24.$$

$$d(11, 9) - d(11, 2) = 32 - 14 = 18.$$

$$d(20, 9) - d(20, 2) = 35 - 6 = 29.$$

$$d(18, 9) - d(18, 15) = 83 - 3 = 80.$$

$$s_{311} = (8^{-34} 3^{84}), s_{312} = (8^{-34} 3^4 17^{27}), s_{313} = (8^{-34} 3^4 17^7 7^{24}),$$

$$s_{314} = (8^{-34} 3^4 17^7 7^{-3} 11^{18}), s_{315} = (8^{-34} 3^4 17^7 7^{-3} 11^3 20^{29}),$$

$$s_{316} = (8^{-34} 3^4 17^7 7^{-3} 11^3 20^0 18^{80}).$$

$$s_{32} = (8^{-34} 3^4 17^7 7^{-3} 11^{18})(20^0 18^6 6^1).$$

The first cycle is s_{314} . The second is non-negative.

The smallest negative cycle is $s_{111} = (8^{-58} 11^{21})$. Thus, $s_4 = (8 \ 11)$.

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20

D_6

7 8 11 16 17 14 5 1 4 12 9 20 19 13 18 6 10 15 3 2

-7 -9 0 -9 -6 0 -3 0 -2 -8 -21 -6 -3 -13 -21 0 0 0 0 0

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20

D_6^{-1}

8 20 19 9 7 16 1 2 11 17 3 10 14 6 18 4 5 15 13 12

TRIAL 1. $\text{MIN}(M)(15, 1) = 16$.

$$d(15, 16) - d(15, 18) = 18 - 39 = -21.$$

$$(15, 16) \rightarrow (15, 4)$$

$$(4, 17) \rightarrow (4, 5)$$

$$(5, 14) \rightarrow (5, 6)$$

$$(6, 14) \rightarrow (6, 6)$$

$(6, 14)$ is an arc of D_6 .

$$(6, 2) \rightarrow (6, 20)$$

$$(20, 2) \rightarrow (20, 20)$$

$(20, 2)$ is an arc of D_6 .

$$(20, 15) \rightarrow (20, 18)$$

$$(18, 15) \rightarrow (18, 18)$$

$(18, 15)$ is an arc of D_6 .

$$(18, 14) \rightarrow (18, 6)$$

$$P = [15, 4, 5, 6, 20, 18, 6].$$

$$d(15, 16) - d(15, 18) = 18 - 39 = -21.$$

$$d(4, 17) - d(4, 16) = 9 - 18 = -9$$

$$d(5, 14) - d(5, 17) = 4 - 9 = -5.$$

$$d(6, 2) - d(6, 14) = 19 - 18 = 1.$$

$$d(20, 15) - d(20, 2) = 6 - 6 = 0.$$

$$s_{II} = (15\ 4\ 5\ 6\ 20\ 18).$$

We can't use s_{II} since $D_6(18) = 18$. Thus, $D_6 s_{II}$ wouldn't be a derangement.

$$d(4, 18) - d(4, 16) = 67 - 18 = 49.$$

$$d(5, 18) - d(5, 17) = 19 - 10 = 9.$$

$$d(6, 18) - d(6, 14) = 70 - 18 = 52.$$

$$d(20, 18) - d(20, 2) = 85 - 6 = 79.$$

$$s_{III} = (15^{-21}4^{49}), s_{II2} = (15^{-21}4^{-9}5^9), s_{II3} = (15^{-21}4^{-9}5^{-5}6^{52}),$$

$$s_{II4} = (15^{-21}4^{-9}5^{-5}6^1 20^{79}) .$$

The respective values of the above permutations are: 28, -21, 17, 45.

$$s_{22} = (15^{-21}4^{-9}5^9)(6^1 20^0 18^6).$$

$$(8, 17) \rightarrow (8, 10)$$

$$(10, 3) \rightarrow (10, 19)$$

$$(19, 3) \rightarrow (19, 19)$$

$(19, 3)$ is an arc of D_7 .

$$(19, 8) \rightarrow (19, 2)$$

$$(2, 15) \rightarrow (2, 18)$$

$$(18, 15) \rightarrow (18, 18)$$

$(18, 15)$ is an arc of D_7 .

$$(18, 14) \rightarrow (18, 6)$$

$$(6, 14) \rightarrow (6, 6)$$

$(6, 14)$ is an arc of D_7 .

$$(6, 2) \rightarrow (6, 20)$$

$$(20, 2) \rightarrow (20, 20)$$

$(20, 2)$ is an arc of D_7 .

$$(20, 15) \rightarrow (20, 18)$$

$$P = [11, 8, 10, 19, 2, 18, 6, 20, 18]$$

$$d(8, 17) - d(8, 1) = 24 - 1 = 23.$$

The path $[11, 8, 10]$ has a positive value $-21 + 23 = 2$.

We need proceed no further in Trial 1. The only possible permutation is $(11\ 8)$.

$$d(11, 1) - d(11, 9) = -21.$$

$$d(8, 9) - d(8, 1) = 59 - 1 = 58.$$

Thus, the value of $(11\ 8)$ is non-negative.

TRIAL 2. $\text{MIN}(M)(11, 2) = 2$.

$$d(11, 2) - d(11, 9) = 14 - 32 = -18.$$

$$(11, 2) \rightarrow (11, 20)$$

$$(20, 2) \rightarrow (20, 20)$$

$(20, 2)$ is an arc of D_7 .

$$(20, 15) \rightarrow (20, 18)$$

$$d(20, 15) - d(20, 2) = 6 - 6 = 0.$$

$$(18, 15) \rightarrow (18, 18)$$

$(18, 15)$ is an arc of D_7 .

$$(18, 14) \rightarrow (18, 6)$$

$$d(18, 14) - d(18, 15) = 9 - 3 = 6.$$

$$(6, 14) \rightarrow (6, 6)$$

$(6, 14)$ is an arc of D_7 .

$$(6, 2) \rightarrow (6, 20)$$

$$d(6, 2) - d(6, 14) = 19 - 18 = 1.$$

$$P = [11, 20, 18, 6, 20]$$

$$d(6, 9) - d(6, 14) = 72 - 18 = 54.$$

$$s_{211} = (11^{-18} 20^0 18^6 6^{54}).$$

The value of s_{211} is non-negative.

$$d(20, 9) - d(20, 2) = 35 - 6 = 29.$$

$$d(18, 9) - d(18, 15) = 83 - 3 = 80.$$

$$s_{211} = (11^{-21} 20^{29}), \quad s_{212} = (11^{-21} 20^0 18^{80}).$$

The value of both of the above permutations are non-negative.

$$s_{22} = (20^0 18^6 6^1).$$

The value of s_{22} is non-negative.

$$\text{TRIAL 3. MIN}(M)(11, 3) = 4.$$

$$d(11, 4) - d(11, 9) = 18 - 32 = -14.$$

$$(11, 4) \rightarrow (11, 9)$$

$$(9, 15) \rightarrow (9, 18)$$

$$d(9, 15) - d(9, 4) = 4 - 6 = -2.$$

$$(18, 15) \rightarrow (18, 18)$$

$(18, 15)$ is an arc of D_7 .

$$(18, 14) \rightarrow (18, 6)$$

$$d(18, 14) - d(18, 15) = 9 - 3 = 6.$$

$$(6, 14) \rightarrow (6, 6)$$

$(6, 14)$ is an arc of D_7 .

$$(6, 2) \rightarrow (6, 20)$$

$$d(6, 2) - d(6, 14) = 19 - 18 = 1.$$

$$(20, 2) \rightarrow (20, 20)$$

$(20, 2)$ is an arc of D_7 .

$$(20, 15) \rightarrow (20, 18)$$

$$d(20, 15) - d(20, 2) = 6 - 6 = 0.$$

$$P = [11, 9, 18, 6, 20, 18].$$

$$d(20, 9) - d(20, 2) = 35 - 6 = 29.$$

$$s_{3I} = (11^{-14}9^{-2}18^66^120^{29}).$$

The value of s_{3I} is non-negative.

$d(9, 9)$ is not permissible.

$$d(18, 9) - d(18, 15) = 83 - 3 = 80.$$

$$d(6, 9) - d(6, 14) = 72 - 18 = 54.$$

$$s_{3I2} = (11^{-14}9^{-2}18^{80}), \quad s_{3I3} = (11^{-14}9^{-2}18^66^{54}).$$

Each of the above permutations has a non-negative value.

$$s_{32} = (18^6 6^1 20^0).$$

The value of s_{32} is non-negative.

Since we were unable to obtain a negatively-valued permutation in all three trials, we can go no further using Phase 1 of our algorithm.

$$D_7^{-1}M^{-}$$

7 8 11 17 18 14 5 1 4 12 9 20 19 13 16 6 10 15 3 2

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
1	0	50	43	-7	63	8	5	∞	88	42	30	56	77	20	56	29	37	39	63	79	1
2	79	0	4	34	42	25	47	60	52	56	38	25	5	12	42	-2	46	-9	8	∞	2
3	60	63	0	43	19	81	21	42	69	20	64	68	61	10	74	13	4	63	∞	27	3
4	85	79	29	0	58	53	83	20	∞	50	78	22	84	85	9	54	9	24	68	67	4
5	2	45	22	-9	0	-15	∞	-2	61	-1	58	16	80	72	10	30	35	16	41	13	5
6	80	76	12	51	52	0	1	3	57	25	54	53	4	30	64	∞	22	76	13	1	6
7	∞	70	77	87	91	29	0	-3	64	55	24	85	54	20	6	31	40	5	78	95	7
8	53	∞	24	23	77	64	51	0	48	47	58	69	66	55	58	29	91	33	70	48	8
9	60	13	84	71	62	15	57	86	0	18	∞	60	78	86	37	73	57	-2	43	93	9
10	82	82	24	63	89	53	70	36	-6	0	62	-5	75	76	30	44	∞	53	-8	55	10
11	14	19	∞	22	53	53	59	-21	-14	58	0	-13	-1	-2	34	5	-9	-14	49	-18	11
12	72	82	70	61	58	35	-1	37	57	∞	32	0	18	71	55	32	37	40	43	-6	12
13	8	74	38	9	60	47	13	12	69	60	8	-3	0	∞	85	73	80	2	86	75	13
14	12	33	4	-13	76	∞	37	65	71	35	9	7	40	0	18	77	7	72	70	54	14
15	44	63	80	61	21	26	47	66	24	34	72	73	32	41	0	36	62	∞	47	35	15
16	83	12	57	64	56	15	76	8	55	7	53	52	15	54	∞	0	43	43	72	0	16
17	82	23	18	∞	70	27	7	45	58	74	27	5	30	16	74	82	0	81	24	47	17
18	64	63	79	65	∞	6	23	28	95	87	80	38	48	39	23	77	77	0	81	47	18
19	20	48	57	26	59	55	52	47	38	57	58	62	∞	81	51	42	82	36	0	83	19
20	8	11	18	85	79	41	43	37	45	74	29	∞	78	59	49	46	12	0	67	0	20
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	

MIN($D_7^+ M^-$)

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
1	4N	7	6	14	16	11	17	18	10	3	2	12	15	5	19	13	20	9	8	1	1
2	18N	16N	3	13	19	14	6	12	4	11	5	15	17	7	9	10	8	1	20	2	2
3	17	14	16	5	10	7	20	8	4	1	13	2	18	11	12	9	15	6	19	3	3
4	15	17	8	12	18	3	10	6	16	5	20	19	11	2	7	13	1	14	9	4	4
5	6N	4N	8N	10N	1	15	20	12	18	3	16	17	19	2	11	9	14	13	7	5	5
6	7	20	8	13	3	19	17	10	14	4	5	12	11	9	15	2	18	1	16	6	6
7	8N	18	15	14	11	6	16	17	13	10	9	2	3	19	12	4	5	20	1	7	7
8	4	3	16	18	10	9	20	7	1	14	15	11	6	13	12	19	5	17	2	8	8
9	18N	2	6	10	15	19	17	7	1	12	5	4	16	13	3	8	14	20	11	9	9
10	19N	9N	12N	3	15	8	16	6	18	20	11	4	7	13	14	1	2	5	17	10	10
11	8N	20N	9N	18N	12N	17N	14N	13N	16	1	2	4	15	19	5	6	7	10	3	11	11
12	20N	7N	13	11	16	6	8	17	18	19	15	9	5	4	3	14	1	2	10	12	12
13	12N	18	1	11	9	8	7	3	6	5	10	9	16	2	20	17	15	19	14	13	13
14	4N	3	12	17	11	1	15	2	10	7	13	20	8	19	9	18	5	16	6	14	14
15	5	6	9	13	10	20	16	14	1	7	19	4	2	17	8	11	12	3	18	15	15
16	20	10	8	7	3	6	13	17	18	12	11	9	5	3	4	19	7	1	15	16	16
17	12	7	14	3	2	19	6	13	11	8	20	9	5	10	15	18	1	16	4	17	17
18	6	7	15	8	12	14	20	13	2	1	4	16	17	3	11	19	10	9	5	18	18
19	1	4	18	9	16	8	2	15	7	6	3	10	11	5	12	14	17	20	13	19	19
20	18	1	2	17	3	11	8	6	7	9	16	15	14	19	10	13	5	4	12	20	20
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	

$$j = 4$$

$$(14\ 4)(4\ 15) = (14\ 15): -4$$

$$(14\ 4)(4\ 17) = (14\ 17): -4$$

$$j = 6$$

$$(5\ 6)(6\ 3) = (5\ 3): -3$$

$$(5\ 6)(6\ 7) = (5\ 7): -14$$

$$(5\ 6)(6\ 8) = (5\ 8): -12$$

$$(5\ 6)(6\ 13) = (5\ 13): -11$$

$$(5\ 6)(6\ 19) = (5\ 19): -2$$

$$(5\ 6)(6\ 20) = (5\ 20): -14$$

$$j = 7$$

$$(5\ 7)(7\ 8) = (5\ 8): -17$$

$$(5\ 7)(7\ 15) = (5\ 15): -8$$

$$(5\ 7)(7\ 18) = (5\ 18): -9$$

$$j = 10$$

$$(5\ 10)(10\ 9) = (5\ 9): -7$$

$$j = 12$$

$$(10\ 12)(12\ 7) = (10\ 7): -6$$

$$(10\ 12)(12\ 20) = (10\ 20): -11$$

$$(11\ 12)(12\ 7) = (11\ 7): -14$$

$$(11\ 12)(12\ 20) = (11\ 20): -19$$

$$(13\ 12)(12\ 7) = (13\ 7): -4$$

$$(13\ 12)(12\ 20) = (13\ 20): -9$$

$$j = 13$$

$$(5\ 13)(13\ 1) = (5\ 1): -3$$

$$j = 14$$

$$(11\ 14)(14\ 4) = (11\ 4): -15$$

$$j = 16$$

$$(2\ 16)(16\ 20) = (2\ 20): -2$$

$$j = 18$$

$$(2\ 18)(18\ 6) = (2\ 6): -3$$

$$(10\ 18)(18\ 6) = (10\ 6): -2$$

$$(11\ 18)(18\ 6) = (11\ 6): -10$$

$$j = 20$$

$$(5\ 20)(20\ 1) = (5\ 1): -6$$

$$(5\ 20)(20\ 2) = (5\ 2): -3$$

$$(5\ 20)(20\ 17) = (5\ 17): -2$$

$$(5\ 20)(20\ 18) = (5\ 18): -14$$

$$(11\ 20)(20\ 18) = (11\ 18): -19$$

$$(12\ 20)(20\ 18) = (13\ 18): -9$$

$$D_7^{-1}M^-(20)$$

7 8 11 17 18 14 5 1 4 12 9 20 19 13 16 6 10 15 3 2

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
1	0	50	43	-7	63	8	5	∞	88	42	30	56	77	20	56	29	37	39	63	79	1
2	79	0	4	34	42	<u>-3</u>	47	60	52	56	38	25	5	12	42	-2	46	-9	8	-2	2
3	60	63	0	43	19	81	21	42	69	20	64	68	61	10	74	13	4	63	∞	27	3
4	85	79	29	0	58	53	83	20	∞	50	78	22	84	85	9	54	9	24	68	67	4
5	<u>-6</u>	<u>-3</u>	<u>-3</u>	-9	0	<u>-15</u>	<u>-14</u>	<u>-17</u>	<u>-7</u>	<u>-1</u>	58	16	<u>-11</u>	72	-8	30	<u>-2</u>	<u>-14</u>	-2	<u>-14</u>	5
6	80	76	12	51	52	0	1	3	57	25	54	53	4	30	64	∞	22	76	13	1	6
7	∞	70	77	87	91	29	0	-3	64	55	24	85	54	20	6	31	40	5	78	95	7
8	53	∞	24	23	77	64	51	0	48	47	58	69	66	55	58	29	91	33	70	48	8
9	60	13	84	71	62	15	57	86	0	18	∞	60	78	86	37	73	57	-2	43	93	9
10	82	82	24	63	89	<u>-2</u>	<u>-6</u>	36	-6	0	62	-5	75	76	30	44	∞	-8	-8	<u>-11</u>	10
11	14	<u>-1</u>	∞	<u>-15</u>	53	<u>-10</u>	<u>-14</u>	<u>-21</u>	<u>-14</u>	58	0	<u>-13</u>	<u>-1</u>	-2	34	5	-9	<u>-19</u>	49	<u>-19</u>	11
12	72	82	70	61	58	35	-	-4	57	∞	32	0	18	71	55	32	37	<u>-6</u>	43	-6	12
13	8	74	38	9	60	47	<u>-4</u>	12	69	60	8	-3	0	∞	-4	73	-4	<u>-9</u>	86	-9	13
14	12	33	4	<u>-13</u>	76	∞	37	65	71	35	9	7	40	0	-4	77	7	-4	70	54	14
15	44	63	80	61	21	26	47	66	24	34	72	73	32	41	0	36	62	∞	47	35	15
16	83	12	57	64	56	15	76	8	55	7	53	52	15	54	∞	0	43	43	72	0	16
17	82	23	18	∞	70	27	7	45	58	74	27	5	30	16	74	82	0	81	24	47	17
18	64	63	79	65	∞	6	23	28	95	87	80	38	48	39	23	77	77	0	81	47	18
19	20	48	57	26	59	55	52	47	38	57	58	62	∞	81	51	42	82	36	0	83	19
20	8	11	18	85	79	41	43	37	45	74	29	∞	78	59	49	46	12	0	67	0	20
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	

P_{20}

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
1																					1
2						18										2		2		16	2
3																					3
4																					4
5	20	20	6			5	6	7	10	5			6		7		20	20	6	6	5
6																					6
7																					7
8																					8
9																					9
10						18	12		10			10						9		12	10
11		9		14		18	12		11			11		11				20		12	11
12																					12
13							12					13						20		12	13
14				14											4		4				14
15																					15
16																					16
17																					17
18																					18
19																					19
20																					20
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	

NEGPATHS(20)

(1) (5 1); (2) (5 2), (7 2), (11 2); (3) (5 3); (4) (11 4); (6) (2 6), (10 6), (11 6);
 (7) (10 7), (11 7), (13 7); (9) (5 9); (17) (5 17); (18) (5 18), (11 18), (12 18), (13 18).

$j = 1$

$$(5 \ 1)(1 \ 4) = (5 \ 4): -13$$

$j = 2$

$$(5 \ 2)(2 \ 16) = (5 \ 16): -5$$

$$(12 \ 2)(2 \ 16) = (11 \ 16): -3$$

$j = 4$

$$(5 \ 4)(4 \ 17) = (5 \ 17): -4$$

$j = 6$

$$(2 \ 6)(6 \ 7) = (2 \ 7): -2$$

$$(11 \ 6)(6 \ 13) = (11 \ 13): -9$$

$j = 7$

$$(2 \ 7)(7 \ 8) = (2 \ 8): -5$$

$$(10 \ 7)(7 \ 8) = (10 \ 8): -9$$

$j = 13$

$$(11 \ 13)(13 \ 1) = (11 \ 1): -1$$

$$(11 \ 13)(13 \ 11) = (11 \ 11): -1$$

We thus have obtained a negative cycle using column 13 and need continue no further in the construction of $D_7^{-1}M^-(40)$.

$$D_7^{-1}M^-(40)$$

7 8 11 17 18 14 5 1 4 12 9 20 19 13 16 6 10 15 3 2

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
1	0	50	43	-7	63	8	5	∞	88	42	30	56	77	20	56	29	37	39	63	79	1
2	79	0	4	34	42	-3	-2	60	52	56	38	25	5	12	42	-2	46	-9	8	-2	2
3	60	63	0	43	19	81	21	42	69	20	64	68	61	10	74	13	4	63	∞	27	3
4	85	79	29	0	58	53	83	20	∞	50	78	22	84	85	9	54	9	24	68	67	4
5	<u>-6</u>	-3	-3	<u>-13</u>	0	<u>-15</u>	<u>-14</u>	<u>-17</u>	-7	<u>-1</u>	58	16	<u>-11</u>	72	-8	<u>-5</u>	<u>-4</u>	<u>-14</u>	-2	<u>-14</u>	5
6	80	76	12	51	52	0	1	3	57	25	54	53	4	30	64	∞	22	76	13	1	6
7	∞	70	77	87	91	29	0	-3	64	55	24	85	54	20	6	31	40	5	78	95	7
8	53	∞	24	23	77	64	51	0	48	47	58	69	66	55	58	29	91	33	70	48	8
9	60	13	84	71	62	15	57	86	0	18	∞	60	78	86	37	73	57	-2	43	93	9
10	82	82	24	63	89	-2	-6	-9	-6	0	62	-5	75	76	30	44	∞	-8	-8	<u>-11</u>	10
11	<u>-1</u>	<u>-1</u>	∞	<u>-15</u>	53	<u>-13</u>	<u>-14</u>	<u>-21</u>	<u>-14</u>	58	-1	<u>-13</u>	-9	-2	<u>-6</u>	<u>-3</u>	<u>-11</u>	<u>-19</u>	49	<u>-19</u>	11
12	72	82	70	61	58	35	-	-4	57	∞	32	0	18	71	55	32	37	<u>-6</u>	43	-6	12
13	8	74	38	9	60	47	-4	-7	69	60	8	-3	0	∞	-4	73	-4	<u>-9</u>	86	-9	13
14	12	33	4	<u>-13</u>	76	∞	37	65	71	35	9	7	40	0	-4	77	7	-4	70	54	14
15	44	63	80	61	21	26	47	66	24	34	72	73	32	41	0	36	62	∞	47	35	15
16	83	12	57	64	56	15	76	8	55	7	53	52	15	54	∞	0	43	43	72	0	16
17	82	23	18	∞	70	27	7	45	58	74	27	5	30	16	74	82	0	81	24	47	17
18	64	63	79	65	∞	6	23	28	95	87	80	38	48	39	23	77	77	0	81	47	18
19	20	48	57	26	59	55	52	47	38	57	58	62	∞	81	51	42	82	36	0	83	19
20	8	11	18	85	79	41	43	37	45	74	29	∞	78	59	49	46	12	0	67	0	20
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	

$$P_{40}$$

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
1																				
2						2	6	7												
3																				
4																				
5	5	5		1												2	4			
6																				
7																				
8																				
9																				
10							10	7												12
11	13	11				11					13		6			2				12
12																				12
13																				12
14																				
15																				
16																				
17																				
18																				
19																				
20																				
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20

From the entry -1 at $(11\ 13)$ of $D_7^{-1}M(40)$, the value of s is -1 . Using P_{40} , checking for further points of s , we obtain $(11\ 6\ 13)$. Going back to P_{20} , $18 \rightarrow 6$, $20 \rightarrow 18$, $12 \rightarrow 20$, $11 \rightarrow 12$. Thus, $s = (11\ 12\ 20\ 18\ 6\ 13)$. $D_8 = D_7s \Rightarrow D_8^{-1} = s^{-1}D_7^{-1}$. Applying s^{-1} to the second row of $D_7^{-1}M$, we obtain $D_8^{-1}M$.

$$D_8^{-1}M^{-}$$

	7	8	11	17	18	19	5	1	4	12	20	2	9	13	16	6	10	14	3	15	
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
1	0	50	43	-7	63	77	5	∞	88	41	56	79	30	20	56	29	37	8	63	39	1
2	79	0	4	34	42	5	47	60	52	56	25	∞	38	12	42	-2	46	25	8	-9	2
3	60	63	0	43	19	61	21	42	49	20	68	27	84	10	74	13	4	81	∞	63	3
4	85	79	29	0	58	84	83	20	∞	50	22	67	78	85	9	54	9	53	68	24	4
5	2	45	22	-9	0	80	∞	-2	61	-1	16	13	58	72	10	30	35	-15	41	16	5
6	76	72	8	47	48	0	-3	-1	53	21	49	-3	50	26	60	∞	18	-2	9	72	6
7	∞	70	77	87	91	54	0	-3	64	55	85	95	24	20	6	31	40	29	78	5	7
8	51	∞	24	23	77	66	51	0	48	47	69	48	58	55	58	29	91	64	70	33	8
9	60	13	84	71	62	78	57	86	0	18	60	93	∞	86	37	73	57	15	43	-2	9
10	82	82	24	63	89	75	70	36	-6	0	-5	55	62	76	30	44	∞	53	-8	53	10
11	27	32	∞	35	66	12	72	-8	-1	71	0	-5	13	11	47	18	4	66	62	-1	11
12	78	88	76	67	64	24	5	42	63	∞	6	0	38	77	61	38	43	41	51	44	12
13	0	66	30	1	52	-8	5	4	61	52	-11	67	0	∞	77	65	72	39	78	-6	13
14	12	33	4	-15	76	40	37	65	71	35	7	54	9	0	18	77	7	∞	70	72	14
15	44	63	80	61	21	32	47	66	24	34	73	35	72	41	0	36	62	26	47	∞	15
16	83	12	57	64	56	15	76	8	55	7	52	0	53	54	∞	0	43	15	72	43	16
17	82	23	18	∞	70	30	7	45	58	74	5	47	27	16	74	82	0	27	24	81	17
18	58	42	73	59	∞	42	17	22	89	81	32	41	74	33	17	71	71	0	75	-6	18
19	20	11	57	26	53	∞	52	47	38	57	62	83	58	81	51	42	82	55	0	-6	19
20	8	50	18	85	79	78	43	37	45	74	∞	0	29	59	49	46	12	41	67	0	20
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	

$\text{MIN}(D_8^{-1}M^-)$

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
1	4	7	18	14	16	30	17	20	10	3	2	11	15	5	19	6	12	9	8	1	1
2	20N	16N	3	6	19	14	11	18	4	13	5	15	17	9	4	10	8	1	12	2	2
3	17	14	16	5	10	7	12	8	4	9	1	6	2	20	11	15	18	13	19	3	3
4	15	17	8	11	20	3	10	18	16	5	12	19	13	2	7	6	14	1	9	4	4
5	18N	4N	8N	10N	1	15	12	11	20	3	16	17	19	2	13	9	14	6	7	5	5
6	7N	12N	18N	8N	3	19	17	10	14	4	5	11	13	9	15	2	20	1	16	6	6
7	8N	20	15	14	13	18	16	17	6	10	9	2	3	19	11	4	5	12	1	7	7
8	4	3	16	20	10	9	12	1	7	14	13	15	18	6	11	19	5	17	2	8	8
9	20N	2	18	10	15	19	7	17	11	1	5	4	16	6	3	8	14	12	13	9	9
10	19N	9N	11N	3	15	8	16	18	20	12	13	4	7	6	14	1	2	5	17	10	10
11	8N	12N	5N	20N	17	14	6	13	16	1	2	4	15	19	5	18	19	7	3	11	11
12	7	11	6	13	16	18	8	17	20	19	15	9	5	4	3	14	1	2	10	12	12
13	11	6	20	13	4	8	7	3	18	5	10	9	16	2	12	17	15	19	14	13	13
14	4	3	11	17	13	1	15	2	10	7	6	12	8	19	9	20	5	16	18	14	14
15	5	9	18	6	10	12	16	14	1	7	19	4	17	2	8	13	11	3	20	15	15
16	12	10	8	2	6	18	17	20	11	13	14	9	5	3	4	19	7	1	15	16	16
17	11	7	14	18	2	19	13	18	6	8	12	9	5	10	15	20	1	16	4	17	17
18	20	7	15	8	11	14	12	2	6	1	4	16	17	3	13	19	10	9	5	18	18
19	20	2	1	4	9	16	8	15	1	5	18	3	10	13	11	14	17	12	6	19	19
20	12	1	17	3	13	8	18	7	9	16	15	2	14	19	10	5	6	4	11	20	20
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	

j = 4

$$(14\ 4)(4\ 15) = (14\ 15): -6$$

$$(14\ 4)(4\ 17) = (14\ 17): -6$$

j = 6

$$(13\ 6)(6\ 7) = (13\ 7): -17$$

$$(13\ 6)(6\ 8) = (13\ 8): -9$$

$$(13\ 6)(6\ 12) = (13\ 12): -11$$

$$(13\ 6)(6\ 18) = (13\ 18): -10$$

j = 7

$$(6\ 7)(7\ 8) = (6\ 8): -6$$

$$(13\ 7)(7\ 8) = (13\ 8): -20$$

$$(13\ 7)(7\ 15) = (13\ 15): -11$$

$$(13\ 7)(7\ 20) = (13\ 20): -12$$

j = 9

$$(10\ 9)(9\ 20) = (10\ 20): -8$$

$$(11\ 9)(9\ 20) = (11\ 20): -3$$

$$(10\ 9)(9\ 20) = (10\ 20): -8$$

j = 10

$$(5\ 10)(10\ 9) = (5\ 9): -7$$

$$(5\ 10)(10\ 11) = (5\ 11): -6$$

$$(5\ 10)(10\ 19) = (5\ 19): -9$$

j = 11

$$(5\ 11)(11\ 8) = (5\ 8): -14$$

$$(5\ 11)(11\ 9) = (5\ 9): -7$$

$$(5\ 11)(11\ 12) = (5\ 12): -11$$

$$(5\ 11)(11\ 17) = (5\ 17): -2$$

$$(5 \ 11)(11 \ 20) = (5 \ 20): -7$$

$$(10 \ 11)(11 \ 12) = (10 \ 12): -10$$

$$(10 \ 11)(11 \ 17) = (10 \ 17): -1$$

$$(13 \ 11)(11 \ 9) = (13 \ 9): -12$$

$$(13 \ 11)(11 \ 12) = (13 \ 12): -16$$

$$(13 \ 11)(11 \ 17) = (13 \ 17): -7$$

$$(13 \ 11)(11 \ 20) = (13 \ 20): -12$$

$$j = 12$$

$$(5 \ 12)(12 \ 7) = (5 \ 7): -6$$

$$(10 \ 12)(12 \ 7) = (10 \ 7): -5$$

$$j = 18$$

$$(5 \ 18)(18 \ 20) = (5 \ 20): -21$$

$$(13 \ 18)(18 \ 20) = (13 \ 20): -16$$

$$j = 20$$

$$(2 \ 20)(20 \ 1) = (2 \ 1): -1$$

$$(2 \ 20)(20 \ 12) = (2 \ 12): -9$$

$$(5 \ 20)(20 \ 1) = (5 \ 1): -13$$

$$(5 \ 20)(20 \ 3) = (5 \ 3): -3$$

$$(5 \ 20)(20 \ 12) = (5 \ 12): -21$$

$$(5 \ 20)(20 \ 17) = (5 \ 17): -9$$

$$(10 \ 20)(20 \ 1) = (10 \ 1): -6$$

$$(10 \ 20)(20 \ 12) = (10 \ 12): -8$$

$$(11 \ 20)(20 \ 12) = (11 \ 12): -3$$

$$(13 \ 20)(20 \ 1) = (13 \ 1): -6$$

$$(13 \ 20)(20 \ 12) = (13 \ 12): -14$$

$$(13 \ 20)(20 \ 17) = (13 \ 17): -2$$

$$(18\ 20)(20\ 12) = (18\ 12): -6$$

$$(19\ 20)(20\ 12) = (19\ 12): -6$$

$$D_8^{-I}M^-(20)$$

	7	8	11	17	18	19	5	1	4	12	20	2	9	13	16	6	10	14	3	15	
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
1	0	50	43	-7	63	77	5	∞	88	41	56	79	30	20	56	29	37	8	63	39	1
2	<u>-1</u>	0	4	34	42	5	47	60	52	56	25	<u>-9</u>	38	12	42	-2	46	25	8	-9	2
3	60	63	0	43	19	61	21	42	49	20	68	27	84	10	74	13	4	81	∞	63	3
4	85	79	29	0	58	84	83	20	∞	50	22	67	78	85	9	54	9	53	68	24	4
5	<u>-13</u>	45	<u>-3</u>	-9	0	80	<u>-6</u>	<u>-14</u>	<u>-7</u>	<u>-7</u>	-6	<u>-21</u>	58	72	10	30	<u>-9</u>	<u>-15</u>	-9	<u>-21</u>	5
6	76	72	8	47	48	0	-3	<u>-6</u>	53	21	49	-3	50	26	60	∞	18	-2	9	72	6
7	∞	70	77	87	91	54	0	-3	64	55	85	95	24	20	6	31	40	29	78	5	7
8	51	∞	24	23	77	66	51	0	48	47	69	48	58	55	58	29	91	64	70	33	8
9	60	13	84	71	62	78	57	86	0	18	60	93	∞	86	37	73	57	15	43	-2	9
10	<u>-6</u>	82	24	63	89	75	<u>-5</u>	36	-6	0	-5	<u>-14</u>	62	76	30	44	<u>-2</u>	53	-8	<u>-14</u>	10
11	27	32	∞	35	66	12	72	-8	<u>-1</u>	71	0	-5	13	11	47	18	4	66	62	-3	11
12	78	88	76	67	64	24	5	42	63	∞	6	0	38	77	61	38	43	41	51	44	12
13	<u>-8</u>	66	30	1	52	-8	<u>-17</u>	<u>-20</u>	<u>-12</u>	52	<u>-11</u>	<u>-16</u>	0	∞	<u>-11</u>	65	<u>-4</u>	<u>-10</u>	78	<u>-16</u>	13
14	12	33	4	<u>-15</u>	76	40	37	65	71	35	<u>-1</u>	54	9	0	-6	77	-6	∞	70	72	14
15	44	63	80	61	21	32	47	66	24	34	73	35	72	41	0	36	62	26	47	∞	15
16	83	12	57	64	56	15	76	8	55	7	52	0	53	54	∞	0	43	15	72	43	16
17	82	23	18	∞	70	30	7	45	58	74	5	47	27	16	74	82	0	27	24	81	17
18	58	42	73	59	∞	42	17	22	89	81	32	<u>-6</u>	74	33	17	71	71	0	75	-6	18
19	20	11	57	26	53	∞	52	47	38	57	62	<u>-6</u>	58	81	51	42	82	55	0	-6	19
20	8	50	18	85	79	78	43	37	45	74	∞	0	29	59	49	46	12	41	67	0	20
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	

$$P_{20}$$

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
1																					1
2	2			1			12				12	2									2
3																					3
4																					4
5	5	19					12	11	10	5	12	5			7				10		5
6																					6
7																					7
8																					8
9																					9
10	10			1			12	10	7		12	10			4		4				10
11																					11
12																					12
13	13			1													4				13
14				14				11			14				4		4				14
15																					15
16																					16
17																					17
18																					18
19																					19
20																					20
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	

NEGPATHS(20)

(1) (2 1), (5 1), (10 1), (13 1); (3) (5 3); (7) (5 7), (10 7); (8) (5 8), (6 8), (9) (5 9), (13 9); (10) (5 10); (11) (14 11); (12) (2 12), (5 12), (10 12), (13 12), (18 12), (19 12); (17) (5 17), (10 17), (13 17).

j = 1

$$(2\ 1)(1\ 4) = (2\ 4): -8$$

$$(5\ 1)(1\ 4) = (5\ 4): -20$$

$$(5\ 1)(1\ 7) = (5\ 7): -7$$

$$(10\ 1)(1\ 4) = (10\ 4): -13$$

$$(13\ 1)(1\ 4) = (13\ 4): -15$$

j = 4

$$(5\ 4)(4\ 15) = (5\ 15): -11$$

$$(5\ 4)(4\ 17) = (5\ 17): -11$$

$$(10\ 4)(4\ 15) = (10\ 15): -4$$

$$(10\ 4)(4\ 17) = (10\ 17): -4$$

$$(12\ 4)(4\ 17) = (13\ 17): -6$$

j = 7

$$(10\ 7)(7\ 8) = (10\ 8): -8$$

j = 10

$$(5\ 10)(10\ 9) = (5\ 9): -13$$

$$(5\ 10)(10\ 11) = (5\ 11): -12$$

$$(5\ 10)(10\ 19) = (5\ 19): -16$$

j = 11

$$(5\ 11)(11\ 8) = (5\ 8): -20$$

$$(5\ 11)(11\ 14) = (5\ 14): -1$$

$$(13\ 11)(11\ 9) = (14\ 9): -2$$

$$j = 12$$

$$(5 \ 12)(12 \ 7) = (5 \ 7): -16$$

$$(5 \ 12)(12 \ 11) = (5 \ 11): -15$$

$$(10 \ 12)(12 \ 7) = (10 \ 7): -9$$

$$(10 \ 12)(12 \ 11) = (10 \ 11): -8$$

$$j = 19$$

$$(5 \ 19)(19 \ 2) = (5 \ 2): -5$$

$$(5 \ 19)(19 \ 20) = (5 \ 19)(19 \ 20): -5$$

$$j = 20$$

$$(5 \ 20)(20 \ 1) = (5 \ 1): -14$$

$$(5 \ 20)(20 \ 12) = (5 \ 12): -22$$

$$D_8^{-1}M^-(40)$$

	7	8	11	17	18	19	5	1	4	12	20	2	9	13	16	6	10	14	3	15	
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
1	0	50	43	-7	63	77	5	∞	88	41	56	79	30	20	56	29	37	8	63	39	1
2	-1	0	4	<u>-8</u>	42	5	-4	60	52	56	-3	-9	38	12	42	-2	46	25	8	-9	2
3	60	63	0	43	19	61	21	42	49	20	68	27	84	10	74	13	4	81	∞	63	3
4	85	79	29	0	58	84	83	20	∞	50	22	67	78	85	9	54	9	53	68	24	4
5	<u>-14</u>	<u>-5</u>	-3	-20	0	80	<u>-16</u>	<u>-20</u>	<u>-13</u>	-7	<u>-16</u>	<u>-22</u>	58	-1	-11	30	-11	-15	-16	-22	5
6	76	72	8	47	48	0	-3	-6	53	21	49	-3	50	26	60	∞	18	-2	9	72	6
7	∞	70	77	87	91	54	0	-3	64	55	85	95	24	20	6	31	40	29	78	5	7
8	51	∞	24	23	77	66	51	0	48	47	69	48	58	55	58	29	91	64	70	33	8
9	60	13	84	71	62	78	57	86	0	18	60	93	∞	86	37	73	57	15	43	-2	9
10	-6	82	24	<u>-13</u>	89	75	<u>-9</u>	-8	-6	0	<u>-8</u>	<u>-14</u>	62	76	-4	44	-4	53	-8	<u>-14</u>	10
11	27	32	∞	35	66	12	72	-8	-1	71	0	-5	13	11	47	18	4	66	62	-3	11
12	78	88	76	67	64	24	5	42	63	∞	6	0	38	77	61	38	43	41	51	44	12
13	-8	66	30	<u>-15</u>	52	-8	<u>-17</u>	<u>-20</u>	<u>-12</u>	52	<u>-11</u>	<u>-16</u>	0	∞	<u>-11</u>	65	-6	<u>-10</u>	78	<u>-16</u>	13
14	12	33	4	<u>-15</u>	76	40	37	65	<u>-2</u>	35	-1	54	9	0	-6	77	-6	∞	70	72	14
15	44	63	80	61	21	32	47	66	24	34	73	35	72	41	0	36	62	26	47	∞	15
16	83	12	57	64	56	15	76	8	55	7	52	0	53	54	∞	0	43	15	72	43	16
17	82	23	18	∞	70	30	7	45	58	74	5	47	27	16	74	82	0	27	24	81	17
18	58	42	73	59	∞	42	17	22	89	81	32	-6	74	33	17	71	71	0	75	-6	18
19	20	11	57	26	53	∞	52	47	38	57	62	-6	58	81	51	42	82	55	0	-6	19
20	8	50	18	85	79	78	43	37	45	74	∞	0	29	59	49	46	12	41	67	0	20
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	

$$P_{40}$$

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
1																					1
2	2			1																	2
3																					3
4																					4
5	20	19		1			12	11	10	5	12	20		11	4		4		10	19	5
6																					6
7																					7
8																					8
9																					9
10	10			1			10	7			12	10			4		4				10
11							12					10									11
12																					12
13	13			1													4				13
14									11		14										14
15																					15
16																					16
17																					17
18																					18
19																					19
20																					20
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	

NEGPATHS(40)

(1) (5 1); (2) (5 2); (4) (2 4); (7) (5 7), (10 7); (8) (5 8); (9) (5 9), (14 9); (11) (5 11), (10 11); (12) (5 12).

j = 1

$$(5\ 1)(1\ 4) = (5\ 4): -21$$

j = 2

$$(5\ 2)(2\ 16) = (5\ 16): -7$$

j = 4

$$(5\ 4)(4\ 15) = (5\ 15): -12$$

j = 7

$$(10\ 7)(7\ 8) = (10\ 8): -12$$

j = 9

$$(14\ 9)(9\ 20) = (14\ 20): -4$$

j = 11

$$(5\ 11)(11\ 8) = (5\ 8): -24$$

$$(5\ 11)(11\ 9) = (5\ 9): -17$$

$$(10\ 11)(11\ 8) = (10\ 8): -16$$

$$(10\ 11)(11\ 9) = (10\ 9): -9$$

j = 12

$$(5\ 12)(12\ 7): -17$$

j = 20

$$(14\ 20)(20\ 12): -4$$

$$D_8^{-1}M^-(60)$$

	7	8	11	17	18	19	5	1	4	12	20	2	9	13	16	6	10	14	3	15	
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
1	0	50	43	-7	63	77	5	∞	88	41	56	79	30	20	56	29	37	8	63	39	1
2	-1	0	4	-8	42	5	-4	60	52	56	-3	-9	38	12	42	-2	46	25	8	-9	2
3	60	63	0	43	19	61	21	42	49	20	68	27	84	10	74	13	4	81	∞	63	3
4	85	79	29	0	58	84	83	20	∞	50	22	67	78	85	9	54	9	53	68	24	4
5	-14	-5	-3	-21	0	80	-17	-24	-17	-7	-16	-22	58	-1	-12	-7	-12	-15	-16	-22	5
6	76	72	8	47	48	0	-3	-6	53	21	49	-3	50	26	60	∞	18	-2	9	72	6
7	∞	70	77	87	91	54	0	-3	64	55	85	95	24	20	6	31	40	29	78	5	7
8	51	∞	24	23	77	66	51	0	48	47	69	48	58	55	58	29	91	64	70	33	8
9	60	13	84	71	62	78	57	86	0	18	60	93	∞	86	37	73	57	15	43	-2	9
10	-6	82	24	-13	89	75	-9	-12	-9	0	-8	-14	62	76	-4	44	-4	53	-8	-14	10
11	27	32	∞	35	66	12	72	-8	-1	71	0	-5	13	11	47	18	4	66	62	-3	11
12	78	88	76	67	64	24	5	42	63	∞	6	0	38	77	61	38	43	41	51	44	12
13	-8	66	30	-15	52	-8	-17	-20	-12	52	-11	-16	0	∞	-11	65	-6	-10	78	-16	13
14	12	33	4	-15	76	40	37	65	-2	35	-1	-4	9	0	-6	77	-6	∞	70	-4	14
15	44	63	80	61	21	32	47	66	24	34	73	35	72	41	0	36	62	26	47	∞	15
16	83	12	57	64	56	15	76	8	55	7	52	0	53	54	∞	0	43	15	72	43	16
17	82	23	18	∞	70	30	7	45	58	74	5	47	27	16	74	82	0	27	24	81	17
18	58	42	73	59	∞	42	17	22	89	81	32	-6	74	33	17	71	71	0	75	-6	18
19	20	11	57	26	53	∞	52	47	38	57	62	-6	58	81	51	42	82	55	0	-6	19
20	8	50	18	85	79	78	43	37	45	74	∞	0	29	59	49	46	12	41	67	0	20
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	

We can't extend any path further as shown in $D_8^{-1}M^-(80)$. It follows that $\sigma_{APOPT} = D_8$.

$$P_{60}$$

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
1																					1
2																					2
3																					3
4																					4
5	5	5		1			12	11	11		5	5			4	2	4				5
6																					6
7																					7
8																					8
9																					9
10							10	11	11		10										10
11																					11
12																					12
13																					13
14									14			20								9	14
15																					15
16																					16
17																					17
18																					18
19																					19
20																					20
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	

$$D_8^{-I}M^-(80)$$

	7	8	11	17	18	19	5	1	4	12	20	2	9	13	16	6	10	14	3	15	
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
1	0	50	43	-7	63	77	5	∞	88	41	56	79	30	20	56	29	37	8	63	39	1
2	-1	0	4	-8	42	5	-4	60	52	56	-3	-9	38	12	42	-2	46	25	8	-9	2
3	60	63	0	43	19	61	21	42	49	20	68	27	84	10	74	13	4	81	∞	63	3
4	85	79	29	0	58	84	83	20	∞	50	22	67	78	85	9	54	9	53	68	24	4
5	-14	-5	-3	-21	0	80	-17	-24	-17	-7	-16	-22	58	-1	-12	-7	-12	-15	-16	-22	5
6	76	72	8	47	48	0	-3	-6	53	21	49	-3	50	26	60	∞	18	-2	9	72	6
7	∞	70	77	87	91	54	0	-3	64	55	85	95	24	20	6	31	40	29	78	5	7
8	51	∞	24	23	77	66	51	0	48	47	69	48	58	55	58	29	91	64	70	33	8
9	60	13	84	71	62	78	57	86	0	18	60	93	∞	86	37	73	57	15	43	-2	9
10	-6	82	24	-13	89	75	-9	-12	-9	0	-8	-14	62	76	-4	44	-4	53	-8	-14	10
11	27	32	∞	35	66	12	72	-8	-1	71	0	-5	13	11	47	18	4	66	62	-3	11
12	78	88	76	67	64	24	5	42	63	∞	6	0	38	77	61	38	43	41	51	44	12
13	-8	66	30	-15	52	-8	-17	-20	-12	52	-11	-16	0	∞	-11	65	-6	-10	78	-16	13
14	12	33	4	-15	76	40	37	65	-2	35	-1	-4	9	0	-6	77	-6	∞	70	-4	14
15	44	63	80	61	21	32	47	66	24	34	73	35	72	41	0	36	62	26	47	∞	15
16	83	12	57	64	56	15	76	8	55	7	52	0	53	54	∞	0	43	15	72	43	16
17	82	23	18	∞	70	30	7	45	58	74	5	47	27	16	74	82	0	27	24	81	17
18	58	42	73	59	∞	42	17	22	89	81	32	-6	74	33	17	71	71	0	75	-6	18
19	20	11	57	26	53	∞	52	47	38	57	62	-6	58	81	51	42	82	55	0	-6	19
20	8	50	18	85	79	78	43	37	45	74	∞	0	29	59	49	46	12	41	67	0	20
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	

We now construct $D_8^{-I}M$.

$$D_8^{-I}M^{-}$$

	7	8	11	17	18	19	5	1	4	12	20	2	9	13	16	6	10	14	3	15	
1	0	50	43	-7	63	77	5	∞	88	41	56	79	30	20	56	29	37	8	63	39	1
2	79	0	4	34	42	5	47	60	52	56	25	∞	38	12	42	-2	46	25	8	-9	2
3	60	63	0	43	19	61	21	42	49	20	68	27	84	10	74	13	4	81	∞	63	3
4	85	79	29	0	58	84	83	20	∞	50	22	67	78	85	9	54	9	53	68	24	4
5	2	45	22	-9	0	80	∞	-2	61	-1	16	13	58	72	10	30	35	⁻¹⁵	41	16	5
6	76	72	8	47	48	0	-3	-1	53	21	49	-3	50	26	60	∞	18	-2	9	72	6
7	∞	70	77	87	91	54	0	-3	64	55	85	95	24	20	6	31	40	29	78	5	7
8	51	∞	24	23	77	66	51	0	48	47	69	48	58	55	58	29	91	64	70	33	8
9	60	13	84	71	62	78	57	86	0	18	60	93	∞	86	37	73	57	15	43	-2	9
10	82	82	24	63	89	75	70	36	-6	0	-5	55	62	76	30	44	∞	53	-8	53	10
11	27	32	∞	35	66	12	72	-8	-1	71	0	-5	13	11	47	18	4	66	62	-1	11
12	78	88	76	67	64	24	5	42	63	∞	6	0	38	77	61	38	43	41	51	44	12
13	0	66	30	1	52	-8	5	4	61	52	⁻¹¹	67	0	∞	77	65	72	39	78	-6	13
14	12	33	4	⁻¹⁵	76	40	37	65	71	35	7	54	9	0	18	77	7	∞	70	72	14
15	44	63	80	61	21	32	47	66	24	34	73	35	72	41	0	36	62	26	47	∞	15
16	83	12	57	64	56	15	76	8	55	7	52	0	53	54	∞	0	43	15	72	43	16
17	82	23	18	∞	70	30	7	45	58	74	5	47	27	16	74	82	0	27	24	81	17
18	58	42	73	59	∞	42	17	22	89	81	32	41	74	33	17	71	71	0	75	-6	18
19	20	11	57	26	53	∞	52	47	38	57	62	83	58	81	51	42	82	55	0	-6	19
20	8	50	18	85	79	78	43	37	45	74	∞	0	29	59	49	46	12	41	67	0	20
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	

$\text{MIN}(D_8^{-I}M^-)$

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
1	4	7	18	14	16	30	17	20	10	3	2	11	15	5	19	6	12	9	8	1	1
2	20N	16N	3	6	19	14	11	18	4	13	5	15	17	9	4	10	8	1	12	2	2
3	17	14	16	5	10	7	12	8	4	9	1	6	2	20	11	15	18	13	19	3	3
4	15	17	8	11	20	3	10	18	16	5	12	19	13	2	7	6	14	1	9	4	4
5	18N	4N	8N	10N	1	15	12	11	20	3	16	17	19	2	13	9	14	6	7	5	5
6	7N	12N	18N	8N	3	19	17	10	14	4	5	11	13	9	15	2	20	1	16	6	6
7	8N	20	15	14	13	18	16	17	6	10	9	2	3	19	11	4	5	12	1	7	7
8	4	3	16	20	10	9	12	1	7	14	13	15	18	6	11	19	5	17	2	8	8
9	20N	2	18	10	15	19	7	17	11	1	5	4	16	6	3	8	14	12	13	9	9
10	19N	9N	11N	3	15	8	16	18	20	12	13	4	7	6	14	1	2	5	17	10	10
11	8N	12N	5N	20N	17	14	6	13	16	1	2	4	15	19	5	18	19	7	3	11	11
12	7	11	6	13	16	18	8	17	20	19	15	9	5	4	3	14	1	2	10	12	12
13	11	6	20	13	4	8	7	3	18	5	10	9	16	2	12	17	15	19	14	13	13
14	4	3	11	17	13	1	15	2	10	7	6	12	8	19	9	20	5	16	18	14	14
15	5	9	18	6	10	12	16	14	1	7	19	4	17	2	8	13	11	3	20	15	15
16	12	10	8	2	6	18	17	20	11	13	14	9	5	3	4	19	7	1	15	16	16
17	11	7	14	18	2	19	13	18	6	8	12	9	5	10	15	20	1	16	4	17	17
18	20	7	15	8	11	14	12	2	6	1	4	16	17	3	13	19	10	9	5	18	18
19	20	2	1	4	9	16	8	15	1	5	18	3	10	13	11	14	17	12	6	19	19
20	12	1	17	3	13	8	18	7	9	16	15	2	14	19	10	5	6	4	11	20	20
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	

$$j = 4$$

$$(5\ 4)(4\ 15) = (5\ 15): 0$$

$$(5\ 4)(4\ 17) = (5\ 17): 0$$

$$(14\ 4)(4\ 15) = (14\ 15): -6$$

$$(14\ 4)(4\ 17) = (14\ 17): -6$$

$$j = 6$$

$$(13\ 6)(6\ 3) = (13\ 3): -14$$

$$(13\ 6)(6\ 7) = (13\ 7): -11$$

$$(13\ 6)(6\ 8) = (13\ 8): -9$$

$$j = 7$$

$$(6\ 7)(7\ 8) = (6\ 8): -6$$

$$(13\ 7)(7\ 8) = (13\ 8): -14$$

$$j = 9$$

$$(11\ 9)(9\ 20) = (11\ 20): -3$$

$$j = 10$$

$$(5\ 10)(10\ 9) = (5\ 9): -7$$

$$(5\ 10)(10\ 11) = (5\ 11): -6$$

$$j = 11$$

$$(5\ 11)(11\ 8) = (5\ 8): -14$$

$$(5\ 11)(11\ 9) = (5\ 9): -7$$

$$(10\ 11)(11\ 8) = (10\ 8): -13$$

$$(10\ 11)(11\ 12) = (10\ 12): -10$$

$$(10\ 11)(11\ 17) = (10\ 17): -1$$

$$(13\ 11)(11\ 8) = (13\ 8): -19$$

$$(13\ 11)(11\ 9) = (13\ 9): -12$$

$$(13\ 11)(11\ 17) = (13\ 17): -7$$

$$(13\ 11)(11\ 20) = (13\ 20): -12$$

$$j = 12$$

$$(10\ 12)(12\ 7) = (10\ 7): -5$$

$$(11\ 12)(12\ 7) = (11\ 7): 0$$

$$j = 18$$

$$(5\ 18)(18\ 20) = (5\ 20): -21$$

$$(6\ 18)(18\ 20) = (6\ 20): -8$$

$$j = 19$$

$$(10\ 19)(19\ 20) = (10\ 20): -14$$

$$j = 20$$

$$(2\ 20)(20\ 1) = (2\ 1): -1$$

$$(2\ 20)(20\ 12) = (2\ 12): -9$$

$$(5\ 20)(20\ 1) = (5\ 1): -13$$

$$(5\ 20)(20\ 3) = (5\ 3): -3$$

$$(5\ 20)(20\ 12) = (5\ 12): -21$$

$$(5\ 20)(20\ 17) = (5\ 17): -9$$

$$(6\ 20)(20\ 1) = (6\ 1): 0$$

$$(6\ 20)(20\ 12) = (6\ 12): -8$$

$$(9\ 20)(20\ 12) = (9\ 12): -2$$

$$(10\ 20)(20\ 1) = (10\ 1): -6$$

$$(10\ 20)(20\ 12) = (10\ 12): -14$$

$$(10\ 20)(20\ 17) = (10\ 17): -2$$

$$(13\ 20)(20\ 1) = (13\ 1): -4$$

$$(13\ 20)(20\ 12) = (13\ 12): -12$$

$$(18\ 20)(20\ 12) = (18\ 12): -6$$

$$(19\ 20)(20\ 12) = (19\ 12): -6$$

$$D_8^{-1}M^-(20)$$

	7	8	11	17	18	19	5	1	4	12	20	2	9	13	16	6	10	14	3	15	
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
1	0	50	43	-7	63	77	5	∞	88	41	56	79	30	20	56	29	37	8	63	39	1
2	<u>-1</u>	0	4	34	42	5	47	60	52	56	25	<u>-9</u>	38	12	42	-2	46	25	8	-9	2
3	60	63	0	43	19	61	21	42	49	20	68	27	84	10	74	13	4	81	∞	63	3
4	85	79	29	0	58	84	83	20	∞	50	22	67	78	85	9	54	9	53	68	24	4
5	<u>-13</u>	45	<u>-3</u>	-9	0	80	∞	<u>-14</u>	<u>-7</u>	-1	-6	<u>-21</u>	58	72	0	30	<u>-9</u>	<u>-15</u>	41	<u>-21</u>	5
6	<u>0</u>	72	8	47	48	0	-3	-6	53	21	49	<u>-8</u>	50	26	60	∞	18	-2	9	-8	6
7	∞	70	77	87	91	54	0	-3	64	55	85	95	24	20	6	31	40	29	78	5	7
8	51	∞	24	23	77	66	51	0	48	47	69	48	58	55	58	29	91	64	70	33	8
9	60	13	84	71	62	78	57	86	0	18	60	<u>-2</u>	∞	86	37	73	57	15	43	-2	9
10	<u>-6</u>	82	24	63	89	75	<u>-5</u>	<u>-13</u>	-6	0	-5	<u>-14</u>	62	76	30	44	<u>-2</u>	53	-8	<u>-14</u>	10
11	27	32	∞	35	66	12	<u>0</u>	-8	-1	71	0	-5	13	11	47	18	4	66	62	-3	11
12	78	88	76	67	64	24	5	42	63	∞	6	0	38	77	61	38	43	41	51	44	12
13	<u>-4</u>	66	<u>0</u>	1	52	-8	<u>-11</u>	<u>-19</u>	<u>-12</u>	52	<u>-11</u>	<u>-12</u>	0	∞	77	65	-7	39	78	<u>-12</u>	13
14	12	33	4	<u>-15</u>	76	40	37	65	71	35	7	54	9	0	-6	77	-6	∞	70	72	14
15	44	63	80	61	21	32	47	66	24	34	73	35	72	41	0	36	62	26	47	∞	15
16	83	12	57	64	56	15	76	8	55	7	52	0	53	54	∞	0	43	15	72	43	16
17	82	23	18	∞	70	30	7	45	58	74	5	47	27	16	74	82	0	27	24	81	17
18	58	42	73	59	∞	42	17	22	89	81	32	<u>-6</u>	74	33	17	71	71	0	75	-6	18
19	20	11	57	26	53	∞	52	47	38	57	62	<u>-6</u>	58	81	51	42	82	55	0	-6	19
20	8	50	18	85	79	78	43	37	45	74	∞	0	29	59	49	46	12	41	67	0	20
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	

$$P_{20}$$

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
1																					1
2	20											20								2	2
3																					3
4																					4
5	20		20	5				11	11	5	10	20			4		20	5		18	5
6	20						6	7				20						6		18	6
7																					7
8																					8
9												20								9	9
10	20						12	11	10		10	20					20		10	19	10
11							12		11			11								9	11
12																					12
13	20		6			13	6	11	11		13	20					11			11	13
14				14											4		4				14
15																					15
16																					16
17																					17
18												20								18	18
19												20								19	19
20																					20
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	

NONNEGPATHS(20)

(1) (2 1), (10 1), (13 1); (3) (3); (4) (6 4); (7) (7 5), (10 5); (8) (5 8), (6 8); (9) (5 9), (13 9); (10) (5 10); (12) (2 12), (5 12), (10 12), (13 12), (18 12), (19 12); (17) (5 17),

(10 17), (13 17).

j = 1

$$(2 \ 1)(1 \ 4) = (2 \ 4): -8$$

$$(5 \ 1)(1 \ 4) = (5 \ 4): -20$$

$$(5 \ 1)(1 \ 7) = (5 \ 7): -8$$

$$(5 \ 1)(1 \ 18) = (5 \ 18): -5$$

$$(6 \ 1)(1 \ 4) = (6 \ 4): -7$$

$$(10 \ 1)(1 \ 4) = (10 \ 4): -13$$

$$(10 \ 1)(1 \ 7) = (10 \ 7): -1$$

$$(13 \ 1)(1 \ 4) = (13 \ 4): -11$$

j = 4

$$(5 \ 4)(4 \ 15) = (5 \ 15): -11$$

$$(5 \ 4)(4 \ 17) = (5 \ 17): -11$$

$$(10 \ 4)(4 \ 15) = (10 \ 15): -4$$

$$(10 \ 4)(4 \ 17) = (10 \ 17): -4$$

$$(13 \ 4)(4 \ 15) = (13 \ 15): -2$$

j = 9

$$(13 \ 9)(9 \ 20) = (13 \ 20): -14$$

j = 12

$$(2 \ 12)(12 \ 7) = (2 \ 7): -4$$

$$(2 \ 12)(12 \ 11) = (2 \ 11): -3$$

$$(5 \ 12)(12 \ 7) = (5 \ 7): -16$$

$$(5 \ 12)(12 \ 11) = (5 \ 11): -15$$

$$(10 \ 12)(12 \ 7) = (10 \ 7): -9$$

$$(10 \ 12)(12 \ 11) = (10 \ 11): -8$$

$$(18 \ 12)(12 \ 11) = (18 \ 11): 0$$

$$(19\ 12)(12\ 7) = (19\ 7): -1$$

$$j = 20$$

$$(13\ 20)(20\ 1) = (13\ 1): -6$$

$$D_8^{-1}M^-(40)$$

	7	8	11	17	18	19	5	1	4	12	20	2	9	13	16	6	10	14	3	15	
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
1	0	50	43	-7	63	77	5	∞	88	41	56	79	30	20	56	29	37	8	63	39	1
2	-1	0	4	-8	42	5	<u>-4</u>	60	52	56	<u>-3</u>	-9	38	12	42	-2	46	25	8	-9	2
3	60	63	0	43	19	61	21	42	49	20	68	27	84	10	74	13	4	81	∞	63	3
4	85	79	29	0	58	84	83	20	∞	50	22	67	78	85	9	54	9	53	68	24	4
5	-13	45	-3	-20	0	80	<u>-16</u>	-14	-7	-1	<u>-15</u>	-21	58	72	-11	30	-11	-15	41	-21	5
6	0	72	8	-7	48	0	-3	-6	53	21	49	<u>-8</u>	50	26	60	∞	18	-2	9	-8	6
7	∞	70	77	87	91	54	0	-3	64	55	85	95	24	20	6	31	40	29	78	5	7
8	51	∞	24	23	77	66	51	0	48	47	69	48	58	55	58	29	91	64	70	33	8
9	60	13	84	71	62	78	57	86	0	18	60	<u>-2</u>	∞	86	37	73	57	15	43	-2	9
10	-6	82	24	-13	89	75	-5	-13	-6	0	<u>-8</u>	-14	62	76	-4	44	-4	53	-8	-14	10
11	27	32	∞	35	66	12	<u>0</u>	-8	-1	71	0	-5	13	11	47	18	4	66	62	-3	11
12	78	88	76	67	64	24	5	42	63	∞	6	0	38	77	61	38	43	41	51	44	12
13	<u>-6</u>	66	0	-11	52	-8	-11	-19	-12	52	-11	-12	0	∞	-2	65	-7	39	78	-14	13
14	12	33	4	-15	76	40	37	65	71	35	7	54	9	0	-6	77	-6	∞	70	72	14
15	44	63	80	61	21	32	47	66	24	34	73	35	72	41	0	36	62	26	47	∞	15
16	83	12	57	64	56	15	76	8	55	7	52	0	53	54	∞	0	43	15	72	43	16
17	82	23	18	∞	70	30	7	45	58	74	5	47	27	16	74	82	0	27	24	81	17
18	58	42	73	59	∞	42	<u>-1</u>	22	89	81	<u>0</u>	-6	74	33	17	71	71	0	75	-6	18
19	20	11	57	26	53	∞	<u>-1</u>	47	38	57	<u>0</u>	-6	58	81	51	42	82	55	0	-6	19
20	8	50	18	85	79	78	43	37	45	74	∞	0	29	59	49	46	12	41	67	0	20
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	

$$P_{40}$$

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
1																					1
2	2						1				12	2									2
3																					3
4																					4
5							12				12	5									5
6																					6
7																					7
8																					8
9																					9
10											12	10									10
11																					11
12																					12
13	20																			13	13
14																					14
15																					15
16																					16
17																					17
18							12				12	18									18
19							12				12	19									19
20																					20
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	

NONNEGPATHS(40)

(1) (13 1); (4) (13 4); (7) (2 7); (11) (2 11), (5 11), (10 11); (12) (6 12).

$$j = 1$$

$$(13 \ 1)(1 \ 4) = (13 \ 4): -13$$

$$(13 \ 1)(1 \ 7) = (13 \ 7): -1$$

$$j = 4$$

$$(13 \ 4)(4 \ 15) = (13 \ 15): -4$$

$$j = 7$$

$$(2 \ 7)(7 \ 8) = (2 \ 8): -7$$

$$j = 11$$

$$(2 \ 11)(11 \ 8) = (2 \ 8): -11$$

$$(5 \ 11)(11 \ 6) = (5 \ 6): -3$$

$$(5 \ 11)(11 \ 8) = (5 \ 8): -23$$

$$(5 \ 11)(11 \ 9) = (5 \ 9): -16$$

$$(10 \ 11)(11 \ 8) = (10 \ 8): -16$$

$$(10 \ 11)(11 \ 9) = (10 \ 9): -9$$

$$j = 12$$

$$(6 \ 12)(12 \ 11) = (6 \ 11): -2$$

NONNEGPATHS(60)

(6) (5 6); (8) (5 8), (10 8); (9) (5 9), (10 9); (11) (6 11).

$$j = 8$$

$$(5 \ 9)(9 \ 2) = (5 \ 2): -3$$

$$j = 11$$

$$(6 \ 11)(11 \ 8) = (6 \ 8): -10$$

$$(6 \ 11)(11 \ 9) = (6 \ 9): -3$$

$$D_8^{-1}M^-(60)$$

	7	8	11	17	18	19	5	1	4	12	20	2	9	13	16	6	10	14	3	15	
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
1	0	50	43	-7	63	77	5	∞	88	41	56	79	30	20	56	29	37	8	63	39	1
2	-1	0	4	-8	42	5	-4	60	52	56	-3	-9	38	12	42	-2	46	25	8	-9	2
3	60	63	0	43	19	61	21	42	49	20	68	27	84	10	74	13	4	81	∞	63	3
4	85	79	29	0	58	84	83	20	∞	50	22	67	78	85	9	54	9	53	68	24	4
5	-13	<u>-3</u>	-3	-20	0	-3	-16	-14	-16	-1	-15	-21	58	72	-11	30	-11	-15	41	-21	5
6	0	72	8	-7	48	0	-3	<u>-10</u>	<u>-3</u>	21	-2	-8	50	26	60	∞	18	-2	9	-8	6
7	∞	70	77	87	91	54	0	-3	64	55	85	95	24	20	6	31	40	29	78	5	7
8	51	∞	24	23	77	66	51	0	48	47	69	48	58	55	58	29	91	64	70	33	8
9	60	13	84	71	62	78	57	86	0	18	60	-2	∞	86	37	73	57	15	43	-2	9
10	-6	82	24	-13	89	75	-5	-16	-9	0	-8	-14	62	76	-4	44	-4	53	-8	-14	10
11	27	32	∞	35	66	12	<u>0</u>	-8	-1	71	0	-5	13	11	47	18	4	66	62	-3	11
12	78	88	76	67	64	24	5	42	63	∞	6	0	38	77	61	38	43	41	51	44	12
13	-6	66	0	-13	52	-8	-11	-19	-12	52	-11	-12	0	∞	-4	65	-7	39	78	-14	13
14	12	33	4	-15	76	40	37	65	71	35	7	54	9	0	-6	77	-6	∞	70	72	14
15	44	63	80	61	21	32	47	66	24	34	73	35	72	41	0	36	62	26	47	∞	15
16	83	12	57	64	56	15	76	8	55	7	52	0	53	54	∞	0	43	15	72	43	16
17	82	23	18	∞	70	30	7	45	58	74	5	47	27	16	74	82	0	27	24	81	17
18	58	42	73	59	∞	42	-1	22	89	81	<u>0</u>	-6	74	33	17	71	71	0	75	-6	18
19	20	11	57	26	53	∞	-1	47	38	57	<u>0</u>	-6	58	81	51	42	82	55	0	-6	19
20	8	50	18	85	79	78	43	37	45	74	∞	0	29	59	49	46	12	41	67	0	20
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	

$$P_{60}$$

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
1																					1
2																					2
3																					3
4																					4
5		9							5												5
6								11	11		6										6
7																					7
8																					8
9																					9
10																					10
11																					11
12																					12
13																					13
14																					14
15																					15
16																					16
17																					17
18																					18
19																					19
20																					20
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	

NONNEGPATH(80)

(2) (5 2); (11) (6 8), (6 9).

$j = 2$

$$(5 \ 2)(2 \ 16) = (5 \ 16): -5$$

Thus, the only path that can be extended is at $(5 \ 16)$. But, checking the possible extensions, none is non-negative. It follows that there exists no cycle, s , in $D_8^{-1}M^-$ such that $D_8 s$ is a tour. It follows that the n -cycle $D_7 = \sigma_{FWTSP_{OPT}}$. Furthermore, from the corollary to theorem 3.6, $D_7 = \sigma_{TSP_{OPT}}$.

As addenda, we include a full list of c-trees.

REFERENCES

- [1] T. H. CORMEN, C. E. LEISERSON, R. L. RIVEST, *An Introduction to Algorithms*, The MIT Press (2000), 527-532.
- [2] W. FELLER, *An Introduction to Probability Theory and its Applications (Vol. I)*, John Wiley and Sons (1950), New York, London, 242-243.
- [3] R. W. FLOYD, *Algorithm 97: Shortest Path*, Comm. ACM, 5, no. 6 (1962), 345.
- [4] C. PAPADIMITROU, K. STIGLITZ, *Combinatorial Optimization: Algorithms and Complexity*, Prentice-Hall (1982); second edition, Dover (1998).
- [5] S. Warshall, A theorem on Boolean matrices, J. ACM., 9, no. 1, (1962)