

Neuronal nicotinic receptors: from structure to function

Sherry Leonard, Daniel Bertrand

Molecular identification of the genes and resulting protein sequences for a large number of nicotinic acetylcholine receptor (nAChR) subunits has stimulated numerous studies highlighting their role in several human behaviors including neurological disorders and nicotine addiction. This receptor gene family is likely to be involved in release of multiple neurotransmitters in both brain and periphery that mediate sensitivity and tolerance to nicotine. Recent findings also suggest that alterations in these receptors may lead to neurological diseases, some associated with increased incidence of smoking. This review addresses current knowledge of nicotinic receptor structure, regulation of expression, and function, in both normal and psychiatric subjects.

Introduction

Expression and function of the nicotinic acetylcholine receptor (nAChR) gene family are likely to be important factors in nicotine addiction. The development of tolerance and eventual addiction to nicotine, as well as the responses to the neuroadaptations that lead to withdrawal symptoms and relapse, are complex and involve multiple neurotransmitter systems. However, the principal effects of nicotine downstream from the cigarette begin with its action at the nicotinic receptor. Interventions, presently available to smokers, do not result in high quit rates at 1 year post-cessation. The addition of antidepressants has made a minor improvement in treatment, but permanent smoking cessation is still rarely achieved, particularly in the mentally ill. Understanding the structure, expression and function of this large gene family and its relationship to other neurotransmitter systems is essential for the formulation of nicotine addiction hypotheses and subsequent treat-

ments that will result in true smoking cessation in both normal subjects and those with psychiatric disorders.

The presence of nicotinic receptors on muscle cells and neurons that display functional differences in response to acetylcholine has been recognized for many years (Ascher, Large, & Rang, 1979; Popot & Changeux, 1984; Rang & Ritter, 1970). However, the gene family is large and the possible subunit combinations numerous. Furthermore, in recent years it has become clear that neuronal nAChRs are widely expressed throughout the body, where they are likely to have diverse but important roles.

To date, 11 genes coding for neuronal nAChR subunits have been identified from the mammalian genome ($\alpha 2$ – $\alpha 7$, $\alpha 9$, $\alpha 10$, and $\beta 2$ – $\beta 4$), reviewed in Lukas and Bencherif (1992), McGehee and Role (1995), Elliott *et al.* (1996), Gotti, Fornasari, and Clementi (1997), and also see GenBank Accession Number AF196344. Phylogenetically linked to the broader family of four transmembrane ligand-gated channels, which include the γ -aminobutyric acid (GABA_A and GABA_C), the glycine and serotonergic receptors (5-HT₃), nicotinic receptors are thought to be among the first of the ligand-gated ion channels to evolve, long before evolution of vertebrates and their complex nervous systems (Le Novère & Changeux, 1995). The nicotinic receptor gene family is even larger in invertebrates (Ballivet, Alliod, Bertrand, & Bertrand, 1996; Jonas, Phannavong, Schuster, Schroder, & Gundelfinger,

Sherry Leonard, Departments of Psychiatry and Pharmacology, University of Colorado Health Sciences Center, and the Veterans Affairs Medical Research Service, Denver, Colorado; Daniel Bertrand, Department of Physiology, C.M.U., Geneva, Switzerland.

Correspondence to: Sherry Leonard, PhD, Department of Psychiatry, Box C-268–71, University of Colorado Health Sciences Center, 4200 E. 9th Avenue, Denver, CO 80262, USA. Tel: 303–315–8412; Fax: 303–315–5347; E-mail: Sherry.Leonard@UCHSC.edu

1994; Schulz *et al.*, 1998; Squire *et al.*, 1995). Since at least 16 genes for nAChR subunits are believed to be expressed in *Caenorhabditis elegans* (Brownlee & Fairweather, 1999), it is possible that additional subunit genes remain to be discovered in vertebrates. In that regard, the sequence for a novel rat neuronal nicotinic receptor cDNA ($\alpha 10$), was recently released by Jim Boulter, UCLA (GenBank Accession Number: AF196344). A human ortholog has also been identified (F. Sgard and P. Fuchs, personal communication).

Neuronal receptors are thought, as their muscle counterparts, to result from the assembly of five subunits around an axis of pseudosymmetry (Bertrand & Changeux, 1992; Miyazawa, Fujiyoshi, Stowell, & Unwin, 1999; Unwin, 1996). Each of the neuronal subunits is likely to span the membrane four times with the second transmembrane domain forming the wall of the ionic pore (Bertrand & Changeux, 1992). While overall structures of the proteins forming the neuronal nicotinic receptors appear quite similar, their genes and exon–intron relationships differ markedly (Lindstrom, 1996). The acetylcholine (ACh) binding sites and membrane spanning regions (MI–IV) are highly conserved between subunit types, but the large cytoplasmic loop between MIII and MIV is unique to each subunit and conserved across species (Elliott *et al.*, 1996; McGehee & Role, 1995). A comparison of exon/intron borders for a subset of the nicotinic receptor subunit gene family is shown in Figure 1A. The exon/intron junction pattern is more complex for the $\alpha 7$ subunit, which has 10 exons, than for the other subunits, which have five or six. Further, two of these splice junctions, unique to the $\alpha 7$ subunit, are found near the border of MI and within the coding region for MII, the latter thought to face the ion channel (Corringer, Le Novère, & Changeux, 2000; Galzi, Revah, Bessis, & Changeux, 1991). The location of the exon/intron splice sites in the $\alpha 7$ gene, relative to the amino acid sequence of MI and MII are shown schematically in the assembled pentamer in Figure 1B.

Heterologous expression studies have revealed that functional neuronal nAChRs can be divided in two groups: (a) the homomeric receptors and (b) the heteromeric receptors (Weiland, Bertrand, & Leonard, 2000). Expression of $\alpha 7$, $\alpha 8$ or $\alpha 9$ subunits alone has been shown to result in functional receptors, while all the other subunits require at least a pair wise expression of an α and β combination (Elgoyhen, Johnson, Boulter, Vetter, & Heinemann, 1994; Gerzanich, Anand, & Lindstrom, 1994; Kuryatov, Olale, Cooper, Choi, & Lindstrom, 2000; Luetje & Patrick, 1991). The newly identified $\alpha 10$ subunit is more homologous to $\alpha 7$ and $\alpha 9$ than to the α subunits that bind nicotine with high affinity ($\alpha 4$, $\alpha 3$). Attempts to express $\alpha 10$ alone yielded no functional receptors but co-expression with its closest homolog $\alpha 9$ caused a marked difference in $\alpha 9$ receptor properties that more closely resembled those of the native receptor in outer hair cells (F. Sgard, personal communication). This suggests that $\alpha 9$ and $\alpha 10$ probably assemble in the same receptor complex.

Although the $\alpha 7$ -receptor subunit may be expressed in the brain as an homomer (Drisdel & Green, 2000), whether it always occurs in this conformation *in vivo* is unclear. Functionally distinct, $\alpha 7$ containing receptors exist in chick sympathetic neurons that may contain other subunits (Girod *et al.*, 1999; Yu & Role, 1998b). It was also recently reported that while neurons from chick cardiac ganglia express a receptor containing the $\alpha 7$ subunit they display a pharmacological profile distinct from that of homomeric receptors reconstituted in *Xenopus* oocytes or in cell lines (Cuevas & Berg, 1998). It is possible that co-assembly of the $\alpha 7$ subunit with other nicotinic receptor subunits is region or tissue specific and may depend on the relative ratio or type of subunits expressed (Nelson & Lindstrom, 1999; Yu & Role, 1998a).

It is, perhaps, not surprising that the $\alpha 7$ subunit is principally expressed as an homomer, since it was probably the first subunit in the gene family to evolve, long before development of the brain. Major rearrangements in the neuronal nicotinic receptor genes have occurred during evolution by gene duplication and mutation, resulting in the variable exon content (Le Novère & Changeux, 1995) and different regulatory mechanisms. The earliest nicotinic receptor subunit gene is thought to be the $\alpha 7$ gene, which maps to human chromosome 15q13–q14. A partial duplication of the $\alpha 7$ gene (*dupa7*) has been cloned and maps less than 1Mb proximal to the full-length $\alpha 7$ gene. *Dupa7* is expressed as mRNA with novel exons and is alternatively spliced, as is the mRNA for the full-length $\alpha 7$ gene (Gault *et al.*, 1998). Three other nicotinic receptor subunit genes map distally at chromosome 15q24 ($\alpha 3$, $\alpha 5$ and $\beta 4$), and form a gene cluster (Boulter *et al.*, 1990; Couturier *et al.*, 1990a). Three subunits of the muscle nicotinic receptor map near each other on chromosome 2 (Lobos, 1993). Table 1 summarizes our current knowledge of the human chromosomal localization of the neuronal nicotinic receptor subunit genes.

Localization of gene expression

Compared to other ligand-gated ion channels, gene expression of the neuronal nicotinic receptors is generally low in the brain. Expression for each subunit varies during development and across brain regions. The most abundant subunits expressed in human brain are the $\alpha 4$, $\alpha 7$, and $\beta 2$ subunits. The highest levels of $\beta 2$ expression were found in caudate, putamen, and hippocampus, while $\alpha 3$ expression was highest in thalamus (Gotti *et al.*, 1997; Rubboli *et al.*, 1994). The $\alpha 4$ subunit was most highly expressed in the cortex (Agulhon *et al.*, 1998; Wevers *et al.*, 1994), while $\alpha 7$ expression was highest in human lateral and medial geniculate and the reticular thalamic nucleus (RTN; Agulhon, Abitbol, Bertrand, & Malafosse, 1999; Breese *et al.*, 1997a; Wevers *et al.*, 1995). Expression of $\alpha 5$ is generally low in the human central nervous system (CNS); except for cerebellum and thalamus, where




Figure 1. Schematic representation of splice junctions for neuronal nicotinic acetylcholine receptor subunit genes and an assembled $\alpha 7$ homomeric receptor. (A) Exon/intron junctions for a subset of the nicotinic acetylcholine receptor subunit gene family. SP, signal peptide; M I to M IV, membrane spanning regions I to IV; A, B, and C, three domains of the ACh binding site. Arrows indicate the intron positions (Elgoyhen, Johnson, Boulter, Vetter, & Heinemann, 1994; Steinlein, Weiland, Stoodt, & Propping, 1996; Rempel, Heyers, Engels, Slegers, & Steinlein, 1998; Gault *et al.*, 1998; F. Sgard, personal communication). (B) Representational structure of an assembled $\alpha 7$ homomeric nicotinic receptor. The ACh binding site is indicated and scissors are positioned at the sites of introns 7 and 8 in the coding region of the gene. Intron 7 lies near the coding region for M I, intron 8 within the coding region for M II, and intron 9 in the coding region for M III.

Table 1. Chromosomal localization for the human nicotinic acetylcholine receptor gene family. Published map locations are shown

Receptor subunit	Subunit type	Map location	Reference
$\alpha 1$	Muscle	2q24-q32	Beeson <i>et al.</i> , 1990; Lobos <i>et al.</i> , 1993
$\beta 1$	Muscle	17p12-p11	Beeson <i>et al.</i> , 1990; Lobos <i>et al.</i> , 1993
γ	Muscle	2q33-q34	Beeson <i>et al.</i> , 1993; Lobos <i>et al.</i> , 1993
δ	Muscle	2q33-q34	Beeson <i>et al.</i> , 1990; Lobos <i>et al.</i> , 1993
ϵ	Muscle	17q13-p12	Beeson <i>et al.</i> , 1993; Lobos <i>et al.</i> , 1993
$\alpha 2$	Neuronal	8p21	Wood <i>et al.</i> , 1995
$\alpha 3$	Neuronal	15q24	Boulter <i>et al.</i> , 1990; Raimondi <i>et al.</i> , 1992
$\alpha 4$	Neuronal	20q13.2-q13.3	Steinlein <i>et al.</i> , 1994
$\alpha 5$	Neuronal	15q24	Boulter <i>et al.</i> , 1990; Raimondi <i>et al.</i> , 1992
$\alpha 7$	Neuronal	15q13-q14	Chini <i>et al.</i> , 1994; Orr-Utregger <i>et al.</i> , 1995
$\beta 2$	Neuronal	1q21.3	Lueders <i>et al.</i> , 1999
$\beta 3$	Neuronal	8p11.2	Koyama <i>et al.</i> , 1994
$\beta 4$	Neuronal	15q24	Boulter <i>et al.</i> , 1990; Raimondi <i>et al.</i> , 1992

moderate levels were observed (Flora *et al.*, 2000a). A disparity in regional expression between the human and rat genes has been noted for $\alpha 7$ in the RTN (Breese *et al.*, 1997a), and for $\alpha 5$ in cerebellum and thalamus (Flora *et al.*, 2000a). Although the pattern of subunit expression is highly conserved across species, disparities between rodents and mammals for expression of the $\alpha 3$ and $\alpha 2$ subunits have been reported (Cimino, Marini, Fornasari, Cattabeni, & Clementi, 1992; Han *et al.*, 2000). These results suggest that species divergence in gene expression has occurred that might affect neuronal circuitry. This possibility merits consideration when pathways regulating behaviors are being studied in multiple species.

It has been generally accepted that the neuronal nicotinic receptor gene family was mainly expressed in the brain and ganglionic synapses of the peripheral nervous system, although a broader localization has recently been suggested. Subunits of principal importance in the chick ciliary ganglia and adrenal chromaffin cells are $\alpha 3$, $\beta 4$, and $\alpha 7$ (Lukas, Norman, & Lucero, 1993; McGehee & Role, 1995; Vernallis, Conroy, & Berg, 1993; Vijayaraghavan, Pugh, Zhang, Rathouz, & Berg, 1992). Expression of the $\alpha 7$ subunit has also been found in developing muscle (Romano, Pugh, McIntosh, & Berg, 1997). Multiple nicotinic receptor subunits, including $\alpha 3$, $\alpha 5$, $\alpha 7$, $\beta 2$, and $\beta 4$ are expressed in keratinocytes, bronchial epithelial cells, and arterial endothelium (Grando *et al.*, 1995; Horton, Nguyen, & Grando, 1997; Macklin, Maus, Pereira, Albuquerque, & Conti-Fine, 1998; Maus *et al.*, 1998; Zia, Ndoeye, Nguyen, & Grando, 1997). A recent survey of $\alpha 5$ mRNA expression in human peripheral tissues indicates this receptor subunit is also expressed in the gut and in the thymus (Flora *et al.*, 2000a,b). In cultured bronchial epithelial cells, nicotine regulates both the integrity of cell shape and motility (Conti-Fine, Navaneetham, Lei, & Maus, 2000).

Nicotinic receptors are found in both lymphocytes and polymorphonuclear (PMN) cells of human peripheral blood where their function is not yet known (Benham-

mou *et al.*, 2000; Hiemke *et al.*, 1996; Lebagry *et al.*, 1996). In some peripheral tissues, cells have been found to express both a cholinergic phenotype and to have nAChRs on their surface; function is thought to involve autocrine or paracrine mechanisms (Nguyen *et al.*, 2000; Wessler, Kirkpatrick, & Racke, 1999). Thus, receptors responding to nicotine have a wide expression pattern and may have non-synaptic roles in addition to those in the CNS.

Receptor assembly

Nicotinic receptor subunits are known to undergo post-translational modification by glycosylation and phosphorylation (Chen, Dang, & Patrick, 1998). Glycosylation has been shown to be necessary for binding of some ligands (Luetje, Maddox, & Harvey, 1998) and perhaps for correct folding of the peptide (Blount & Merlie, 1990; Rickert & Imperiali, 1995), but may not be required for receptor assembly (Chen *et al.*, 1998). However, subunits that do not acquire ligand-binding capacity may be rapidly degraded (Blount & Merlie, 1990), affecting efficient receptor assembly (Green & Millar, 1995). Indeed, nicotinic receptor assembly appears to be exquisitely sensitive to correct folding; only 30% of synthesized subunits in the muscle become incorporated into competent surface receptors (Blount & Merlie, 1990).

Although several stable cell lines expressing human neuronal nicotinic receptors have been developed (Buisson, Gopalakrishnan, & Bertrand, 1999), obtaining competent receptor expression of $\alpha 7$ receptors in stably transfected cell lines is considered to be more difficult than for other subunits. Additionally, receptor expression in the transfected cultures is not generally uniform. Studies using chimaeras of the $\alpha 7$ subunit, usually with the 5-HT₃ receptor that is also a ligand-gated ion channel, suggest that the membrane-spanning regions may contribute to the low assembly efficiency for this receptor (Cooper, Harkness, Baker, & Millar,




Figure 2. Schematic representation of the $\beta 4$, $\alpha 3$, $\alpha 5$ gene cluster. (A) Map of the $\beta 4$, $\alpha 3$, $\alpha 5$ gene cluster. Arrows indicate the transcription polarities. (B) Enlargement of $\beta 4$ gene map showing the principal known promoters and enhancer elements. (C) Schematic drawing of the $\beta 4$ transcription regulatory elements.

1999; Corringer *et al.*, 1995; Eisele *et al.*, 1993). Indeed, specific amino acids in the $\alpha 7$ subunit have been found to control receptor distribution between the cell surface and intracellular pools (Dineley & Patrick, 2000). Amino acids near and within membrane-spanning region I (MI) were found to affect surface receptor expression. Since overexpression, or a significant increase in function of the $\alpha 7$ receptor has been shown to result in increased neuronal cell death (Berger, Gage, & Vijayaraghavan, 1998; Orr-Urtreger *et al.*, 2000), the tight regulation of functional surface expression demonstrated for the $\alpha 7$ receptor may be important in development.

Despite the stringency apparently required for competent surface $\alpha 7$ -receptor expression and the fact that the stability of $\alpha 7$ -receptor expression in cultured cell lines derived from tumors has also been questioned (Kassner & Berg, 1997), a rat $\alpha 7$ cDNA has been successfully expressed in several cell lines including rat pituitary (Quik, Choremis, Komourian, Lukas, & Puchacz, 1996), neuronally derived human cell line SH-SY5Y (Puchacz, Buisson, Bertrand, & Lukas, 1994), and a human $\alpha 7$ clone in embryonic kidney cell line HEK 293 (Gopalakrishnan *et al.*, 1995).

Regulation of gene expression

Levels of gene expression and tissue specificity are thought to be principally determined by regulatory elements in the promoter and 5'-untranslated regions (UTRs) of genes. The diversity of transcription factor binding sites, enhancer and repressor sequences in the nicotinic receptor subunit genes, thus far characterized, suggest exquisite and generally individualized regulation by these promoter elements (Fornasari, Battaglioli, Terzano, & Clementi, 1999). The $\beta 4$ and $\alpha 3$ genes lie in a gene cluster on human chromosome 15q24, which also includes $\alpha 5$ (Table 1; Figure 2A). Conservation of this cluster suggests there are likely to be shared regulatory mechanisms (Boulter *et al.*, 1990). All three of these genes are GC-rich and have promoter elements that bind the transcription factor Sp1 (Bigger, Casanova, & Gardner, 1996; Fornasari, Battaglioli, Flora, Terzano, & Clementi, 1997), as do many of the nicotinic receptor subunit gene promoters (Fornasari *et al.*, 1999). Elements that coordinately regulate the expression of the $\alpha 3$ and $\beta 4$ genes have been identified in the region between the two genes. A neuron-selective enhancer, the $\beta 43'$, functions in both orientations and coordinately activates

both the $\alpha 3$ and $\beta 4$ genes, which are transcribed in the same direction (Fyodorov, Nelson, & Deneris, 1998; McDonough & Deneris, 1997). The $\beta 43'$ enhancer also appears to be activated by ETS factors such as Pet-1 (Figure 2B; Deneris *et al.*, 2000), as was found for muscle subunits γ and ϵ (Schaeffer, Duclert, Huchet-Dymanus, & Changeux, 1998).

Transcription of the $\alpha 5$ gene is known to occur in the opposite direction to that of $\alpha 3$ and $\beta 4$ and thus $\alpha 5$ may not share promoter or enhancer elements with the other two genes in the cluster. The $\alpha 5$ gene is expressed in both neuronal and non-neuronal tissues and the 5'-UTR of the $\alpha 5$ promoter can act as either an activator or an inhibitor of gene expression in neuronal or non-neuronal cells, respectively (Flora *et al.*, 2000b). Expression of the $\alpha 5$ subunit is observed as early as E11 in the chick. Homeodomain transcription factor Phox 2B, necessary for neurogenesis in the autonomic nervous system (Pattyn, Morin, Cremer, Goridis, & Brunet, 1999), activates $\alpha 3$, but not $\alpha 5$.

A core promoter for the $\beta 4$ gene of 226 bp has been isolated, containing both a CT-box binding Pura and hnRNPK (Du, Tomkinson, & Gardner, 1997), and a CA-box binding transcription factors Sp1 and Sp3 (Figure 1C; Bigger, Melnikova, & Gardner, 1997; Du, Melnikova, & Gardner, 1998). The binding site for another factor, Sox 10, overlaps the CT and CA boxes (Melnikova, Yang, & Gardner, 2000). In addition to Sp1 sites, the $\alpha 3$ promoter is activated by the Pou domain transcription factor SCIP, apparently through a protein-protein interaction (Deneris *et al.*, 2000).

A core promoter of 143 bp has been isolated for the $\beta 3$ gene (Matter *et al.*, 1998), containing an E box that binds basic helix-loop-helix (bHLH) proteins. Specifically, the *atonal* homolog cATH5, whose expression is principally confined to developing retina, activates the $\beta 3$ core promoter more than 10-fold. The time courses of expression during development for both cATH5 and $\beta 3$ are similar.

The sequences for the bovine (Carrasco-Serrano, Campos-Caro, Viniegra, Ballesta, & Criado, 1998) and human (Gault *et al.*, 1998) $\alpha 7$ gene promoters have been reported. The two core promoter sequences are similar; both are GC-rich and contain several Egr-1 transcription factor consensus sites, as well as a CREB site, suggesting that the promoters may be steroid regulatable. This is consistent with results in the mouse, showing that the $\alpha 7$ gene is regulated by neurosteroids (Bullock *et al.*, 1997; Pauly, Grun, & Collins, 1990). The Egr-1 component of transcription factor binding to the bovine $\alpha 7$ promoter is increased in bovine adrenomedullary cells on treatment with phorbol esters, indicating that the $\alpha 7$ gene may be a specific target for *egr-1* gene expression (Carrasco-Serrano, Viniegra, Ballesta, & Criado, 2000).

Although functional investigation of regulatory elements for most of the nicotinic receptor subunit genes is far from complete, there are similarities and differences between the elements controlling the expression of the

different subunits. Differences are expected to play a fundamental role during development and tissue specific activation, likely controlling the gene specific roles of the subunits. For example, since it is known that the $\alpha 3\beta 4$ combination is the most abundant in ganglionic receptors, coordinated regulation for the expression of these two genes may be expected in the autonomic system.

Function

One of the interesting features of the neuronal nAChRs is their localization in both the brain and periphery, and how this relates to function. Although individual nAChR subtypes may have a more restricted localization, one or more subtypes are found in most tissues. Observations made on rat brain synaptosomes have shown that application of nicotinic agonists can provoke the release of dopamine (Grady, Marks, Wonnacott, & Collins, 1992; Grady, Marks, & Collins, 1994; Misu *et al.*, 1993; Wonnacott, 1997). However, strong evidence suggests that nicotinic receptors are expressed both pre- and post-synaptically, and in complex patterns that include both axo-axonic and somato-dendritic synapses (Amador & Dani, 1995; Bordey, Feltz, & Trouslard, 1996; Dolezal, Lee, Schobert, & Hertting, 1996; Guo, Tredway, & Chiappinelli, 1998; Lena & Changeux, 1997; MacDermott, Role, & Siegelbaum, 1999; Marchi & Raiteri, 1996; McGehee & Role, 1996; Ullian & Sargent, 1995). Additionally, the nicotinic receptor subunit gene family appears to be growing with the discovery of the $\alpha 10$ subunit (Boulter *et al.*, GenBank Accession Number AF196344; F. Sgard personal communication), and it is not yet known whether all of the subunits have been cloned. With the completion of the genome sequencing efforts, this question should be resolved in the near future.

Examination of the physiological properties of heterologously expressed nicotinic receptors, composed of the known subunits, reveals large differences between receptor subtypes. At one extreme are the receptors that desensitize slowly, composed of $\alpha 4\beta 2$ subunits (Bertrand, Ballivet, & Rungger, 1990; Buisson, Gopalakrishnan, Arneric, Sullivan, & Bertrand, 1996) while at the other extreme is the fast desensitizing $\alpha 7$ receptor (Couturier *et al.*, 1990b; Gopalakrishnan *et al.*, 1995; Séguéla, Wadiche, Dineley-Miller, Dani, & Patrick, 1993; Vijayaraghavan *et al.*, 1992; Zhang, Vijayaraghavan, & Berg, 1994). Receptors composed of $\alpha 3\beta 4$ or $\alpha 3\beta 4\alpha 5$ display a slower time course of desensitization than $\alpha 7$ receptors (Couturier *et al.*, 1990a; Role, 1992).

Such differences have also been observed in brain receptors measured either on cultured cells or in slices (Alkondon & Albuquerque, 1993, 1995; Alkondon *et al.*, 1994; Alkondon, Rocha, Maelicke, & Albuquerque, 1996; Frazier, Buhler, Weiner, & Dunwiddie, 1998a; Frazier *et al.*, 1998b; Lena *et al.*, 1999; Zaninetti,

Tribollet, Bertrand, & Raggenbass, 1999; Zoli, Lena, Picciotto, & Changeux, 1998). Differences in pharmacological properties have been further documented *in vivo* (Decker, Brioni, Bannon, & Arneric, 1995). Speculations concerning the role of the nicotinic receptors, therefore, depend upon the subtype considered, the tissue in which they are expressed, and the subcellular localization.

High sensitivity to nicotine and sustained responses, which are characteristic of the major brain $\alpha 4\beta 2$ receptor, result in prolonged tonic responses while low sensitivity and highly desensitizing receptors typical of $\alpha 7$ are thought to provoke a fast, phasic depolarization (Boulter *et al.*, 1987; Monteggia *et al.*, 1995), and for $\alpha 7$ (Couturier *et al.*, 1990b; Peng, Katz, Gerzanich, Anand, & Lindstrom, 1994; Séguéla *et al.*, 1993). Responses of either $\alpha 4\beta 2$ or $\alpha 7$ receptors are accompanied by a significant increase in the intracellular calcium concentration (Bertrand, Galzi, Devillers, Bertrand, & Changeux, 1993; Haghighi & Cooper, 2000; Ragozzino, Barabino, Fucile, & Eusebi, 1998; Rogers & Dani, 1995; Sands, Costa, & Patrick, 1993; Tsuneki, Klink, Lena, Korn, & Changeux, 2000; Vernino, Amador, Luetje, Patrick, & Dani, 1992). Thus, for receptors expressed pre-synaptically it can be expected that they play a role in modulating the response of the synaptic bouton, and either directly or indirectly increasing the intracellular calcium concentration. Elevation of the intracellular calcium concentration increases the probability of neurotransmitter release. Indeed, as demonstrated by several groups, neuronal nAChRs play a major role in controlling release of many different neurotransmitters (Guo *et al.*, 1998; Lena & Changeux, 1997; McGehee & Role, 1995; Pidoplichko, DeBiasi, Williams, & Dani, 1997; Wonnacott, 1997). In contrast, fast synaptic transmission caused by neuronal nicotinic receptors has been demonstrated in only a few cases in the CNS, but is expected to play an important role that awaits further characterization (Alkondon, Pereira, Eisenberg, & Albuquerque, 1999; Frazier *et al.*, 1998a, b; Hefft, Hulo, Bertrand, & Muller, 1999; Roerig, Nelson, & Katz, 1997). Additional modulatory roles of nicotinic receptors can also be anticipated in view of their properties and subcellular localization. For example, it is widely documented that nAChRs are expressed on the cell soma and ACh-evoked currents have been classified (Albuquerque *et al.*, 1995; Lena *et al.*, 1999; Yakel & Jackson, 1988; Zaninetti *et al.*, 1999; Zoli *et al.*, 1998). Stimulation of these receptors is expected to cause effects both through cell depolarization and/or increase of the intracellular calcium concentration. Evidence of calcium modulation has already been provided for NMDA receptors (Fisher & Dani, 2000). The effect of nAChR activation in stratum radiatum interneurons was recently found to induce either inhibition or disinhibition of pyramidal neurons that may be dependent on activation of specific receptor subtypes (Ji & Dani, 2000).

Recent reports also suggest that single cells can express more than one receptor subtype. In approximately 30% of the ventral tegmental (VTA) dopaminergic neurons, both fast and slow desensitizing currents have been found, indicating that these neurons may express both $\alpha 7$ and $\alpha 4\beta 2$ receptors (Pidoplichko *et al.*, 1997). In contrast, only 4% of neurons in the substantia nigra display these two types of responses. The activation and desensitization kinetics in the VTA varied significantly with low concentrations of nicotine, altering the firing of the neurons on both a shorter and a longer time scale (Pidoplichko *et al.*, 1997). In this same brain area it was also shown that activation of nicotinic receptors on presynaptic terminals enhanced glutamatergic input to dopaminergic neurons (Mansvelder & McGehee, 2000). From the sensitivity to methyllycconitine (MLA) these investigators concluded that both short- and long-term effects of nicotine required activation of an $\alpha 7$ -containing receptor. Additional evidence for the contribution of $\alpha 7$ receptors to the intracellular calcium concentration was also provided in mouse brain slices (Tsuneki *et al.*, 2000).

In rat superior cervical ganglia two types of $\alpha 7$ containing, MLA sensitive receptors have been characterized (Cuevas, Roth, & Berg, 2000). One class has fast desensitization properties similar to homomeric $\alpha 7$ receptors. However, the second class, also blockable by MLA, desensitized slowly and differed in the reversibility of α -bungarotoxin blockade.

In the striatum, pharmacological evidence for two types of nicotinic receptors on dopaminergic terminals has also been found. Using α -conotoxin-MII, which has been shown to display a high selectivity for $\alpha 3\beta 2$ and the anatoxin-a/epibatidine hybrid UB-165, $\alpha 3\beta 2$ and $\alpha 4\beta 2$ receptors were found to play a role in dopamine release (Kaiser, Soliakov, Harvey, Luetje, & Wonnacott, 1998; Sharples *et al.*, 2000). Dopamine release was partially blocked by $\alpha 7$ antagonists and glutamate antagonists, suggesting that the $\alpha 7$ effect may be modulated through glutamate release onto dopaminergic terminals (Kaiser & Wonnacott, 2000). However, *in vivo* glutamate release may be dependent on nicotine dose; at low nicotine concentrations, tonic activation of NMDA receptors in the VTA may be necessary for dopamine release (Fu, Matta, Gao, Brower, & Sharp, 2000). The major subunit expressed in the mesolimbic system appears to be the $\alpha 6$ receptor (Goldner, Dineley, & Patrick, 1997; Le Novère *et al.*, 1999; Le Novère, Zoli, & Changeux, 1996). Since the physiological and pharmacological characterization of $\alpha 6$ -containing receptors remains to be determined, evidence regarding the specificity of the α -conotoxin-MII effects on $\alpha 6$ -containing vs. $\alpha 3\beta 2$ nAChRs has not yet been firmly established.

Evidence for more than one receptor subtype on GABAergic neurons from cultured rat hippocampus and in hippocampal slices from both human and rat has been reported (Albuquerque, Pereira, Alkondon, Schrattenholz, & Maelicke, 1997; Alkondon, Pereira, & Albuquerque, 1998; Hilmas, Alkondon, Eisenberg, &

Albuquerque, 1999). Acetylcholine induced both a rapidly decaying current and a slowly decaying whole-cell current; the latter was sensitive to dihydro- β -erythroidine (DH β E). The fast current was likely to be α 7 mediated, as it was blocked by MLA, and the slow was likely an α 4 β 2 receptor. Approximately 45% of interneurons in hippocampus may exhibit ACh currents, mediating effects on GABA release by both receptor subtypes.

A further complexity in evaluating the effects of neuronal nAChRs is their possible direct interaction with other ligand-gated channels. For example, occlusion of P2X responses by neuronal nAChR activation has been documented by several laboratories (Johnson, Shum, Thornton, & Bornstein, 1999; Searl, Redman, & Siliński, 1998; Zhou & Galligan, 1998). Co-activation of α 3 β 4 nAChR and P2X receptors results in cross-inhibition of both channel types (Khakh, Zhou, Sydes, Galligan, & Lester, 2000). These authors have demonstrated that inhibition was observed only at high channel density and was found to be state dependent.

Localization of α 7 receptors on dendritic spines associated with cytoskeletal elements has been found in chick ciliary ganglia, where apparent function may involve receptor stability and prevention of run-down. The clustering on the spines may have the effect of protecting the cell from rapid Ca^{2+} influx. Intracellular injection of the calcium-chelating agent BAPTA caused a significant reduction of the response. Run-down appears to be under the influence of calmodulin, through a calmodulin kinase II pathway. Blockade of the phosphatase calcineurin increased run-down (Chang & Berg, 1999; Conroy & Berg, 1999; Liu & Berg, 1999). The actin cytoskeleton has been shown to be important for retaining these receptor-rich spines (Shoop, Yamada, & Berg, 2000).

Pharmacology

Although acetylcholine is the principal endogenous ligand for the neuronal nicotinic receptors, other pharmacological agents may activate or inhibit the receptors. It is important to recall that while α 4 β 2 receptors display a sensitivity to ACh and nicotine in the micromolar range (Buisson *et al.*, 1996), α 7 receptor half activation is observed only at about 150 μ M for ACh and 40 μ M for nicotine (Gopalakrishnan *et al.*, 1995). In slices of both human cerebral cortex and rat hippocampus, choline (10 mM) induced an inward current that decayed to baseline prior to the termination of the agonist pulse and was sensitive to 50 nM MLA (Alkondon, Pereira, Cortes, Maelicke, & Albuquerque, 1997; Alkondon *et al.*, 1999). Lower concentrations (1 mM) have been found to potentiate the effects of ACh at α 4 β 4 receptors expressed in oocytes (Zwart & Vijverberg, 2000). Choline concentration in the CSF ranges between 4 and 12 μ M (Klein, Gonzales, Koppen & Loffelholz, 1993), suggesting that choline may not affect the response of

endogenous nAChR receptors. However, choline concentration at the synapse and in the vicinity of perisynaptic receptors is not known and may reach millimolar concentrations immediately following acetylcholine hydrolysis. Several compounds have been found that appear to modulate nicotinic receptors. Galanthamine is a reversible, competitive cholinesterase inhibitor that allosterically increases activity of nicotinic receptors and has shown some promise in treatment of Alzheimer's disease (Maelicke & Albuquerque, 2000; Raskind, Peskind, Wessel, & Yuan, 2000; Tariot *et al.*, 2000). Nefiracetam, and its parent alkaloids piracetam and aniracetam, have been recently shown to facilitate hippocampal neurotransmission, as well (Oyaizu & Narahashi, 1999; Nishizaki *et al.*, 2000; Nomura & Nishizaki, 2000). The facilitatory action was blocked by antagonists of both high- and low-affinity nicotinic receptors, suggesting that these compounds might be effective at multiple receptor subtypes.

Substance P and other tachykinins also interact with multiple nicotinic receptor subunits (Cuevas & Adams, 2000; Lukas & Eisenhour, 1996; Valenta, Downing, & Role, 1993). Substance P has been shown to inhibit epibatidine stimulated release of catecholamine from chromaffin cells with an IC_{50} of 5 μ M (Krause, Michael, Lubke, Livett, & Oehme, 1997). Receptor inhibition was found to be non-competitive and to involve the β subunits (Stafford, Oswald, Figl, Cohen, & Weiland, 1998; Stafford, Oswald, & Weiland, 1994). The α 7-receptor inhibition by substance P was similar to that of the high-affinity nicotinic receptors. Although substance P inhibition was always observed in the micromolar range, this interaction may play an important physiological role as suggested by the recent findings in rat cardiac ganglia neurons (Cuevas & Adams, 2000).

Several studies suggest that endogenous neurosteroids may be allosteric inhibitors of nicotinic receptors (Bertrand, Valera, Bertrand, Ballivet, & Rungger, 1991; Ke & Lukas, 1996; Valera, Ballivet, & Bertrand, 1992). Progesterone was found to be an inhibitor of α 4 β 2 receptors, while A-ring metabolites of progesterone appear to also inhibit α 3-containing receptors that regulate dopamine release (Bullock *et al.*, 1997). A recent comparison of several other neurosteroids indicated that compounds with the β -orientation of groups at the C-17 position are more potent than the α -oriented diastereomers (Paradiso, Sabey, Evers, Zorumski, Covey, & Steinbach, 2000). The most powerful allosteric modulation was reported for the antihelminth ivermectin on the α 7 receptor (Krause *et al.*, 1998). When exposed to low concentrations of this compound, human or chick α 7 receptors displayed an increase in their sensitivity to ACh by two orders of magnitude.

Cocaine has been found to selectively inhibit specific nAChR subtypes with a potency in the μ M range. The α 4 and β 4 subunits appear to be targeted with specific residues implicated (Francis, Vazquez, Papke, & Oswald, 2000). A structural analog of morphine, dextromethor-

phan, blocked nicotine-stimulated $Rb^{86}(+)$ efflux in a non-competitive manner in membranes prepared from a stable nAChR-transfected human embryonic kidney cell line (Hernandez *et al.*, 2000). This drug is widely used as a cough suppressant.

Another group of drugs that have been shown to interact with nicotinic receptors are antidepressants. Serotonin reuptake inhibitors such as sertraline and fluoxetine inhibit nicotinic receptor function in a non-competitive manner in TE671/RD and SHSY-5Y cell lines (Fryer & Lukas, 1999a). Bupropion and phencyclidine act in a similar manner (Fryer & Lukas, 1999b). Several of these drugs have shown promise in smoking cessation trials (Benowitz & Peng, 2000).

Many other drugs are expected to interact with the nicotinic receptors. For example, it was documented that both memantine and amantadine are extremely potent open-channel blockers at the human $\alpha 4\beta 2$ receptors (Buisson & Bertrand, 1998). Modulatory sensitivity of the neuronal nicotinic receptor family by this wide range of pharmacological agents suggests that there is much yet to learn concerning the endogenous regulation of these ion channels, and effects on release of multiple kinds of neurotransmitters in both brain and periphery.

Nicotine addiction

Understanding the processes underlying nicotine addiction requires characterization of the neuroadaptive effects of sustained exposure to nicotine, on both nicotinic receptors and the other neurotransmitter systems affected by nicotinic receptor activation and desensitization. The paradoxical up-regulation of high-affinity nicotinic receptors by nicotine has been recognized for some time in both rodent brain (Flores, Davila-Garcia, Ulrich, & Kellar, 1997; Marks, Stitzel, & Collins, 1986; Pauly, Marks, Robinson, van de Kamp, & Collins, 1996; Sanderson, Drasdo, McCrea, & Wonnacott, 1993) and more recently in human brain (Benwell, Balfour, & Anderson, 1988; Breese *et al.*, 1997b; Perry, Davila-Garcia, Stockmeier, & Kellar, 1999) and human blood leukocytes (Benhammou *et al.*, 2000). This up-regulation of receptor number is dose dependent (Benhammou *et al.*, 2000; Breese *et al.*, 1997b; Marks, Stitzel, & Collins, 1986). Low-affinity nicotinic receptors, measured by [125 I] α -bungarotoxin binding, are increased in rodents at higher nicotine concentration than required for high-affinity receptor up-regulation (Marks, Stitzel, & Collins, 1986). In human brain, the low-affinity $\alpha 7$ receptor is expressed at low levels, making accurate ligand binding determinations difficult. A study from Court *et al.* (1998) showed no increase in [125 I] α -bungarotoxin binding in smokers compared to non-smokers. However, Breese *et al.* (2000) found increased [3 H]MLA binding in smokers, although the difference between smokers and non-smokers was not significant. These results suggest that in many human smokers, nicotine concentrations in the brain may not reach levels that induce receptor up-

regulation. The actual role that receptor up-regulation in smokers plays in the process of nicotine addiction and cessation or relapse is not known. Receptor increases following chronic nicotine administration in animals do not appear to be due to increases in subunit transcription (Pauly *et al.*, 1996), but were rather attributed to increased stability of the receptor itself (Peng, Gerzanich, Anand, Whiting, & Lindstrom, 1994b). It has been suggested that the nicotine-bound receptor may assume an inactivated conformation that is more stable, but is not functional following nicotine binding (Dani & Heinemann, 1996).

Most receptor subunit combinations are generally up-regulated by chronic nicotine treatment (Bhat, Turner, Selvaag, Marks, & Collins, 1991; Peng, Gerzanich, Anand, Wang, & Lindstrom, 1997; Sanderson *et al.*, 1993). However, receptor increases have been found to be both brain region-specific and may be restricted to defined subunit combinations (Rogers, Mandelzys, Deneris, Cooper, & Heinemann, 1992). It has recently been found that the β subunit may affect desensitization and receptor up-regulation. Receptors formed from $\alpha 3\beta 2$ subunits in transfected cells are up-regulated by chronic nicotine; $\alpha 3\beta 4$ receptors desensitize to a lesser extent and are not increased by chronic nicotine in this *in vitro* model. It is clear from these results that the subunit combinations expressed can regulate tissue and cellular responses to nicotine.

Activation of nicotinic receptors evokes the release of multiple types of neurotransmitters, including dopamine, norepinephrine, GABA, glutamate, acetylcholine, and neuropeptides (Guo *et al.*, 1998; Wonnacott, 1997). It is likely that neuroadaptations of these neurotransmitter systems, including their receptors, contribute to the development of tolerance to nicotine and the maintenance of addiction (Watkins, Koob, & Markou, 2000).

Dopamine pathways in the pedunculopontine nucleus, VTA, and nucleus accumbens have been implicated in the positive reinforcement provided by nicotine (Balfour, Benwell, Birrell, Kelly, & Al-Aloul, 1998; Nisell, Nomikos, & Svensson, 1994). Nicotine stimulated release of dopamine in the nucleus accumbens is blockable by NMDA receptor antagonists, suggesting that glutamate neurotransmission is also involved in reward pathways. Differential effects of nicotine have been noted in the nucleus accumbens core and shell (Nisell, Marcus, Nomikos, & Svensson, 1997). These disparate responses may be due to reciprocal changes that occur in the medioventral shell and the laterodorsal core (Bassareo & Di Chiara, 1999). In rats chronically treated with nicotine, a sensitization to nicotine responses occurs in the core, but not in the shell. Conversely, dopamine release in accumbens, in response to acute nicotine treatment, is likely to be from the shell. The development of sensitization in the core is thought to strengthen nicotine associations leading to addiction (Balfour *et al.*, 1998; Balfour, Wright, Benwell, & Birrell, 2000; Di Chiara, 2000).

Drugs that directly affect receptor function may be useful in smoking cessation treatment. The antidepressants bupropion and nortriptyline have already been used in this regard (Balfour & Ridley, 2000; Benowitz & Peng, 2000; Prochazka *et al.*, 1998). The nicotinic receptor antagonist, mecamylamine, has also been proposed as a therapy for smoking cessation (Rose, Behm, & Westman, 1998). A recent study suggests that mecamylamine can inhibit the cardioacceleratory and epinephrine-releasing effects of nicotine, likely reducing receptor binding and distribution of coadministered nicotine (Zevin, Jacob, & Benowitz, 2000).

Cytochrome P450–2A6 (CYP2A6) is the principal cytochrome P450-metabolizing enzyme for nicotine (Messina, Tyndale, & Sellers, 1997). Subjects with null alleles for *cyp2a6* increase the bioavailability of nicotine and, thus, smoke less (Pianezza, Sellers, & Tyndale, 1998). Inhibitors of CYP2A6, such as methoxsalen, reduced nicotine metabolism and resulted in decreased smoking in a recent study, suggesting that nicotine metabolism may be an important target for development of smoking cessation interventions (Messina *et al.*, 1997; Sellers, Kaplan, & Tyndale, 2000a; Sellers, Zeman, Kaplan, & Tyndale, 2000b).

New tools for nicotinic receptor research

A need for better tools to dissect the different receptor subtypes constitutes one of the important challenges for both basic science and applied pharmaceutical research. An expanding group of toxins that have differential effects at subtypes of nicotinic receptors are the snail toxins from *Conus geographicus* better known as α -conotoxins (α -CTX). These short peptides of less than 20 amino acids display a high selectivity for the different nAChR subtypes (McIntosh, Santos, & Olivera, 1999). For example, α -CTX MI displays a high affinity for the muscle type receptor but a low affinity for the homomeric $\alpha 7$ receptor. In contrast, the α -CTX IMI from the worm *Conus imperialis* is selective for the $\alpha 7$ receptor (Johnson, Martinez, Elgoyhen, Heinemann, & McIntosh, 1995). Analysis of different toxins and their spectrum of activity on receptor subtypes suggests that α -CTX MII displays a preferential activity for $\alpha 3\beta 2$ nAChRs (Cartier *et al.*, 1996; Whiteaker, McIntosh, Luo, Collins, & Marks, 2000a). The primary mechanism accounting for the large differences observed between toxin affinities seems to reside in their specific off rates.

Extensive studies of neuromuscular transmission have revealed that numerous natural toxins are interacting with the nAChRs expressed by the muscle cell but less with ganglionic receptors. While sensitivity to the snake toxin α -bungarotoxin was used for some time as a characterizing tool for neuronal nAChR, it was later demonstrated that this snake toxin can also exhibit differences in affinity to receptor subtypes. Investigation of the short vs. long snake toxins (Erabutoxin/Cobra-toxin) revealed that, while these two toxins bind almost equally to the muscle receptor, only the long toxin

displays a high affinity at the $\alpha 7$ nAChR (Servent *et al.*, 1997).

Further investigation of natural products interacting with the nAChRs has shown that these toxins can be divided into two main classes by function, as inhibitors or activators of the receptors. Plant toxins include nicotine, cytisine, and MLA, whereas animal toxins include epibatidine and anabaseine. Originally purified from the equatorial frog skin (*Epipedobates tricolor*; Badio & Daly, 1994), epibatidine was found to be one of the most potent agonists of the heteromeric neuronal nAChRs. Epibatidine binds to several subsets of nicotine receptors with different affinities for cytisine (Marks, Smith, & Collins, 1998). Alpha-CTX MII binds with high affinity to one of these subsets with a low cytisine affinity, resolving epibatidine binding into three populations. These are the high-affinity binding sites representing $\alpha 4\beta 2$ receptors, and the two low-affinity cytisine sites identified by alpha CTX-MII, likely containing $\alpha 3$ subunits (Whiteaker *et al.*, 2000a).

Understanding the differences between an agonist and a competitive antagonist is also an important issue in the field of ligand-gated channels. The analysis *in vitro* of chimaeric receptors where presence or absence of binding sites can be engineered continues to be helpful in this regard (Corringer *et al.*, 1995). Analysis, based on the allosteric model originally formulated by Monod, Wyman, and Changeux (1965), revealed that agonists are compounds that stabilize the active state whereas competitive antagonists are defined by the functional state that they stabilize. For example, while DH β E and MLA are both competitive inhibitors of the $\alpha 7$ receptor, DH β E was shown to stabilize a desensitized state whereas MLA stabilizes the resting state (Bertrand *et al.*, 1997). Newly synthesized compounds such as epibatidine derivatives can be similarly analyzed by functional comparisons (Bertrand *et al.*, 1999; Spang *et al.*, 2000). Although presented as clear-cut differences between agonists and antagonists, it should be recalled that some compounds, including epibatidine, may display complex pharmacological patterns that vary in efficacy at different receptor subtypes. A typical and further complexity of the pharmacological profile is illustrated by the partial agonist behavior of some compounds. A recent illustration of the partial agonist effect of epibatidine was provided in studies of the rat $\alpha 4\beta 2$ nAChRs (Buisson, Vallejo, Green, & Bertrand, 2000). While of extreme importance for drug selectivity, an extensive discussion of pharmacological profile would require an entire review and is not further developed herein.

A series of genetic manipulations in mice has allowed the production of knockouts of specific nAChR genes (Orr-Urtreger *et al.*, 1997; Picciotto *et al.*, 1995; Ross *et al.*, 2000; Xu *et al.*, 1999), as well as the first knock-in of a mutated $\alpha 7$ gene (Orr-Urtreger *et al.*, 2000). These provide valuable tools for study of receptor function in live mammalian systems (Cordero-Erausquin, Marubio, Klink, & Changeux, 2000). The first published nicotinic

receptor knockout was the disruption of the gene coding for the $\beta 2$ subunit (Picciotto *et al.*, 1995). Designed to suppress the most abundant high-affinity nicotinic receptors, these studies revealed surprisingly little change in behavior of the mouse. The most important alteration, observed thus far in the $\beta 2$ knockout, is modification of passive avoidance. Further, the $\beta 2$ knockout mouse does not self-administer nicotine to the same extent as the wild-type mouse, suggesting that $\beta 2$ -containing receptors (presumably $\alpha 4\beta 2$), are likely to be important in the reinforcing properties of nicotine (Picciotto *et al.*, 1998).

Rubidium efflux studies of $\beta 2$ null-mutants show that the $\beta 2$ subunit appears to be a component of all receptors that modulate dopamine and GABA release. Acetylcholine release, however, which was detectable in the inferior colliculus and interpeduncular nucleus (IPN) of the $\beta 2$ null-mutant may be modulated by an $\alpha 3\beta 4$ -type receptor (Whiteaker *et al.*, 2000b).

The development of a null-mutant for the $\alpha 4$ gene showed that loss of this subunit results in a reduction of the nicotine antinociceptive effects (Marubio *et al.*, 1999), confirming initial observations about the antinociceptive effects of nicotinic agonists in the CNS. An independently generated knockout of the $\alpha 4$ nicotinic receptor subunit has recently been reported, and has been examined in several behavioral paradigms (Ross *et al.*, 2000). Increases in exploratory behavior, as well as heightened anxiety, were found in these mice. It was suggested that the $\alpha 4$ subunit might be required for activation of inhibitory neural circuits. As anxiety and motor restlessness are associated with nicotine withdrawal (Hughes, Higgins, & Bickel, 1994; Jorenby *et al.*, 1996; McKenna & Cox, 1992; Shiffman *et al.*, 2000), this mouse may be useful in drug development for smoking cessation.

Knockouts of $\alpha 3$ and $\alpha 7$ genes have also been generated (Orr-Urtreger *et al.*, 1997; Xu *et al.*, 1999). Suppression of $\alpha 3$ gene expression considerably reduced survival; only one-third of the homozygous littermates survived but displayed megacystis as well as mydriasis. Attributed to the alteration of the ganglionic nicotinic receptor activity, these impairments caused marked modification of the bladder size, apparition of bladder stones, and incontinence (De Biasi, Nigro, & Xu, 2000; Xu *et al.*, 1999). Mice lacking the $\alpha 7$ -subunit gene show only subtle phenotypic changes. The baroreflex appears to be affected (Franceschini *et al.*, 2000) and the mice exhibit increased hippocampal excitability (Orr-Urtreger *et al.*, 1997).

While conditional knockouts may provide a better insight into the role of nicotinic receptors in the CNS, alternative information can be obtained using knock-ins with mutated receptors. A mutation in the MII region of the $\alpha 7$ -receptor subunit, equivalent to the chick L247T mutation originally described by Revah *et al.* (Bertrand *et al.*, 1992; Revah *et al.*, 1991), has been recently introduced into a tissue culture cell line, resulting in cell death (apparently due to a gain of function that causes an

excessive entry of calcium; Briggs *et al.*, 1999). Treatment of a cultured fetal hippocampal cell line with nicotine has been shown to induce apoptosis, mediated by the $\alpha 7$ nicotinic receptor (Berger *et al.*, 1998). The mechanism was again thought to be calcium mediated, due to a lack of calcium buffering in the fetal cells, suggesting that nicotine is apoptotic at very early embryonic stages and may be neuroprotective later in development. The $\alpha 7$ MII mutation has recently been introduced into a transgenic animal where it results in lethality in the first 24 h of life (Orr-Urtreger *et al.*, 2000). Electrophysiological investigations of cultured hippocampal neurons from these mice confirmed the gain of function in the homozygous animal with an increase of the ACh sensitivity, decrease of desensitization, and conversion of competitive antagonists into agonists. These results suggest that regulated expression of the $\alpha 7$ receptor during development may be important for programmed cell death, and perhaps neuronal migration. In that regard, prenatal administration of nicotine has been shown to induce aberrant lung development. In a primate model, prenatal nicotine resulted in lung hypoplasia and reduced surface complexity of the developing alveoli. Increases in expression of $\alpha 7$ -containing receptors were also noted (Sekhon *et al.*, 1999).

To examine in more detail the contribution of $\alpha 9$ in hearing function, Vetter and collaborators have developed a mouse knockout of this gene (Vetter *et al.*, 1999). Morphological evidence has confirmed a localization of the $\alpha 9$ receptor in the outer cochlear hair cells (Elgoyhen *et al.*, 1994). In the $\alpha 9$ knockout, mature innervation of these hair cells was lost and resulted in the failure of suppression of cochlear responses, following fiber activation.

The importance of nAChRs in disease

Nicotinic receptors may play a role in several human diseases, as well as in cognition. If not causal, the involvement of nicotinic receptor function, in many cases, is likely to be important. The first disease in which nicotinic receptors were carefully studied was myasthenia gravis (Beeson *et al.*, 1998; Croxen *et al.*, 1997; Vincent *et al.*, 1998). In this autoimmune disease, antibodies against the muscle receptors are developed causing either a progressive reduction of the receptor number or blockade of neurotransmission. In both cases muscle weakness with excessive muscle fatigue is observed together with impairment of muscle control. A few rare cases of genetically transmissible myasthenia gravis have been reported, and to date at least 14 mutations have been found for the muscle receptors in this disease (Engel *et al.*, 1996; Ohno *et al.*, 1995, 1996, 1997; Ohno, Anlar, Ozdirim, Brengman & Engel, 1998). Electrophysiological investigations of receptors reconstituted with these altered subunits have further revealed that mutations are accompanied by a reduction of receptor function that can explain the diminution of synaptic transmission.

Genetic linkage of a disease to the chromosomal map site of a nicotinic receptor has been used to implicate nicotinic receptor roles in two diseases, autosomal dominant nocturnal frontal lobe epilepsy (ADNFLE) and schizophrenia. A gene locus was assigned for ADNFLE to chromosome 20q13.2 the locus of the $\alpha 4$ nicotinic receptor subunit. Subsequently a point mutation leading to a change in an amino acid (Ser248Phe) was identified in the second transmembrane region of the $\alpha 4$ subunit in the linkage family (Steinlein *et al.*, 1995). Another $\alpha 4$ -Ser248Phe-linked ADNFLE family has been found, but was genetically distinct from the first pedigree, based on other inherited markers (Steinlein *et al.*, 2000). A second type of mutant with an insertion of an amino acid into the extracellular end of the second transmembrane region of CHR4 that reduces the calcium permeability has been identified (Steinlein *et al.*, 1997). Both types of mutants lead to hypoactivity of $\alpha 4$ -containing nAChRs (Bertrand, Weiland, Berkovic, Steinlein, & Bertrand, 1998). While supporting the linkage between a mutation in the $\alpha 4$ subunit and ADNFLE, these data also illustrate the importance of the second transmembrane segment. Another epilepsy, juvenile myoclonic epilepsy (JME) has been linked to the gene locus of the $\alpha 7$ nicotinic receptor on chromosome 15q14 (Elmslie *et al.*, 1997). JME is a distinctive and common form of familial idiopathic generalized epilepsy.

The locus of the $\alpha 7$ -subunit gene at 15q13-q14 (Table 1) has also been genetically linked to schizophrenia in several independent studies. A polymorphic marker, less than 100 kb from the $\alpha 7$ nicotinic receptor gene, was linked [logarithm of the odds ratio (lod) score of 5.3, $\Theta = 0.00$] to an auditory gating deficit in schizophrenia (Freedman *et al.*, 1997). Linkage to schizophrenia was also found in a second cohort by the same group (Leonard *et al.*, 1998a), and has been replicated in several laboratories (Craddock *et al.*, 1999; Kaufmann *et al.*, 1998; Riley *et al.*, 2000; Stassen *et al.*, 2000). The $\alpha 7$ gene is partially duplicated and the duplication maps within the linkage region on chromosome 15, making it a candidate gene also (Gault *et al.*, 1998). Schizophrenic subjects have been reported to have reduced expression of the $\alpha 7$ -receptor subunit in hippocampus (Freedman, Hall, Adler, & Leonard, 1995), in the RTN (Court *et al.*, 1999) and in frontal cortex (Guan, Zhang, Blennow, & Nordberg, 1999). It is of interest that a related disorder, manic depression, has recently been genetically linked to this same locus (Edenberg *et al.*, 1997).

Schizophrenics have also been shown to have reduced levels of high-affinity nicotinic receptors in several brain regions, as measured by either [3 H]nicotine (Breese *et al.*, 2000) or by [3 H]cytisine (Durany *et al.*, 2000; Freedman *et al.*, 1995). These decreases are not likely to be a result of typical neuroleptic medication (Lee, Breese, & Leonard, 2000). The low levels of both high- and low-affinity nicotinic receptors in schizophrenia may be related to the very high incidence of smoking seen in this disease (Dalack, Healy, & Meador-Woodruff, 1998; Glassman, Covey, Dalack, & Stetner, 1992; Hughes,

Hatsukami, Mitchell, & Dahlgren, 1986; Leonard *et al.*, 1998b). Nicotine normalizes a sensory gating deficit found in schizophrenics that is thought to be related to their inability to filter out extraneous or intrusive information in their environment (Adler, Hoffer, Griffith, Waldo, & Freedman, 1992; Adler, Hoffer, Wiser, & Freedman, 1993; Adler, Smas, Fiandaca, Frim, & Majzoub, 1990; Freedman, Waldo, Bickford-Wimer, & Nagamoto, 1991). It has been suggested that smoking is, thus, an attempt at self-medication in which patients are stimulating a reduced population of nicotinic receptors (Adler *et al.*, 1998; Leonard *et al.*, 1998b). The atypical neuroleptic clozapine, which also normalizes this gating deficit (Nagamoto, Adler, Hea, Griffith, McRae, & Freedman, 1996), appears to decrease smoking in schizophrenic patients (McEvoy, Freudenreich, McGee, Vanderzwaag, Levin, & Rose, 1995). The incidence of smoking in depression is also higher than in subjects with no mental illness and smoking cessation attempts are less successful (Glassman *et al.* 1990, 1992). In depression, the effects of nicotine are thought to involve modulation of stress-induced responses that return when the subject is not smoking (Balfour & Ridley, 2000).

Compounds that may be therapeutically specific for the $\alpha 7$ nAChR have been isolated as derivatives of the snail toxin anabaseine, such as GTS-21 (Li, Papke, He, Millard, & Meyer, 1999; Meyer, King, & Meyers, 1998; Meyer *et al.*, 1997; van Haaren, Anderson, Haworth, & Kem, 1999). These drugs appear to protect cells from a range of necrotic and apoptotic signals by increasing calcium influx over a narrow concentration range, and mediating the activation of protein kinase C. GTS-21 has been found to normalize an auditory evoked-potential deficit in mice that have decreased levels of $\alpha 7$ expression (Stevens, Kem, Mahnir, & Freedman, 1998), similar to the auditory deficit found in most schizophrenics that is normalized by nicotine (Adler *et al.*, 1992, 1993; Freedman *et al.*, 1991; Leonard *et al.*, 1998b). These results suggest that GTS-21 might also normalize gating deficits in human subjects.

The nicotinic receptor gene family may be important in both Alzheimer's and Parkinson's diseases. Expression, of both high- and low-affinity nicotinic receptor subunits, declines with aging (Court *et al.*, 1997; Marutle, Warpman, Bogdanovic, & Nordberg, 1998). In Alzheimer's disease, however, there are further selective reductions that appear to be region specific. The $\alpha 4$ subunit was found to be reduced in temporal cortex, but $\alpha 7$ was selectively lost from the RTN where $\alpha 7$ is particularly abundant (Court *et al.*, 1999). High-affinity binding to $\alpha 7$ receptors of the β -amyloid peptide (A β 1–42) suggests a role for nicotinic receptors in Alzheimer's disease (Wang *et al.*, 2000a; Wang, Lee, Davis, & Shank, 2000b). Nicotinic-receptor expression is also affected in Parkinson's disease (Perry *et al.*, 1995). In MPTP-treated squirrel monkeys (*Saimiri sciureus*), a Parkinson's disease model, $\alpha 6$ and $\beta 3$ subunits were selectively decreased, suggesting that receptor ligands directed to these subunits might be useful in treatment of this disorder (Quik & Jeyarasasingam, 2000).

Tourette's syndrome responds to transdermal nicotine therapy and the treatment appears to be long lasting. Motor and vocal tics present in the disease are decreased by nicotine given as either gum or patch (Shytle, Silver, Philipp, McConville, & Sanberg, 1996), and the effects can last up to 4 weeks following the drug application (Dursun & Reveley, 1997). Recently, a 2-year trial using the high-affinity antagonist mecamlamine has shown significant effect on tics and other behavioral disturbances in Tourette's, suggesting that peripheral nicotinic receptor antagonists may be useful in this disease (Silver, Shytle, & Sanberg, 2000).

In a common smoking-associated lung cancer, small cell lung carcinoma (SCLC), the release of autocrine growth factors serotonin and mammalian bombesin/gastrin-releasing peptide are regulated by a neuronal nicotinic receptor containing $\alpha 7$ subunits (Codignola *et al.*, 1994; Schuller, 1991; Schuller, Nylén, Park, & Becker, 1990). Both nicotine and NNK, a nicotine-derived nitrosamine, increased Ca^{2+} influx and activated a Raf-1 MAP kinase pathway in fetal pulmonary neuroendocrine and SCLC cells in culture, leading to increased DNA synthesis. Thus, *in utero* exposure of the fetal lung to nicotine or NNK could be an etiological factor in the development of smoking-associated pediatric lung disorders.

Cognition is likely to also be a target for therapeutic approaches involving the nicotinic receptor gene family. Although improvement of cognitive performance by nicotine in both human and rat is widely accepted, its mode of action still remains obscure. Responses to visual stimuli are improved by nicotine (Levin, Bettgeowda, Weaver, & Christopher, 1998; Levin & Simon, 1998). Nicotine patches at 7 mg/day in human subjects significantly reduced the number of errors and increased consistency of cognitive responses. Nicotine has also been shown to produce a small but significant improvement in the overall cognitive performance of Alzheimer's patients (White & Levin, 1999; Potter *et al.*, 1999). The competitive antagonist DH β E at 6 mg/kg blocks the effects induced by 0.05–0.15 mg/kg of nicotine on attention in an animal model, a nicotine concentration compatible with that found in human plasma (Mirza & Stoleran, 1998; Shoaib & Stoleran, 1999; Stoleran, Mirza, Hahn, & Shoaib, 2000). In contrast, MLA, a specific competitive antagonist of $\alpha 7$ receptors, was inefficient, indicating that nicotine effects were probably mediated through high-affinity receptors.

Two compounds from Targacept (Winston-Salem, North Carolina), RJR-2483 and RJR-1784 have effects on memory. Broad screening of these substances revealed that RJR-2483 resembles nicotine, is a full agonist at the $\alpha 4\beta 2$ nAChR, improves both short- and long-term memory and has long-lasting effects (Abdulla *et al.*, 1996; Papke, Bencherif, & Lippiello, 1996; Summers, Lippiello, & Giacobini, 1996; Bencherif *et al.*, 1996). Additionally, at 0.1 $\mu\text{M/kg}$, RJR-1734 causes a major improvement in the performance of rats placed in a radial arm maze compared to RJR-2483. Effects lasting

as long as 18 h following a single oral administration could be observed, supporting the hypothesis that compounds acting on the nicotinic receptor may become of clinical relevance. Compounds from SIBIA may also be effective in cognition (Cosford *et al.*, 1996; Sacaan *et al.*, 1997). SIB-1508 displayed a higher specificity for $\alpha 4\beta 2$ receptors than did SIB-1553. The latter compound is more selective for the $\alpha 4\beta 4$ nAChRs. Broad cognitive improvements have been reported for the SIB-1553, whereas SIB-1508 could bring more benefit in the control of locomotor activity.

Since it is known that the ligand-binding pocket in the nAChRs is made at the interface between the α subunit and its adjacent subunit, it is reasonable to think that ligands specific for a given receptor composition can be developed (Corringer *et al.*, 1995). Compounds with an inter-nitrogen distance of 4.79 Å have been examined (Decker & Meyer, 1999). One group of derivatives was constructed from the 3-pyridyl-ethers which conserve some of the features of the natural agonist epibatidine. These compounds display a subnanomolar affinity for $\alpha 3$ - or $\alpha 4$ -containing receptors, comparable to that of the natural toxin epibatidine (Abreo *et al.*, 1996; Badio & Daly, 1994). Design of drugs specifically targeted to one receptor subtype is well illustrated by the identification of ABT594 (Bannon *et al.*, 1998). Designed to mimic the antinociceptive effects of epibatidine while displaying lower toxicity, ABT594 was shown to significantly relieve pain in animal models. Contribution of the $\alpha 4\beta 2$ receptor in the pain pathway was later confirmed in the $\alpha 4$ knockout mouse model (Marubio *et al.*, 1999).

Future prospects

Although progress in understanding the role that neuronal nAChRs play in neuronal function, nicotine addiction, and disease has been remarkable, a monumental task yet remains for this field. An exhaustive search must be made for other receptor subunits, both in the brain and in the periphery, before we can begin to assign specific functions to subunit combinations *in vivo*. Continued development of new ligands with defined specificities for subunit combinations will provide new tools for pharmacologists and for informed drug development for smoking cessation. The study of gene regulation for this family of receptor subunits is just beginning, as our knowledge of promoter structure and transcription factor interactions grows. Lastly, the finding of nicotinic receptor involvement in multiple illnesses suggests that nicotinic-receptor mutations might play a role through interaction with other gene products and environmental factors to define a disease.

References

- Abdulla FA, Bradbury E, Calaminici MR, Lippiello PM, Wonnacott S, Gray JA, Sinden JD. 1996. Relationship between up-regulation of nicotine binding sites in rat brain and delayed cognitive enhancement observed after chronic or acute nicotinic receptor stimulation. *Psychopharmacology* 124:323–331.

- Abreo MA, Lin NH, Garvey DS, Gunn DE, Hettinger AM, Wasicak JT, Pavlik PA, Martin YC, Donnelly-Roberts DL, Anderson DJ, Sullivan JP, Williams M, Arneric SP, Holladay MW. 1996. Novel 3-pyridyl ethers with subnanomolar affinity for central neuronal nicotinic acetylcholine receptors. *Journal of Medicinal Chemistry* 39:817–825.
- Adler GK, Smas CM, Fiandaca M, Frim DM, Majzoub JA. 1990. Regulated expression of the human corticotropin releasing hormone gene by cyclic AMP. *Molecular and Cellular Endocrinology* 70:165–174.
- Adler LE, Hoffer LJ, Griffith J, Waldo MC, Freedman R. 1992. Normalization by nicotine of deficient auditory sensory gating in the relatives of schizophrenics. *Biological Psychiatry* 32:607–616.
- Adler LE, Hoffer LD, Wiser A, Freedman R. 1993. Normalization of auditory physiology by cigarette smoking in schizophrenic patients. *American Journal of Psychiatry* 150:1856–1861.
- Adler LE, Olincy A, Waldo MC, Harris JG, Griffith J, Stevens K, Flach K, Nagamoto H, Bickford P, Leonard S, Freedman R. 1998. Schizophrenia, sensory gating, and nicotinic receptors. *Schizophrenia Bulletin* 24:189–202.
- Agulhon C, Charnay Y, Vallet P, Abitbol M, Kobetz A, Bertrand D, Malafosse A. 1998. Distribution of mRNA for the alpha4 subunit of the nicotinic acetylcholine receptor in the human fetal brain. *Brain Research, Molecular Brain Research* 58:123–131.
- Agulhon C, Abitbol M, Bertrand D, Malafosse A. 1999. Localization of mRNA for CHRNA7 in human fetal brain. *Neuroreport* 10:2223–2227.
- Albuquerque EX, Pereira EF, Castro NG, Alkondon M, Reinhardt S, Schroder H, Maelicke A. 1995. Nicotinic receptor function in the mammalian central nervous system. *Annals of the New York Academy of Sciences* 757:48–72.
- Albuquerque EX, Pereira EF, Alkondon M, Schratzenholz A, Maelicke A. 1997. Nicotinic acetylcholine receptors on hippocampal neurons: distribution on the neuronal surface and modulation of receptor activity. *Journal of Receptor & Signal Transduction Research* 17:243–266.
- Alkondon M, Albuquerque EX. 1993. Diversity of nicotinic acetylcholine receptors in rat hippocampal neurons. I. Pharmacological and functional evidence for distinct structural subtypes. *Journal of Pharmacology & Experimental Therapeutics* 265:1455–1473.
- Alkondon M, Albuquerque EX. 1995. Diversity of nicotinic acetylcholine receptors in rat hippocampal neurons. III. Agonist actions of the novel alkaloid epibatidine and analysis of type II current. *Journal of Pharmacology & Experimental Therapeutics* 274:771–782.
- Alkondon M, Reinhardt S, Lobron C, Hermsen B, Maelicke A, Albuquerque EX. 1994. Diversity of nicotinic acetylcholine receptors in rat hippocampal neurons. II. The rundown and inward rectification of agonist-elicited whole-cell currents and identification of receptor subunits by in situ hybridization. *Journal of Pharmacology & Experimental Therapeutics* 271:494–506.
- Alkondon M, Rocha ES, Maelicke A, Albuquerque EX. 1996. Diversity of nicotinic acetylcholine receptors in rat brain. V. Alpha-Bungarotoxin-sensitive nicotinic receptors in olfactory bulb neurons and presynaptic modulation of glutamate release. *Journal of Pharmacology & Experimental Therapeutics* 278:1460–1471.
- Alkondon M, Pereira EF, Cortes WS, Maelicke A, Albuquerque EX. 1997. Choline is a selective agonist of alpha7 nicotinic acetylcholine receptors in the rat brain neurons. *European Journal of Neuroscience* 9:2734–2742.
- Alkondon M, Pereira EF, Albuquerque EX. 1998. Distinct nicotinic receptors (nAChRs) modulate GABA release from CA1 interneurons in rat hippocampal slices. *Society for Neuroscience Abstract* 24:1581.
- Alkondon M, Pereira EF, Eisenberg HM, Albuquerque EX. 1999. Choline and selective antagonists identify two subtypes of nicotinic acetylcholine receptors that modulate GABA release from CA1 interneurons in rat hippocampal slices. *Journal of Neuroscience* 19:2693–2705.
- Amador M, Dani JA. 1995. Mechanism for modulation of nicotinic acetylcholine receptors that can influence synaptic transmission. *Journal of Neuroscience* 15:4525–4532.
- Ascher P, Large WA, Rang HP. 1979. Studies on the mechanism of action of acetylcholine antagonists on rat parasympathetic ganglion cells. *Journal of Physiology* 295:139–170.
- Badio B, Daly JW. 1994. Epibatidine, a potent analgesic and nicotinic agonist. *Molecular Pharmacology* 45:563–569.
- Balfour DJ, Benwell ME, Birrell CE, Kelly RJ, Al-Aloul M. 1998. Sensitization of the mesoaccumbens dopamine response to nicotine. *Pharmacology, Biochemistry & Behavior* 59:1021–1030.
- Balfour DJK, Ridley DL. 2000. The effects of nicotine on neural pathways implicated in depression: a factor in nicotine addiction? *Pharmacology Biochemistry & Behavior* 66:79–85.
- Balfour DJK, Wright AE, Benwell ME, Birrell CE. 2000. The putative role of extra-synaptic mesolimbic dopamine in the neurobiology of nicotine dependence. *Behavioural Brain Research* 113:73–83.
- Ballivet M, Alliod C, Bertrand S, Bertrand D. 1996. Nicotinic acetylcholine receptors in the nematode *Caenorhabditis elegans*. *Journal of Molecular Biology* 258:261–269.
- Bannon AW, Decker MW, Holladay MW, Curzon P, Donnelly-Roberts D, Puttfarcken PS, Bitner RS, Diaz A, Dickenson AH, Porsolt RD, Williams M, Arneric SP. 1998. Broad-spectrum, non-opioid analgesic activity by selective modulation of neuronal nicotinic acetylcholine receptors. *Science* 279:77–81.
- Bassareo V, Di Chiara G. 1999. Differential responsiveness of dopamine transmission to food-stimuli in nucleus accumbens shell/core compartments. *Neuroscience* 89:637–641.
- Beeson D, Brydson M, Betty M, Jeremiah S, Povey S, Vincent A, Newsom-Davis J. 1993. Primary structure of the human muscle acetylcholine-receptor — cDNA cloning of the gamma-subunit and epsilon-subunit. *European Journal of Biochemistry* 215:229–238.
- Beeson D, Jeremiah S, West LF, Povey S, Newsom-Davis J. 1990. Assignment of the human nicotinic acetylcholine receptor genes: the alpha and delta subunit genes to chromosome 2 and the beta subunit gene to chromosome 17. *Annals of Human Genetics* 54:199–208.
- Beeson D, Newland C, Croxen R, Buckel A, Li FY, Larsson C, Tariq M, Vincent A, Newsom-Davis J. 1998. Congenital myasthenic syndromes. Studies of the AChR and other candidate genes. *Annals of the New York Academy of Sciences* 841:181–183.
- Bencherif M, Lovette ME, Fowler KW, Arrington S, Reeves Caldwell WS, Lippello, PM. 1996. RJR-2403: a nicotinic agonist with CNS selectivity I. In vitro characterization. *Journal of Pharmacology & Experimental Therapeutics* 279:1413–1421.
- Benhammou K, Lee MJ, Strook M, Sullivan B, Logel J, Raschen K, Gotti C, Leonard S. 2000. [³H]Nicotine binding in peripheral blood cells of smokers is correlated with the number of cigarettes smoked per day. *Neuropharmacology* 39:2818–2829.
- Benowitz NL, Peng, MW. 2000. Non-nicotine pharmacotherapy for smoking cessation — mechanisms and prospects. *CNS Drugs* 13:265–285.
- Benwell ME, Balfour DJ, Anderson JM. 1988. Evidence that tobacco smoking increases the density of (–)[³H]nicotine binding sites in human brain. *Journal of Neurochemistry* 50:1243–1247.
- Berger F, Gage FH, Vijayaraghavan S. 1998. Nicotinic receptor-induced apoptotic cell death of hippocampal progenitor cells. *Journal of Neuroscience* 18:6871–6881.
- Bertrand D, Changeux JP. 1992. Nicotinic receptor: An allosteric protein specialized for intercellular communication. *Seminar in Neuroscience* 7:75–90.
- Bertrand D, Ballivet M, Rungger D. 1990. Activation and blocking of neuronal nicotinic acetylcholine receptor reconstituted in *Xenopus* oocytes. *Proceedings of the National Academy of Sciences USA* 87:1993–1997.
- Bertrand D, Valera S, Bertrand S, Ballivet M, Rungger D. 1991. Steroids inhibit nicotinic acetylcholine-receptors. *Neuroreport* 2:277–280.
- Bertrand D, Devillers-Thiery A, Revah F, Galzi J-L, Hussy N, Mulle C, Bertrand S, Ballivet M, Changeux J-P. 1992. Unconventional pharmacology of a neuronal nicotinic receptor mutated in the channel domain. *Proceedings of the National Academy of Sciences USA* 89:1261–1265.
- Bertrand D, Galzi JL, Devillers TA, Bertrand S, Changeux JP. 1993. Mutations at two distinct sites within the channel domain M2 alter calcium permeability of neuronal alpha 7 nicotinic receptors. *Proceedings of the National Academy of Sciences USA* 90:6971–6975.
- Bertrand S, Devillers-Thiery A, Palma E, Buisson B, Edelstein SJ, Corringer PJ, Changeux JP, Bertrand D. 1997. Paradoxical allosteric effects of competitive inhibitors on neuronal alpha7 nicotinic receptor mutants. *Neuroreport* 8:3591–3596.
- Bertrand S, Weiland S, Berkovic SF, Steinlein OK, Bertrand D. 1998. Properties of neuronal nicotinic acetylcholine receptor mutants from humans suffering from autosomal dominant nocturnal frontal lobe epilepsy. *British Journal of Pharmacology* 125:751–760.
- Bertrand S, Patt JT, Spang JE, Westera G, Schubiger PA, Bertrand D. 1999. Neuronal nAChR stereoselectivity to non-natural epibatidine derivatives. *FEBS Letters* 450:273–279.

- Bhat RV, Turner SL, Selvaag SR, Marks MJ, Collins AC. 1991. Regulation of brain nicotinic receptors by chronic agonist infusion. *Journal of Neurochemistry* 56:1932-1939.
- Bigger CB, Casanova EA, Gardner PD. 1996. Transcriptional regulation of neuronal nicotinic acetylcholine receptor genes. Functional interactions between Sp1 and the rat beta4 subunit gene promoter. *Journal of Biological Chemistry* 271:32842-32848.
- Bigger CB, Melnikova IN, Gardner PD. 1997. Sp1 and Sp3 regulate expression of the neuronal nicotinic acetylcholine receptor beta4 subunit gene. *Journal of Biological Chemistry* 272:25976-25982.
- Blount P, Merlie JP. 1990. Mutational analysis of muscle nicotinic acetylcholine receptor subunit assembly. *Journal of Cell Biology* 111:2613-2622.
- Bordey A, Feltz P, Trouslard J. 1996. Nicotinic actions on neurones of the central autonomic area in rat spinal cord slices. *Journal of Physiology* 497:175-187.
- Boulter J, Connolly J, Deneris E, Goldman D, Heinemann S, Patrick J. 1987. Functional expression of two neuronal nicotinic acetylcholine receptors from cDNA clones identifies a gene family. *Proceedings of the National Academy of Sciences USA* 84:7763-7767.
- Boulter J, O'Shea-Greenfield A, Duvoisin RM, Connolly JG, Wada E, Jensen A, Gardner PD, Ballivet M, Deneris ES, McKinnon D, Heinemann S, Patrick J. 1990. α -3, α -5 and β -4: Three members of the rat neuronal nicotinic acetylcholine receptor-related gene family form a gene cluster. *Journal of Biological Chemistry* 265:4472-4482.
- Breese CR, Adams C, Logel J, Drebing C, Rollins Y, Barnhart M, Sullivan B, DeMasters BK, Freedman R, Leonard S. 1997a. Comparison of the regional expression of nicotinic acetylcholine receptor α 7 mRNA and [125 I]- α -bungarotoxin binding in human postmortem brain. *Journal of Comparative Neurology* 387:385-398.
- Breese CR, Marks MJ, Logel J, Adams CE, Sullivan B, Collins AC, Leonard S. 1997b. Effect of smoking history on [3 H]nicotine binding in human postmortem brain. *Journal of Pharmacology & Experimental Therapeutics* 282:7-13.
- Breese CR, Lee MJ, Adams CE, Sullivan B, Logel J, Gillen KM, Marks MJ, Collins AC, Leonard S. 2000. Abnormal regulation of the high affinity nicotinic receptors in subjects with schizophrenia. *Neuropsychopharmacology* 23:351-364.
- Briggs CA, McKenna DG, Monteggia LM, Touma E, Roch JM, Arneric SP, Gopalakrishnan M, Sullivan JP. 1999. Gain of function mutation of the alpha 7 nicotinic receptor: distinct pharmacology of the human alpha 7V274T variant. *European Journal of Pharmacology* 366:301-308.
- Brownlee DJA, Fairweather I. 1999. Exploring the neurotransmitter labyrinth in nematodes. *Trends in Neurosciences* 22:16-24.
- Buisson B, Bertrand D. 1998. Open-channel blockers at the human alpha 4 beta 2 neuronal nicotinic acetylcholine receptor. *Molecular Pharmacology* 53:555-563.
- Buisson B, Gopalakrishnan M, Arneric SP, Sullivan JP, Bertrand D. 1996. Human alpha4beta2 neuronal nicotinic acetylcholine receptors in HEK 293 cells: A patch-clamp study. *Journal of Neuroscience* 16:7880-7891.
- Buisson B, Gopalakrishnan M, Bertrand D. 1999. Stable expression of human neuronal nicotinic receptors. In: Arneric SP, Brioni JD, eds. *Neuronal Nicotinic Acetylcholine Receptors: Pharmacology And Therapeutic Opportunities*. New York: Wiley-Liss, Inc., pp. 99-124.
- Buisson B, Vallejo YF, Green WN, Bertrand D. 2000. The unusual nature of epibatidine responses at the alpha4beta2 nicotinic acetylcholine receptor. *Neuropharmacology* 39:2561-2569.
- Bullock AE, Clark AL, Grady SR, Robinson SF, Slobe BS, Marks MJ, Collins AC. 1997. Neurosteroids modulate nicotinic receptor function in mouse striatal and thalamic synaptosomes. *Journal of Neurochemistry* 68:2412-2423.
- Carrasco-Serrano C, Campos-Caro A, Viniegra S, Ballesta JJ, Criado M. 1998. GC- and E-box motifs as regulatory elements in the proximal promoter region of the neuronal nicotinic receptor alpha7 subunit gene. *Journal of Biological Chemistry* 273:20021-20028.
- Carrasco-Serrano C, Viniegra S, Ballesta JJ, Criado M. 2000. Phorbol ester activation of the neuronal nicotinic acetylcholine receptor alpha 7 sub-unit gene: Involvement of transcription factor Egr-1. *Journal of Neurochemistry* 74:932-939.
- Cartier GE, Yoshikami D, Gray WR, Luo S, Olivera BM, McIntosh JM. 1996. A new alpha-conotoxin which targets alpha3beta2 nicotinic acetylcholine receptors. *Journal of Biological Chemistry* 271:7522-7528.
- Chang KT, Berg DK. 1999. Nicotinic acetylcholine receptors containing alpha7 subunits are required for reliable synaptic transmission *in situ*. *Journal of Neuroscience* 19:3701-3710.
- Chen D, Dang H, Patrick JW. 1998. Contributions of N-linked glycosylation to the expression of a functional alpha7-nicotinic receptor in *Xenopus* oocytes. *Journal of Neurochemistry* 70:349-357.
- Chini B, Raimond E, Elgoyhen AB, Moralli D, Balzaretti M, Heinemann S. 1994. Molecular cloning and chromosomal localization of the human α 7-nicotinic receptor subunit gene (CHRNA7). *Genomics* 19:379-381.
- Cimino M, Marini P, Fornasari D, Cattabeni F, Clementi F. 1992. Distribution of nicotinic receptors in cynomolgus monkey brain and ganglia - localization of alpha-3 subunit messenger-RNA, alpha-bungarotoxin and nicotine binding-sites. *Neuroscience* 51:77-86.
- Codignola A, Tarroni P, Cattaneo MG, Vicentini LM, Clementi F, Sher E. 1994. Serotonin release and cell proliferation are under the control of α -bungarotoxin-sensitive nicotinic receptors in small-cell lung carcinoma cell lines. *FEBS Letters* 342:286-290.
- Conroy, WG, Berg, DK. 1999. Rapsyn variants in ciliary ganglia and their possible effects on clustering of nicotinic receptors. *Journal of Neurochemistry* 73:1399-1408.
- Conti-Fine BM, Navaneetham D, Lei S, Maus ADJ. 2000. Neuronal nicotinic receptors in non-neuronal cells: new mediators of tobacco toxicity? *European Journal of Pharmacology* 393:279-294.
- Cooper ST, Harkness PC, Baker ER, Millar NS. 1999. Up-regulation of cell-surface alpha 4 beta 2 neuronal nicotinic receptors by lower temperature and expression of chimeric subunits. *Journal of Biological Chemistry* 274:27145-27152.
- Cordero-Erausquin M, Marubio LM, Klink R, Changeux J-P. 2000. Nicotinic receptor function: new perspectives from knockout mice. *Trends in Pharmacological Sciences* 21:211-217.
- Corringer PJ, Galzi JL, Eisele JL, Bertrand S, Changeux JP, Bertrand D. 1995. Identification of a new component of the agonist binding site of the nicotinic alpha 7 homooligomeric receptor. *Journal of Biological Chemistry* 270:11749-11752.
- Corringer PJ, Le Novère N, Changeux J-P. 2000. Nicotinic receptors at the amino acid level. *Annual Review of Pharmacology & Toxicology* 40:431-458.
- Cosford ND, Bleicher L, Herbaut A, McCallum JS, Vernier JM, Dawson H, Whitten JP, Adams P, Chavez-Noriega L, Correa LD, Crona JH, Mahaffy LS, Menzaghi F, Rao TS, Reid R, Sacca AI, Santori E, Stauderman KA, Whelan K, Lloyd GK, McDonald IA. 1996. (S)-(-)-5-ethynyl-3-(1-methyl-2-pyrrolidinyl)pyridine malate (SIB-1508Y): a novel anti-parkinsonian agent with selectivity for neuronal nicotinic acetylcholine receptors. *Journal of Medicinal Chemistry* 39:3235-3237.
- Court J, Spurdin D, Lloyd S, McKeith I, Ballard C, Cairns N, Kerwin R, Perry R, Perry E. 1999. Neuronal nicotinic receptors in dementia with Lewy bodies and schizophrenia: alpha-bungarotoxin and nicotine binding in the thalamus. *Journal of Neurochemistry* 73:1590-1597.
- Court JA, Lloyd S, Johnson M, Griffiths M, Birdsall NJM, Piggott MA, Oakley AE, Ince PG, Perry EK, Perry RH. 1997. Nicotinic and muscarinic cholinergic receptor binding in the human hippocampal formation during development and aging. *Developmental Brain Research* 101:93-105.
- Court JA, Lloyd S, Thomas N, Piggott MA, Marshall EF, Morris CM, Lamb H, Perry RH, Johnson M, Perry EK. 1998. Dopamine and nicotinic receptor binding and the levels of dopamine and homovanillic acid in human brain related to tobacco use. *Neuroscience* 87:63-78.
- Couturier S, Erkman L, Valera S, Rungger D, Bertrand S, Boulter J, Ballivet M, Bertrand D. 1990a. Alpha 5, alpha 3, and non-alpha 3. Three clustered avian genes encoding neuronal nicotinic acetylcholine receptor-related subunits. *Journal of Biological Chemistry* 265:17560-17567.
- Couturier S, Bertrand D, Matter J-M, Hernandez M-C, Bertrand S, Millar N, Valera S, Barkas T, Ballivet M. 1990b. A neuronal nicotinic acetylcholine receptor subunit α 7 is developmentally regulated and forms a homo-oligomeric channel blocked by α -BTX. *Neuron* 5:847-856.
- Craddock N, Lendon C, Cichon S, Culverhouse R, Detera-Wadleigh S, Devon R, Faraone S, Foroud T, Gejman P, Leonard S, McInnis M, Owen MJ, Riley B, Armstrong C, Barden N, van Broeckhoven C, Ewald H, Folstein S, Gerhard D, Goldman D, Gurling H, Kelsoe J, Levinson D, Muir W, Philippe A. 1999. Chromosome Workshop: Chromosomes 11, 14, and 15. *American Journal of Medical Genetics* 88:244-254.

- Croxen R, Newland C, Beeson D, Oosterhuis H, Chauplannaz G, Vincent A, Newsom-Davis J. 1997. Mutations in different functional domains of the human muscle acetylcholine receptor alpha subunit in patients with the slow-channel congenital myasthenic syndrome. *Human Molecular Genetics* 6:767-774.
- Cuevas J, Adams DJ. 2000. Substance P preferentially inhibits large conductance nicotinic ACh receptor channels in rat intracardiac ganglion neurons. *Journal of Neurophysiology* 84:1961-1970.
- Cuevas J, Berg DK. 1998. Mammalian nicotinic receptors with alpha7 subunits that slowly desensitize and rapidly recover from alpha-bungarotoxin blockade. *Journal of Neuroscience* 18:10335-10344.
- Cuevas J, Roth AL, Berg DK. 2000. Two distinct classes of functional alpha 7-containing nicotinic receptor on rat superior cervical ganglion neurons. *Journal of Physiology-London* 525:735-746.
- Dalack GW, Healy DJ, Meador-Woodruff JH. 1998. Nicotine dependence in schizophrenia: clinical phenomena and laboratory findings. *American Journal of Psychiatry* 155:1490-1501.
- Dani JA, Heinemann S. 1996. Molecular and cellular aspects of nicotine abuse. *Neuron* 16:905-908.
- De Biasi M, Nigro F, Xu W. 2000. Nicotinic acetylcholine receptors in the autonomic control of bladder function. *European Journal of Pharmacology* 393:137-140.
- Decker MW, Meyer MD. 1999. Therapeutic potential of neuronal nicotinic acetylcholine receptor agonists as novel analgesics. *Biochemical Pharmacology* 58:917-923.
- Decker MW, Brioni JD, Bannon AW, Arneric SP. 1995. Diversity of neuronal nicotinic acetylcholine receptors: lessons from behavior and implications for CNS therapeutics. *Life Sciences* 56:545-570.
- Deneris ES, Francis N, McDonough J, Fyodorov D, Miller T, Yang XD. 2000. Transcriptional control of the neuronal nicotinic acetylcholine receptor gene cluster by the beta 43' enhancer, Sp1, SCIP and ETS transcription factors. *European Journal of Pharmacology* 393:69-74.
- Di Chiara G. 2000. Role of dopamine in the behavioural actions of nicotine related to addiction. *European Journal of Pharmacology* 393:295-314.
- Dineley KT, Patrick JW. 2000. Amino acid determinants of alpha 7 nicotinic acetylcholine receptor surface expression. *Journal of Biological Chemistry* 275:13974-13985.
- Dolezal V, Lee K, Schobert A, Hertting G. 1996. The influx of Ca²⁺ and the release of noradrenaline evoked by the stimulation of presynaptic nicotinic receptors of chick sympathetic neurons in culture are not mediated via L-, N-, or P-type calcium channels. *Brain Research* 740:75-80.
- Drisdel RC, Green WN. 2000. Neuronal α -bungarotoxin receptors are α 7 subunit homomers. *Journal of Neuroscience* 20:133-139.
- Du Q, Tomkinson AE, Gardner PD. 1997. Transcriptional regulation of neuronal nicotinic acetylcholine receptor genes. A possible role for the DNA-binding protein Puralpha. *Journal of Biological Chemistry* 272:14990-14995.
- Du Q, Melnikova IN, Gardner PD. 1998. Differential effects of heterogeneous nuclear ribonucleoprotein K on Sp1- and Sp3-mediated transcriptional activation of a neuronal nicotinic acetylcholine receptor promoter. *Journal of Biological Chemistry* 273:19877-19883.
- Durany N, Zochling R, Boissl KW, Paulus W, Ransmayr G, Tatschner T, Danielczyk W, Jellinger K, Deckert J, Riederer P. 2000. Human post-mortem striatal alpha 4 beta 2 nicotinic acetylcholine receptor density in schizophrenia and Parkinson's syndrome. *Neuroscience Letters* 287:109-112.
- Dursun SM, Reveley MA. 1997. Differential effects of transdermal nicotine on microstructured analyses of tics in Tourette's syndrome: an open study. *Psychological Medicine* 27:483-487.
- Edenberg HJ, Foroud T, Conneally PM, Sorbel JJ, Carr K, Crose C, Willis C, Zhao J, Miller M, Bowman E, Mayeda A, Rau NL, Smiley C, Rice JP, Goate A, Reich T, Stine OC, McMahon F, DePaulo JR, Meyers D, Detera-Wadleigh SD, Goldin LR, Gershon ES, Blehar MC, Nurnberger Jr, Jr. 1997. Initial genomic scan of the NIMH genetics initiative bipolar pedigrees: chromosomes 3, 5, 15, 16, 17, and 22. *American Journal of Medical Genetics* 74:238-246.
- Eisele JL, Bertrand S, Galzi JL, Devillers-Thiery A, Changeux J-P, Bertrand D. 1993. Chimeric nicotinic-serotonergic receptor combines distinct ligand binding and channel specificities. *Nature* 366:479-483.
- Elgoyhen AB, Johnson DS, Boulter J, Vetter DE, Heinemann S. 1994. Alpha 9: an acetylcholine receptor with novel pharmacological properties expressed in rat cochlear hair cells. *Cell* 79:705-715.
- Elliott KJ, Ellis SB, Berckhan KJ, Urrutia A, Chavez-Noriega LE, Johnson EC, Velicelebi G, Harpold MM. 1996. Comparative structure of human neuronal alpha 2-alpha 7 and beta 2-beta 4 nicotinic acetylcholine receptor subunits and functional expression of the alpha 2, alpha 3, alpha 4, alpha 7, beta 2, and beta 4 subunits. *Journal of Molecular Neuroscience* 7:217-228.
- Elmslie FV, Rees M, Williamson MP, Kerr M, Kjeldsen MJ, Pang KA, Sundqvist A, Friis ML, Chadwick D, Richens A, Covanis A, Santos M, Arzimanoglou A, Panayiotopoulos CP, Curtis D, Whitehouse WP, Gardiner RM. 1997. Genetic mapping of a major susceptibility locus for juvenile myoclonic epilepsy on chromosome 15q. *Human Molecular Genetics* 6:1329-1334.
- Engel AG, Ohno K, Milone M, Wang HL, Nakano S, Bouzat C, Pruitt JN, Hutchinson DO, Brengman JM, Bren N, Sieb JP, Sine SM. 1996. New mutations in acetylcholine receptor subunit genes reveal heterogeneity in the slow-channel congenital myasthenic syndrome. *Human Molecular Genetics* 5:1217-1227.
- Fisher JL, Dani JA. 2000. Nicotinic receptors on hippocampal cultures can increase synaptic glutamate currents while decreasing the NMDA-receptor component. *Neuropharmacology* 39:2756-2769.
- Flora A, Schulz R, Benfante R, Battaglini E, Terzano S, Clementi F, Fornasari D. 2000a. Neuronal and extraneuronal expression and regulation of the human alpha 5 nicotinic receptor subunit gene. *Journal of Neurochemistry* 75:18-27.
- Flora A, Schulz R, Benfante R, Battaglini E, Terzano S, Clementi F, Fornasari D. 2000b. Transcriptional regulation of the human alpha 5 nicotinic receptor subunit gene in neuronal and non-neuronal tissues. *European Journal of Pharmacology* 393:85-95.
- Flores CM, Davila-Garcia MI, Ulrich YM, Kellar KJ. 1997. Differential regulation of neuronal nicotinic receptor binding sites following chronic nicotine administration. *Journal of Neurochemistry* 69:2216-2219.
- Fornasari D, Battaglini E, Flora A, Terzano S, Clementi F. 1997. Structural and functional characterization of the human alpha3 nicotinic subunit gene promoter. *Molecular Pharmacology* 51:250-261.
- Fornasari D, Battaglini E, Terzano S, Clementi F. 1999. Transcriptional regulation of neuronal nicotinic receptor subunit genes. In: Arneric SP, Brioni JD, eds. *Neuronal Nicotinic Receptors: Pharmacology and Therapeutic Opportunities*. New York: Wiley-Liss, pp. 25-42.
- Franceschini D, Orr-Urtreger A, Yu W, Mackey LY, Bond RA, Armstrong D, Patrick JW, Beaudet AL, De Biasi M. 2000. Altered baroreflex responses in alpha 7 deficient mice. *Behavioural Brain Research* 113:3-10.
- Francis MM, Vazquez RW, Papke RL, Oswald RE. 2000. Subtype-selective inhibition of neuronal nicotinic acetylcholine receptors by cocaine is determined by the alpha 4 and beta 4 subunits. *Molecular Pharmacology* 58:109-119.
- Frazier CJ, Buhler AV, Weiner JL, Dunwiddie TV. 1998a. Synaptic potentials mediated via alpha-bungarotoxin-sensitive nicotinic acetylcholine receptors in rat hippocampal interneurons. *Journal of Neuroscience* 18:8228-8235.
- Frazier CJ, Rollins YD, Breese CR, Leonard S, Freedman R, Dunwiddie TV. 1998b. Acetylcholine activates an alpha-bungarotoxin-sensitive nicotinic current in rat hippocampal interneurons, but not pyramidal cells. *Journal of Neuroscience* 18:1187-1195.
- Freedman R, Waldo M, Bickford-Wimer P, Nagamoto H. 1991. Elementary neuronal dysfunctions in schizophrenia. *Schizophrenia Research* 4:233-243.
- Freedman R, Hall M, Adler LE, Leonard S. 1995. Evidence in postmortem brain tissue for decreased numbers of hippocampal nicotinic receptors in schizophrenia. *Biological Psychiatry* 38:22-33.
- Freedman R, Coon H, Myles-Worsley M, Orr-Urtreger A, Olincy A, Davis A, Polymeropoulos M, Holik J, Hopkins J, Hoff M, Rosenthal J, Waldo MC, Reimherr R, Wender P, Yaw J, Young DA, Breese CR, Adams C, Patterson D, Adler LE, Kruglyak L, Leonard S, Byerley W. 1997. Linkage of a neurophysiological deficit in schizophrenia to a chromosome 15 locus. *Proceedings of the National Academy of Sciences USA* 94:587-592.
- Fryer JD, Lukas RJ. 1999a. Antidepressants noncompetitively inhibit nicotinic acetylcholine receptor function. *Journal of Neurochemistry* 72:1117-1124.
- Fryer JD, Lukas RJ. 1999b. Noncompetitive functional inhibition at diverse, human nicotinic acetylcholine receptor subtypes by bupropion, phencyclidine, and ibogaine. *Journal of Pharmacology and Experimental Therapeutics* 288:88-92.

- Fu YT, Matta SG, Gao WQ, Brower WG, Sharp BM. 2000. Systemic nicotine stimulates dopamine release in nucleus accumbens: re-evaluation of the role of N-methyl-D-aspartate receptors in the ventral tegmental area. *Journal of Pharmacology & Experimental Therapeutics* 294:458-465.
- Fyodorov D, Nelson T, Deneris E. 1998. Pet-1, a novel ETS domain factor that can activate neuronal nAChR gene transcription. *Journal of Neurobiology* 34:151-163.
- Galzi J-L, Revah F, Bessis A, Changeux J-P. 1991. Functional architecture of the nicotinic acetylcholine receptor: from electric organ to brain. *Annual Review of Pharmacology* 31:37-72.
- Gault J, Robinson M, Berger R, Drebing C, Logel J, Hopkins J, Moore T, Jacobs S, Meriwether J, Choi MJ, Kim EJ, Walton K, Buiting K, Davis A, Breese CR, Freedman R, Leonard S. 1998. Genomic organization and partial duplication of the human $\alpha 7$ neuronal nicotinic acetylcholine receptor gene. *Genomics* 52:173-185.
- Gerzanich V, Anand R, Lindstrom J. 1994. Homomers of alpha 8 and alpha 7 subunits of nicotinic receptors exhibit similar channel but contrasting binding site properties. *Molecular Pharmacology* 45:212-220.
- Girod R, Crabtree G, Ernstrom G, Ramirez-Latorre J, McGehee D, Turner J, Role L. 1999. Heteromeric complexes of alpha 5 and/or alpha 7 subunits: effects of calcium and potential role in nicotine-induced presynaptic facilitation. *Annals of the New York Academy of Sciences* 868:578-590.
- Glassman AH, Helzer JE, Covey LS, Cottler LB, Stetner F, Tipp JE, Johnson J. 1990. Smoking, smoking cessation, and major depression. *Journal of the American Medical Association* 264:1546-1549.
- Glassman AH, Covey LS, Dalack GW, Stetner F. 1992. Cigarette smoking, major depression, and schizophrenia. *Clinical Neuropsychopharmacology* 15 Suppl 1 Pt A:560-561A.
- Goldner FM, Dineley KT, Patrick JW. 1997. Immunohistochemical localization of the nicotinic acetylcholine receptor subunit alpha 6 to dopaminergic neurons in the substantia nigra and ventral tegmental area. *Neuroreport* 8:2739-2742.
- Gopalakrishnan M, Buisson B, Touma E, Giordano T, Campbell JE, Hu IC, Donnelly-Roberts D, Americ SP, Bertrand D, Sullivan JP. 1995. Stable expression and pharmacological properties of the human alpha 7 nicotinic acetylcholine receptor. *European Journal of Pharmacology* 290:237-246.
- Gotti C, Fornasari D, Clementi F. 1997. Human neuronal nicotinic receptors. *Progress in Neurobiology* 53:199-237.
- Grady S, Marks MJ, Wonnacott S, Collins AC. 1992. Characterization of nicotinic receptor-mediated [3 H]dopamine release from synaptosomes prepared from mouse striatum. *Journal of Neurochemistry* 59:848-856.
- Grady SR, Marks MJ, Collins AC. 1994. Desensitization of nicotine-stimulated [3 H]dopamine release from mouse striatal synaptosomes. *Journal of Neurochemistry* 62:1390-1398.
- Grando SA, Horton RM, Pereira EFR, Diethelmokita BM, George PM, Albuquerque EX, Contifine BM. 1995. A nicotinic acetylcholine-receptor regulating cell-adhesion and motility is expressed in human keratinocytes. *Journal of Investigative Dermatology* 105:774-781.
- Green WN, Millar NS. 1995. Ion-channel assembly. *Trends in Neuroscience* 18:280-287.
- Guan ZZ, Zhang X, Blennow K, Nordberg A. 1999. Decreased protein level of nicotinic receptor alpha 7 subunit in the frontal cortex from schizophrenic brain. *Neuroreport* 10:1779-1782.
- Guo JZ, Tredway TL, Chiappinelli VA. 1998. Glutamate and GABA release are enhanced by different subtypes of presynaptic nicotinic receptors in the lateral geniculate nucleus. *Journal of Neuroscience* 18:1963-1969.
- Haghighi AP, Cooper E. 2000. A molecular link between inward rectification and calcium permeability of neuronal nicotinic acetylcholine alpha3beta4 and alpha4beta2 receptors. *Journal of Neuroscience* 20:529-541.
- Han ZY, Le Novere N, Zoli M, Hill JA, Champtiaux N, Changeux JP. 2000. Localization of nAChR subunit mRNAs in the brain of macaca mulatta. *European Journal of Neuroscience* 12:3664-3674.
- Hefft S, Hulo S, Bertrand D, Muller D. 1999. Synaptic transmission at nicotinic acetylcholine receptors in rat hippocampal organotypic cultures and slices. *Journal of Physiology* 515:769-776.
- Hernandez SC, Bertolino M, Xiao YX, Pringle KE, Caruso FS, Kellar KJ. 2000. Dextromethorphan and its metabolite dextrorphan block alpha 3 beta 4 neuronal nicotinic receptors. *Journal of Pharmacology & Experimental Therapeutics* 293:962-967.
- Hiemke C, Stolp M, Reuss S, Wevers A, Reinhardt S, Maelicke A, Schlegel S, Schroder H. 1996. Expression of alpha subunit genes of nicotinic acetylcholine receptors in human lymphocytes. *Neuroscience Letters* 214:171-174.
- Hilmas C, Alkondon M, Eisenberg HM, Albuquerque EX. 1999. Nicotinic receptor activation modulates GABA release from neurons in human cortical slices. *Society for Neuroscience Abstract* 24:1581.
- Horton RM, Nguyen VT, Grando SA. 1997. A survey of cholinergic receptor types expressed in cultured human keratinocytes. *Journal of Investigative Dermatology* 108:603 (Abstract).
- Hughes JR, Hatsukami DK, Mitchell JE, Dahlgren IA. 1986. Prevalence of smoking among psychiatric outpatients. *American Journal of Psychiatry* 143:603.
- Hughes JR, Higgins ST, Bickel WK. 1994. Nicotine withdrawal versus other drug-withdrawal syndromes - similarities and dissimilarities. *Addiction* 89:1461-1470.
- Ji DY, Dani JA. 2000. Inhibition and disinhibition of pyramidal neurons by activation of nicotinic receptors on hippocampal interneurons. *Journal of Neurophysiology* 83:2682-2690.
- Johnson DS, Martinez J, Elgoyhen AB, Heinemann SF, McIntosh M. 1995. Alpha-Conotoxin Iml exhibits subtype-specific nicotinic acetylcholine receptor blockade: preferential inhibition of homomeric alpha 7 and alpha 9 receptors. *Molecular Pharmacology* 48:194-199.
- Johnson PJ, Shum OR, Thornton PD, Bornstein JC. 1999. Evidence that inhibitory motor neurons of the guinea-pig small intestine exhibit fast excitatory synaptic potentials mediated via P2X receptors. *Neuroscience Letters* 266:169-172.
- Jonas PE, Phannavong B, Schuster R, Schroder C, Gundelfinger ED. 1994. Expression of the ligand-binding nicotinic acetylcholine receptor subunit D alpha 2 in the *Drosophila* central nervous system. *Journal of Neurobiology* 25:1494-1508.
- Jorenby DE, Hatsukami DK, Smith SS, Fiore MC, Allen S, Jensen J, & Baker TB. 1996. Characterization of tobacco withdrawal symptoms: transdermal nicotine reduces hunger and weight gain. *Psychopharmacology* 128:130-138.
- Kaiser S, Wonnacott S. 2000. alpha-Bungarotoxin-sensitive nicotinic receptors indirectly modulate [3 H]dopamine release in rat striatal slices via glutamate release. *Molecular Pharmacology* 58:312-318.
- Kaiser SA, Soliakov L, Harvey SC, Luetje CW, Wonnacott S. 1998. Differential inhibition by alpha-conotoxin-MII of the nicotinic stimulation of [3 H]dopamine release from rat striatal synaptosomes and slices. *Journal of Neurochemistry* 70:1069-1076.
- Kassner PD, Berg DK. 1997. Differences in the fate of neuronal acetylcholine receptor protein expressed in neurons and stably transfected cells. *Journal of Neurobiology* 33:968-982.
- Kaufmann CA, Suarez B, Malaspina D, Pepple J, Svrakic D, Mangel PD, Meyer J, Zambuto CT, Schmitt K, Matise TC, Harkavy FJ, Hampe C, Lee H, Shore D, Wynne D, Faraone SV, Tsuang MT, Cloninger CR. 1998. NIMH Genetics Initiative Millenium Schizophrenia Consortium: linkage analysis of African-American pedigrees. *American Journal of Medical Genetics* 81:282-289.
- Ke L, Lukas RJ. 1996. Effects of steroid exposure on ligand binding and functional activities of diverse nicotinic acetylcholine receptor subtypes. *Journal of Neurochemistry* 67:1100-1112.
- Khakh BS, Zhou XP, Sydes J, Galligan JJ, Lester HA. 2000. State-dependent cross-inhibition between transmitter-gated cation channels. *Nature* 406:405-410.
- Klein J, Gonzales R, Koppen A, Loffelholz K. 1993. Free choline and choline metabolites in rat-brain and body-fluids-sensitive determination and implications for choline supply to the brain. *Neurochemistry International* 22:293-300.
- Koyama K, Sudo K, Nakamura Y. 1994. Mapping of the human nicotinic acetylcholine receptor beta 3 gene (CHRNA3) within chromosome 8p11.2. *Genomics* 21:460-461.
- Krause RM, Buisson B, Bertrand S, Corring PJ, Galzi JL, Changeux JP, Bertrand D. 1998. Ivermectin: a positive allosteric effector of the alpha 7 neuronal nicotinic acetylcholine receptor. *Molecular Pharmacology* 53:283-294.
- Krause W, Michael N, Lubke C, Livett BG, Oehme P. 1997. Substance p and epibatidine-evoked catecholamine release from fractionated chromaffin cells. *European Journal of Pharmacology* 328:249-254.
- Kuryatov A, Olale F, Cooper J, Choi C, Lindstrom J. 2000. Human alpha6 AChR subtypes: subunit composition, assembly and pharmacological responses. *Neuropharmacology* 39:2570-2590.

- Le Novère N, Changeux J-P. 1995. Molecular evolution of the nicotinic acetylcholine receptor: an example of multigene family in excitable cells. *Journal of Molecular Evolution* 40:155–172.
- Le Novère N, Zoli M, Changeux J-P. 1996. Neuronal nicotinic receptor alpha 6 subunit mRNA is selectively concentrated in catecholaminergic nuclei of the rat brain. *European Journal of Neuroscience* 8:2428–2439.
- Le Novère N, Zoli M, Lena C, Ferrari R, Picciotto MR, Merlo-Pich E, Changeux J-P. 1999. Involvement of alpha 6 nicotinic receptor subunit in nicotine-elicited locomotion, demonstrated by in vivo antisense oligonucleotide infusion. *Neuroreport* 10:2497–2501.
- Lebargy F, Benhamou K, Morin D, Zini R, Urien S, Bree F, Bignon J, Branellec A, Lagrue G. 1996. Tobacco smoking induces expression of very-high-affinity nicotine binding sites on blood polymorphonuclear cells. *American Journal of Respiratory & Critical Care Medicine* 153:1056–1063.
- Lee MJ, Breese CR, Leonard S. 2000. The effect of nicotine and haloperidol co-treatment on high and low affinity nicotinic and dopamine receptor levels in the rat brain. *Brain Research* (in press).
- Lena C, Changeux J-P. 1997. Role of Ca^{2+} ions in nicotinic facilitation of GABA release in mouse thalamus. *Journal of Neuroscience* 17:576–585.
- Lena C, d'Exaerde AD, Cordero-Erausquin M, Le Novère N, Arroyo-Jimenez MD, Changeux J-P. 1999. Diversity and distribution of nicotinic acetylcholine receptors in the locus ceruleus neurons. *Proceedings of the National Academy of Sciences USA* 96:12126–12131.
- Leonard S, Gault J, Adams C, Breese CR, Rollins Y, Adler LE, Olincy A, Freedman R. 1998a. Nicotinic receptors, smoking and schizophrenia. *Restorative Neurology & Neuroscience* 12:195–201.
- Leonard S, Gault J, Moore T, Hopkins J, Robinson M, Olincy A, Adler LE, Cloninger CR, Kaufmann CA, Tsuang MT, Faraone SV, Malaspina D, Svarkic DM, Freedman R. 1998b. Further investigation of a chromosome 15 locus in schizophrenia: analysis of affected sibpairs from the NIMH Genetics Initiative. *American Journal of Medical Genetics* 81:308–312.
- Levin ED, Simon BB. 1998. Nicotinic acetylcholine involvement in cognitive function in animals. *Psychopharmacology* 138:217–230.
- Levin ED, Bettgowda C, Weaver T, Christopher NC. 1998. Nicotine-dizocilpine interactions and working and reference memory performance of rats in the radial-arm maze. *Pharmacology, Biochemistry & Behavior* 61:335–340.
- Li Y, Papke RL, He YJ, Millard WJ, Meyer EM. 1999. Characterization of the neuroprotective and toxic effects of alpha7 nicotinic receptor activation in PC12 cells. *Brain Research* 830:218–225.
- Lindstrom J. 1996. Neuronal nicotinic acetylcholine receptors. In T. Narahashi, ed. *Ion Channels*. New York: Plenum Press, pp. 377–450.
- Liu QS, Berg DK. 1999. Actin filaments and the opposing actions of CaM kinase II and calcineurin in regulating alpha 7-containing nicotinic receptors on chick ciliary ganglion neurons. *Journal of Neuroscience* 19:10280–10288.
- Lobos EA. 1993. Five subunit genes of the human muscle nicotinic acetylcholine receptor are mapped to two linkage groups on chromosomes 2 and 17. *Genomics* 17:642–650.
- Lueders KK, Elliott RW, Marenholz I, Mischke D, DuPree M, Hamer D. 1999. Genomic organization and mapping of the human and mouse neuronal beta2-nicotinic acetylcholine receptor genes. *Mammalian Genome* 10:900–905.
- Luetje CW, Patrick J. 1991. Both α - and β -subunits contribute to the agonist sensitivity of neuronal nicotinic acetylcholine receptors. *Journal of Neuroscience* 11:837–845.
- Luetje CW, Maddox FN, Harvey SC. 1998. Glycosylation within the cysteine loop and six residues near conserved Cys192/Cys193 are determinants of neuronal bungarotoxin sensitivity on the neuronal nicotinic receptor alpha3 subunit. *Molecular Pharmacology* 53:1112–1119.
- Lukas RJ, Bencherif M. 1992. Heterogeneity and regulation of nicotinic acetylcholine receptors. *International Review of Neurobiology* 34:25–131.
- Lukas RJ, Eisenhour CM. 1996. Interactions between tachykinins and diverse, human nicotinic acetylcholine receptor subtypes. *Neurochemical Research* 21:1245–1257.
- Lukas RJ, Norman SA, Lucero L. 1993. Characterization of nicotinic acetylcholine receptors expressed by cells of the SH-SY5Y human neuroblastoma clonal line. *Molecular & Cellular Neuroscience* 4:1–12.
- MacDermott AB, Role LW, Siegelbaum SA. 1999. Presynaptic ionotropic receptors and the control of transmitter release. *Annual Review of Neuroscience* 22:443–485.
- Macklin KD, Maus ADJ, Pereira EFR, Albuquerque EX, Conti-Fine BM. 1998. Human vascular endothelial cells express functional nicotinic acetylcholine receptors. *Journal of Pharmacology & Experimental Therapeutics* 287:435–439.
- Maelicke A, Albuquerque EX. 2000. Allosteric modulation of nicotinic acetylcholine receptors as a treatment strategy for Alzheimer's disease. *European Journal of Pharmacology* 393:165–170.
- Mansvelder HD, McGehee DS. 2000. Long-term potentiation of excitatory inputs to brain reward areas by nicotine. *Neuron* 27:349–357.
- Marchi M, Raiteri M. 1996. Nicotinic autoreceptors mediating enhancement of acetylcholine release become operative in conditions of "impaired" cholinergic presynaptic function. *Journal of Neurochemistry* 67:1974–1981.
- Marks MJ, Stitzel JA, Collins AC. 1986. A dose-response analysis of nicotine tolerance and receptor changes in two inbred mouse strains. *Journal of Pharmacology & Experimental Therapeutics* 239:358–364.
- Marks MJ, Smith KW, Collins AC. 1998. Differential agonist inhibition identifies multiple epibatidine binding sites in mouse brain. *Journal of Pharmacology & Experimental Therapeutics* 285:377–386.
- Marubio LM, Arroyo-Jimenez MD, Cordero-Erausquin M, Lena C, Le Novère N, d'Exaerde AD, Huchet M, Damaj MI, Changeux JP. 1999. Reduced antinociception in mice lacking neuronal nicotinic receptor subunits. *Nature* 398:805–810.
- Marutle A, Warpmann U, Bogdanovic N, Nordberg A. 1998. Regional distribution of subtypes of nicotinic receptors in human brain and effect of aging studied by (+/-)-[³H]epibatidine. *Brain Research* 801:143–149.
- Matter J-M, Matter-Sadzinski L, Roztocil T, Hernandez M-C, Couturier S, Ong MT, Ballivet M. 1998. On the transcriptional regulation of neuronal nAChR genes. *Journal of Physiology (Paris)* 92:245–248.
- Maus ADJ, Pereira EFR, Karachunski PI, Horton RM, Navaneetham D, Macklin K, Cortes WS, Albuquerque EX, Conti-Fine BM. 1998. Human and rodent bronchial epithelial cells express functional nicotinic acetylcholine receptors. *Molecular Pharmacology* 54:779–788.
- McDonough J, Deneris E. 1997. Beta43': An enhancer displaying neural-restricted activity is located in the 3'-untranslated exon of the rat nicotinic acetylcholine receptor beta4 gene. *Journal of Neuroscience* 17:2273–2283.
- McEvoy J, Freudenreich O, McGee M, Vanderzwaag C, Levin, Rose J. 1995. Clozapine decreases smoking in patients with chronic schizophrenia. *Biological Psychiatry* 37:550–552.
- McGehee DS, Role LW. 1995. Physiological diversity of nicotinic acetylcholine receptors expressed by vertebrate neurons. *Annual Review of Physiology* 57:521–546.
- McGehee DS, Role LW. 1996. Presynaptic ionotropic receptors. *Current Opinion in Neurobiology* 6:342–349.
- McIntosh JM, Santos AD, Olivera BM. 1999. Conus peptides targeted to specific nicotinic acetylcholine receptor subtypes. *Annual Review of Biochemistry* 68:59–88.
- McKenna JP, Cox JL. 1992. Transdermal nicotine replacement and smoking cessation. *American Family Physician* 45:2595–2601.
- Melnikova IN, Yang Y, Gardner PD. 2000. Interactions between regulatory proteins that bind to the nicotinic receptor beta 4 subunit gene promoter. *European Journal of Pharmacology* 393:75–83.
- Messina ES, Tyndale RF, Sellers EM. 1997. A major role for CYP2A6 in nicotine C-oxidation by human liver microsomes. *Journal of Pharmacology & Experimental Therapeutics* 282:1608–1614.
- Meyer EM, Tay ET, Papke RL, Meyers C, Huang GL, de Fiebre CM. 1997. 3-[2,4-Dimethoxybenzylidene]anabaseine (DMXB) selectively activates rat alpha7 receptors and improves memory-related behaviors in a mecamylamine-sensitive manner. *Brain Research* 768:49–56.
- Meyer EM, King MA, Meyers C. 1998. Neuroprotective effects of 2,4-dimethoxybenzylidene anabaseine (DMXB) and tetrahydroaminoacridine (THA) in neocortices of nucleus basalis lesioned rats. *Brain Research* 786:252–254.
- Mirza NR, Stolerman IP. 1998. Nicotine enhances sustained attention in the rat under specific task conditions. *Psychopharmacology* 138:266–274.

- Misu Y, Nakamura S, Goshima Y, Yue JL, Miyamae T, Kubo T. 1993. Review on the relationship between nicotinic acetylcholine receptors and dopaminergic neurotransmission in the central nervous system—dopa is an endogenous neuroactive substance. *Japanese Journal of Psychopharmacology* 13:199–210.
- Miyazawa A, Fujiyoshi Y, Stowell M, Unwin N. 1999. Nicotinic acetylcholine receptor at 4.6 Å resolution: transverse tunnels in the channel wall. *Journal of Molecular Biology* 288:765–786.
- Monod J, Wyman J, Changeux JP. 1965. On the nature of allosteric transitions: a plausible model. *Journal of Molecular Biology* 12:88–118.
- Monteggia LM, Gopalakrishnan M, Touma E, Idler KB, Nash H, Arneric SP, Sullivan JP, Giordano T. 1995. Cloning and transient expression of genes encoding the human alpha 4 and beta 2 neuronal nicotinic acetylcholine receptor (nAChR) subunits. *Gene* 155:189–193.
- Nagamoto HT, Adler LE, Hea RA, Griffith JM, McRae KA, Freedman R. 1996. Gating of auditory P50 in schizophrenics: unique effects of clozapine. *Biological Psychiatry* 40:181–188.
- Nelson ME, Lindstrom J. 1999. Single channel properties of human alpha 3 AChRs: impact of beta 2, beta 4 and alpha 5 subunits. *Journal of Physiology (London)* 516:657–678.
- Nguyen VT, Hall LL, Gallacher G, Ndoye A, Jolkovsky DL, Webber RJ, Buchli R, Grando SA. 2000. Choline acetyltransferase, acetylcholinesterase, and nicotinic acetylcholine receptors of human gingival and esophageal epithelia. *Journal of Dental Research* 79:939–949.
- Nisell M, Marcus M, Nomikos GG, Svensson TH. 1997. Differential effects of acute and chronic nicotine on dopamine output in the core and shell of the rat nucleus accumbens. *Journal of Neural Transmission* 104:1–10.
- Nisell M, Nomikos GG, Svensson TH. 1994. Systemic nicotine-induced dopamine release in the rat nucleus-accumbens is regulated by nicotinic receptors in the ventral tegmental area. *Synapse* 16:36–44.
- Nishizaki T, Matsuoka T, Nomura T, Kondoh T, Watabe S, Shiotani T, Yoshii M. 2000. Presynaptic nicotinic acetylcholine receptors as a functional target of nefiracetam in inducing a long-lasting facilitation of hippocampal neurotransmission. *Alzheimer Disease & Associated Disorders* 14:S82–S94.
- Nomura T, Nishizaki T. 2000. Nefiracetam facilitates hippocampal neurotransmission by a mechanism independent of the piracetam and aniracetam action. *Brain Research* 870:157–162.
- Ohno K, Hutchinson DO, Milone M, Brengman JM, Bouzat C, Sine SM, Engel AG. 1995. Congenital myasthenic syndrome caused by prolonged acetylcholine receptor channel openings due to a mutation in the M2 domain of the epsilon subunit. *Proceedings of the National Academy of Sciences USA* 92:758–762.
- Ohno K, Wang HL, Milone M, Bren N, Brengman JM, Nakano Quiram P, Pruitt JN, Sine SM, Engel AG. 1996. Congenital myasthenic syndrome caused by decreased agonist binding affinity due to a mutation in the acetylcholine receptor epsilon subunit. *Neuron* 17:157–170.
- Ohno K, Quiram PA, Milone M, Wang HL, Harper MC, Pruitt JN, Brengman JM, Pao L, Fischbeck KH, Crawford TO, Sine S, Engel AG. 1997. Congenital myasthenic syndromes due to heteroallelic nonsense/missense mutations in the acetylcholine receptor epsilon subunit gene: identification and functional characterization of six new mutations. *Human Molecular Genetics* 6:753–766.
- Ohno K, Anlar B, Ozdirim E, Brengman JM, Engel AG. 1998. Frameshifting and splice-site mutations in the acetylcholine receptor epsilon subunit gene in three Turkish kinships with congenital myasthenic syndromes. *Annals of the New York Academy of Sciences* 841:189–194.
- Orr-Urtreger A, Seldin MF, Baldini A, Beaudet AL. 1995. Cloning and mapping of the mouse alpha 7-neuronal nicotinic acetylcholine receptor. *Genomics* 26:399–402.
- Orr-Urtreger A, Goldner FM, Saeki M, Lorenzo I, Goldberg L, De Biasi M, Dani JA, Patrick JW, Beaudet AL. 1997. Mice deficient in the alpha 7 neuronal nicotinic acetylcholine receptor lack alpha-bungarotoxin binding sites and hippocampal fast nicotinic currents. *Journal of Neuroscience* 17:9165–9171.
- Orr-Urtreger A, Broide RS, Kasten MR, Dani JA, Beaudet AL, Patrick JW. 2000. Mice homozygous for the L250T mutation in the $\alpha 7$ nicotinic acetylcholine receptor show increased neuronal apoptosis and die within one day of birth. *Journal of Neurochemistry* 74:2154–2166.
- Oyaizu M, Narahashi T. 1999. Modulation of the neuronal nicotinic acetylcholine receptor-channel by the nootropic drug nefiracetam. *Brain Research* 822:72–79.
- Papke RL, Bencherif M, Lippiello P. 1996. An evaluation of neuronal nicotinic acetylcholine receptor activation by quaternary nitrogen compounds indicates that choline is selective for the alpha 7 subtype. *Neuroscience Letters* 213:201–204.
- Paradiso K, Sabey K, Evers AS, Zorumski CF, Covey DF, Steinbach JH. 2000. Steroid inhibition of rat neuronal nicotinic alpha 4 beta 2 receptors expressed in HEK 293 cells. *Molecular Pharmacology* 58:341–351.
- Pattyn A, Morin X, Cremer H, Goridis C, Brunet JF. 1999. The homeobox gene Phox2b is essential for the development of autonomic neural crest derivatives. *Nature* 399:366–370.
- Pauly JR, Grun EU, Collins AC. 1990. Chronic corticosterone administration modulates nicotine sensitivity and brain nicotinic receptor binding in C3H mice. *Psychopharmacology* 101:310–316.
- Pauly JR, Marks MJ, Robinson SF, van de Kamp JL, Collins AC. 1996. Chronic nicotine and mecamylamine treatment increase brain nicotinic receptor binding without changing alpha 4 or beta 2 mRNA levels. *Journal of Pharmacology & Experimental Therapeutics* 278:361–369.
- Peng X, Katz M, Gerzanich V, Anand R, Lindstrom J. 1994a. Human alpha 7 acetylcholine receptor: Cloning of the alpha 7 subunit from the SH-SY5Y cell line and determination of pharmacological properties of native receptors and functional alpha 7 homomers expressed in *Xenopus* oocytes. *Molecular Pharmacology* 45:546–554.
- Peng X, Gerzanich V, Anand R, Whiting PJ, Lindstrom J. 1994b. Nicotine-induced increase in neuronal nicotinic receptors results from a decrease in the rate of receptor turnover. *Molecular Pharmacology* 46:523–530.
- Peng X, Gerzanich V, Anand R, Wang F, Lindstrom J. 1997. Chronic nicotine treatment up-regulates alpha3 and alpha7 acetylcholine receptor subtypes expressed by the human neuroblastoma cell line SH-SY5Y. *Molecular Pharmacology* 51:776–784.
- Perry DC, Davila-Garcia MI, Stockmeier CA, Kellar KJ. 1999. Increased nicotinic receptors in brains from smokers: membrane binding and autoradiography studies. *Journal of Pharmacology & Experimental Therapeutics* 289:1545–1552.
- Perry EK, Morris CM, Court JA, Cheng A, Fairbairn AF, McKeith IG, Irving D, Brown A, Perry RH. 1995. Alteration in nicotine binding-sites in Parkinson's disease, Lewy body dementia and Alzheimer's disease – possible index of early neuropathology. *Neuroscience* 64:385–395.
- Pianezza ML, Sellers EM, Tyndale RF. 1998. Nicotine metabolism defect reduces smoking. *Nature* 393:750.
- Picciotto MR, Zoli M, Lena C, Bessis A, Lallemand Y, Lenovere N, Vincent P, Pich EM, Brulet P, Changeux JP. 1995. Abnormal avoidance learning in mice lacking functional high-affinity nicotine receptors in the brain. *Nature* 374:65–67.
- Picciotto MR, Zoli M, Rimondini R, Lena C, Marubio LM, Pich EM, Fuxe K, Changeux J-P. 1998. Acetylcholine receptors containing the beta2 subunit are involved in the reinforcing properties of nicotine. *Nature* 391:173–177.
- Pidoplichko VI, DeBiasi M, Williams JT, Dani JA. 1997. Nicotine activates and desensitizes midbrain dopamine neurons. *Nature* 390:401–404.
- Popot JL, Changeux J-P. 1984. Nicotinic receptor of acetylcholine: structure of an oligomeric integral membrane protein. *Physiology Review* 64:1162–1239.
- Potter A, Corwin J, Lang J, Piasecki M, Lenox R, Newhouse PA. 1999. Acute effects of the selective cholinergic channel activator (nicotinic agonist) ABT-418 in Alzheimer's disease. *Psychopharmacology* 142:334–342.
- Prochazka AV, Weaver MJ, Keller RT, Fryer GE, Licari PA, Lofaso D. 1998. A randomized trial of nortriptyline for smoking cessation. *Archives of Internal Medicine* 158:2035–2039.
- Puchacz E, Buisson B, Bertrand D, Lukas RJ. 1994. Functional expression of nicotinic acetylcholine receptors containing rat alpha 7 subunits in human SH-SY5Y neuroblastoma cells. *FEBS Letters* 354:155–159.
- Quik M, Jeyarasasingam G. 2000. Nicotinic receptors and Parkinson's disease. *European Journal of Pharmacology* 393:223–230.
- Quik M, Choremis J, Komourian J, Lukas RJ, Puchacz E. 1996. Similarity between rat brain nicotinic alpha-bungarotoxin receptors and stably expressed alpha-bungarotoxin binding sites. *Journal of Neurochemistry* 67:145–154.

- Ragozzino D, Barabino B, Fucile S, Eusebi F. 1998. Ca^{2+} permeability of mouse and chick nicotinic acetylcholine receptors expressed in transiently transfected human cells. *Journal of Physiology (London)* 507:749-757.
- Raimondi E, Rubboli F, Moralli D, Chini B, Fornasari D, Tarroni P, De Carli L, Clementi F. 1992. Chromosomal localization and physical linkage of the genes encoding the human alpha 3, alpha 5, and beta 4 neuronal nicotinic receptor subunits. *Genomics* 12:849-850.
- Rang HP, Ritter JM. 1970. On the mechanism of desensitization at cholinergic receptors. *Molecular Pharmacology* 6:357-382.
- Raskind MA, Peskind ER, Wessel T, Yuan W. 2000. Galantamine in AD - a 6-month randomized, placebo-controlled trial with a 6-month extension. *Neurology* 54:2261-2268.
- Rempel N, Heyers S, Engels H, Slegers E, Steinlein OK. 1998. The structures of the human neuronal nicotinic acetylcholine receptor beta2- and alpha3-subunit genes (CHRNA2 and CHRNA3). *Human Genetics* 103:645-653.
- Revah F, Bertrand D, Galzi JL, Devillers-Thiery A, Mulle C, Hussy N, Bertrand S, Ballivet M, Changeux J-P. 1991. Mutations in the channel domain alter desensitization of a neuronal nicotinic receptor. *Nature* 353, 846-849.
- Rickert KW, Imperiali B. 1995. Analysis of the conserved glycosylation site in the nicotinic acetylcholine receptor: potential roles in complex assembly. *Chemistry & Biology* 2:751-759.
- Riley BP, Makoff AM, Magudi-Carter M, Jenkins TJ, Williamson R, Collier DA, Murray RM. 2000. Haplotype transmission disequilibrium and evidence for linkage of the CHRNA7 gene region to schizophrenia in Southern African Bantu families. *American Journal of Medical Genetics* 96:196-201.
- Roerig B, Nelson DA, Katz LC. 1997. Fast synaptic signaling by nicotinic acetylcholine and serotonin 5-HT3 receptors in developing visual cortex. *Journal of Neuroscience* 17:8353-8362.
- Rogers M, Dani JA. 1995. Comparison of quantitative calcium flux through NMDA, ATP, and ACh receptor channels. *Biophysical Journal* 68:501-506.
- Rogers SW, Mandelzys A, Deneris ES, Cooper E, Heinemanns S. 1992. The expression of nicotinic acetylcholine receptors by PC12 cells treated with NGF. *Journal of Neuroscience* 12:4611-4623.
- Role LW. 1992. Diversity in primary structure and function of neuronal nicotinic acetylcholine receptor channels. *Current Opinion in Neurobiology* 2:254-262.
- Romano SJ, Pugh PC, McIntosh JM, Berg DK. 1997. Neuronal-type acetylcholine receptors and regulation of alpha 7 gene expression in vertebrate skeletal muscle. *Journal of Neurobiology* 32:69-80.
- Rose JE, Behm FM, Westman EC. 1998. Nicotine-mecamylamine treatment for smoking cessation: the role of pre-cessation therapy. *Experimental & Clinical Psychopharmacology* 6:331-343.
- Ross SA, Wong JYF, Clifford JJ, Kinsella A, Massalas JS, Horne MK, Scheffer IE, Kola I, Waddington JL, Berkovic SF, Drago J. 2000. Phenotypic characterization of an $\alpha 4$ neuronal nicotinic acetylcholine receptor subunit knock-out mouse. *Journal of Neuroscience* 20:6431-6441.
- Rubboli F, Court JA, Sala C, Morris C, Perry E, Clementi F. 1994. Distribution of neuronal nicotinic receptor subunits in human brain. *Neurochemistry International* 25:69-71.
- Sacaan AI, Reid RT, Santori EM, Adams P, Correa LD, Mahaffy LS, Bleicher L, Cosford ND, Stauderman KA, McDonald IA, Rao TS, Lloyd GK. 1997. Pharmacological characterization of SIB-1765F: a novel cholinergic ion channel agonist. *Journal of Pharmacology & Experimental Therapeutics* 280:373-383.
- Sanderson EM, Drasdo AL, McCrear K, Wonnacott S. 1993. Upregulation of nicotinic receptors following continuous infusion of nicotine is brain-region-specific. *Brain Research* 617:349-352.
- Sands SB, Costa AC, Patrick JW. 1993. Barium permeability of neuronal nicotinic receptor alpha 7 expressed in *Xenopus* oocytes. *Biophysical Journal* 65:2614-2621.
- Schaeffer L, Duclert N, Huchet-Dymanus M, Changeux JP. 1998. Implication of a multisubunit Ets-related transcription factor in synaptic expression of the nicotinic acetylcholine receptor. *EMBO Journal* 17:3078-3090.
- Schuller HM. 1991. Receptor-mediated mitogenic signals and lung cancer. *Cancer Cells* 3:496-503.
- Schuller HM, Nylen E, Park P, Becker KL. 1990. Nicotine, acetylcholine and bombesin are trophic growth factors in neuroendocrine cell lines derived from experimental hamster lung tumors. *Life Sciences* 47:571-578.
- Schulz R, Sawruk E, Mulhardt C, Bertrand S, Baumann A, Phannavong B, Betz H, Bertrand D, Gundelfinger ED, Schmitt B. 1998. D alpha3, a new functional alpha subunit of nicotinic acetylcholine receptors from *Drosophila*. *Journal of Neurochemistry* 71:853-862.
- Searl TJ, Redman RS, Silinsky EM. 1998. Mutual occlusion of P2X ATP receptors and nicotinic receptors on sympathetic neurons of the guinea-pig. *Journal of Physiology (London)* 510:783-791.
- Séguéla P, Wadiche J, Dineley-Miller K, Dani JA, Patrick JW. 1993. Molecular cloning, functional properties, and distribution of rat brain $\alpha 7$: a nicotinic cation channel highly permeable to calcium. *Journal of Neuroscience* 13:596-604.
- Sekhon HS, Jia YB, Raab R, Kuryatov A, Pankow JF, Whitsett JA, Lindstrom J, Spindel ER. 1999. Prenatal nicotine increases pulmonary alpha 7 nicotinic receptor expression and alters fetal lung development in monkeys. *Journal of Clinical Investigation* 103:637-647.
- Sellers EM, Kaplan HL, Tyndale RF. 2000a. Inhibition of cytochrome P450 2A6 increases nicotine's oral bioavailability and decreases smoking. *Clinical Pharmacology & Therapeutics* 68:35-43.
- Sellers EM, Zeman MV, Kaplan HL, Tyndale RF. 2000b. Inhibition of cytochrome P450 2A6 (CYP2A6) decreases smoking and activation of the procarcinogen 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK). *Clinical Pharmacology & Therapeutics* 67:OIA3.
- Servent D, Wincklerdiedrich V, Hu HY, Kessler P, Drevet P, Bertrand D, Menez A. 1997. Only snake curaremimetic toxins with a fifth disulfide bond have high affinity for the neuronal alpha-7 nicotinic receptor. *Journal of Biological Chemistry* 272:24279-24286.
- Sharples CGV, Kaiser S, Soliakov L, Marks MJ, Collins AC, Washburn M, Wright E, Spencer JA, Gallagher T, Whiteaker P, Wonnacott S. 2000. UB-165: A novel nicotinic agonist with subtype selectivity implicates the alpha 4 beta 2 subtype in the modulation of dopamine release from rat striatal synaptosomes. *Journal of Neuroscience* 20:2783-2791.
- Shiffman S, Johnston JA, Khayrallah M, Elash CA, Gwaltney CJ, Paty JA, Gnys M, Evoniuk G, DeVaugh-Geiss J. 2000. The effect of bupropion on nicotine craving and withdrawal. *Psychopharmacology* 148:33-40.
- Shoaib M, Stolerman IP. 1999. Plasma nicotine and cotinine levels following intravenous nicotine self-administration in rats. *Psychopharmacology* 143:318-321.
- Shoop RD, Yamada N, Berg DK. 2000. Cytoskeletal links of neuronal acetylcholine receptors containing alpha 7 subunits. *Journal of Neuroscience* 20:4021-4029.
- Shytle RD, Silver AA, Philipp MK, McConville BJ, Sanberg PR. 1996. Transdermal nicotine for Tourette's syndrome. *Drug Development Research* 38:290-298.
- Silver AA, Shytle RD, Sanberg PR. 2000. Mecamylamine in Tourette's syndrome: a two-year retrospective case study. *Journal of Child and Adolescent Psychopharmacology* 10:59-68.
- Spang JE, Bertrand S, Westera G, Patt JT, Schubiger PA, Bertrand D. 2000. Chemical modification of epibatidine causes a switch from agonist to antagonist and modifies its selectivity for neuronal nicotinic acetylcholine receptors. *Chemistry & Biology* 7:545-555.
- Squire MD, Tornoe C, Baylis HA, Fleming JT, Barnard EA, Sattelle DB. 1995. Molecular cloning and functional co-expression of a *Caenorhabditis elegans* nicotinic acetylcholine receptor subunit (acr-2). *Receptors & Channels* 3:107-115.
- Stafford GA, Oswald RE, Weiland GA. 1994. The beta-subunit of neuronal nicotinic acetylcholine-receptors is a determinant of the affinity for substance-P inhibition. *Molecular Pharmacology* 45:758-762.
- Stafford GA, Oswald RE, Figl A, Cohen BN, Weiland GA. 1998. Two domains of the beta subunit of neuronal nicotinic acetylcholine receptors contribute to the affinity of substance P. *Journal of Pharmacology & Experimental Therapeutics* 286:619-626.
- Stassen HH, Bridler R, Hägele S, Hergersberg M, Mehmman B, Schinzel A, Weisbrod M, Scharfetter C. 2000. Schizophrenia and smoking: evidence for a common neurobiological basis. *American Journal of Medical Genetics (Neuropsychiatric Genetics)* 96:173-177.
- Steinlein O, Weiland S, Stoodt J, Propping P. 1996. Exon-intron structure of the human neuronal nicotinic acetylcholine receptor alpha 4 subunit (CHRNA4). *Genomics* 32:289-294.
- Steinlein OK, Magnusson A, Stoodt J, Bertrand S, Weiland, Berkovic SF, Nakken KO, Propping P, Bertrand D. 1997. An insertion mutation of the CHRNA4 gene in a family with autosomal dominant nocturnal frontal lobe epilepsy. *Human Molecular Genetics* 6:943-947.

- Steinlein OK, Mulley JC, Propping P, Wallace RH, Phillips HA, Sutherland GR, Scheffer IE, Berkovic SF. 1995. A missense mutation in the neuronal nicotinic acetylcholine receptor alpha 4 subunit is associated with autosomal dominant nocturnal frontal lobe epilepsy. *Nature Genetics* 11:201-203.
- Steinlein OK, Stoodt J, Mulley J, Berkovic S, Scheffer IE, Brodtkorb E. 2000. Independent occurrence of the CHRNA4 Ser248Phe mutation in a Norwegian family with nocturnal frontal lobe epilepsy. *Epilepsia* 41:529-535.
- Stevens KE, Kem WR, Mahnir VM, Freedman R. 1998. Selective alpha7-nicotinic agonists normalize inhibition of auditory response in DBA mice. *Psychopharmacology* 136:320-327.
- Stolerman IP, Mirza NR, Hahn B, Shoaib M. 2000. Nicotine in an animal model of attention. *European Journal of Pharmacology* 393:147-154.
- Summers KL, Lippio P, Giacobini E. 1996. A microdialysis study of the effects of the nicotinic agonist RJR-2403 on cortical release of acetylcholine and biogenic amines. *Neurochemical Research* 21:1181-1186.
- Tariot PN, Solomon PR, Morris JC, Kershaw P, Lilienfeld S, Ding C. 2000. A 5-month, randomized, placebo-controlled trial of galantamine in AD. *Neurology* 54:2269-2276.
- Tsuneki H, Klink R, Lena C, Korn H, Changeux JP. 2000. Calcium mobilization elicited by two types of nicotinic acetylcholine receptors in mouse substantia nigra pars compacta. *European Journal of Neuroscience* 12:2475-2485.
- Ullian EM, Sargent, PB. 1995. Pronounced cellular diversity and extrasynaptic location of nicotinic acetylcholine receptor subunit immunoreactivities in the chicken pretectum. *Journal of Neuroscience* 15:7012-7023.
- Unwin N. 1996. Projection structure of the nicotinic acetylcholine receptor: distinct conformations of the alpha subunits. *Journal of Molecular Biology* 257:586-596.
- Valenta DC, Downing JE, Role LW. 1993. Peptide modulation of ACh receptor desensitization controls neurotransmitter release from chicken sympathetic neurons. *Journal of Neurophysiology* 69:928-942.
- Valera S, Ballivet M, Bertrand D. 1992. Progesterone modulates a neuronal nicotinic acetylcholine-receptor. *Proceedings of the National Academy of Sciences USA* 89:9949-9953.
- van Haaren F, Anderson KG, Haworth SC, Kem WR. 1999. GTS-21, a mixed nicotinic receptor agonist/antagonist, does not affect the nicotine cue. *Pharmacology, Biochemistry & Behavior* 64:439-444.
- Vernall AB, Conroy WG, Berg DK. 1993. Neurons assemble acetylcholine receptors with as many as three kinds of subunits while maintaining subunit segregation among receptor subtypes. *Neuron* 10:451-464.
- Vernino S, Amador M, Luetje CW, Patrick J, Dani JA. 1992. Calcium modulation and high calcium permeability of neuronal nicotinic acetylcholine receptors. *Neuron* 8:127-134.
- Vetter DE, Liberman MC, Mann J, Barhanin J, Boulter J, Brown MC, Saffioti-Kolman J, Heinemann SF, Elgoyhen AB. 1999. Role of alpha 9 nicotinic ACh receptor subunits in the development and function of cochlear efferent innervation. *Neuron* 23:93-103.
- Vijayaraghavan S, Pugh PC, Zhang Z-W, Rathouz MM, Berg DK. 1992. Nicotinic receptors that bind α -bungarotoxin on neurons raise intracellular free Ca^{++} . *Neuron* 8:353-362.
- Vincent A, Jacobson L, Plested P, Polizzi A, Tang T, Riemersma S, Newland C, Ghorazian S, Farrar J, MacLennan C, Willcox N, Beeson D, Newsom-Davis J. 1998. Antibodies affecting ion channel function in acquired neuromyotonia, in seropositive and seronegative myasthenia gravis, and in antibody-mediated arthrogryposis multiplex congenita. *Annals of the New York Academy of Sciences* 841:482-496.
- Wang HY, Lee DHS, D'Andrea MR, Peterson PA, Shank RP, Reitz AB. 2000a. beta-amyloid(1-42) binds to alpha 7 nicotinic acetylcholine receptor with high affinity - implications for Alzheimer's disease pathology. *Journal of Biological Chemistry* 275:5626-5632.
- Wang HY, Lee DHS, Davis CB, Shank RP. 2000b. Amyloid peptide A beta(1-42) binds selectively and with picomolar affinity to alpha 7 nicotinic acetylcholine receptors. *Journal of Neurochemistry* 75:1155-1161.
- Watkins SS, Koob GF, Markou A. 2000. Neural mechanisms underlying nicotine addiction: acute positive reinforcement and withdrawal. *Nicotine & Tobacco Research* 2:19-37.
- Weiland S, Bertrand D, Leonard S. 2000. Neuronal nicotinic acetylcholine receptors: from the gene to the disease. *Behavioural Brain Research* 113:43-56.
- Wessler I, Kirkpatrick CJ, Racke K. 1999. The cholinergic 'pitfall': acetylcholine, a universal cell molecule in biological systems, including humans. *Clinical and Experimental Pharmacology & Physiology* 26:198-205.
- Wevers A, Jeske A, Lobron C, Birtsch C, Heinemann S, Maelicke A, Schroder R, Schroder H. 1994. Cellular distribution of nicotinic acetylcholine receptor subunit mRNAs in the human cerebral cortex as revealed by non-isotopic in situ hybridization. *Brain Research, Molecular Brain Research* 25:122-128.
- Wevers A, Sullivan JP, Giordano T, Birtsch C, Monteggia LM, Nowacki S, Arneric SP, Schroder H. 1995. Cellular distribution of the mRNA for the alpha 7 subunit of the nicotinic acetylcholine receptor in the human cerebral cortex. *Drug Development Research* 36:103-110.
- White HK, Levin ED. 1999. Four week nicotine skin patch treatment effects on cognitive performance in Alzheimer's disease. *Psychopharmacology* 143:158-165.
- Whiteaker P, McIntosh JM, Luo SQ, Collins AC, Marks MJ. 2000a. I-125-alpha-conotoxin MII identifies a novel nicotinic acetylcholine receptor population in mouse brain. *Molecular Pharmacology* 57:913-925.
- Whiteaker P, Marks MJ, Grady SR, Lu Y, Picciotto MR, Changeux JP, Collins AC. 2000b. Pharmacological and null mutation approaches reveal nicotinic receptor diversity. *European Journal of Pharmacology* 393:123-135.
- Wonnacott S. 1990. The paradox of nicotinic acetylcholine receptor upregulation by nicotine. *Trends in Pharmacological Science* 11:216-218.
- Wonnacott S. 1997. Presynaptic nicotinic ACh receptors. *Trends in Neurosciences* 20:92-98.
- Wood S, Schertzer M, Yaremko ML. 1995. Identification of the human neuronal nicotinic cholinergic alpha 2 receptor locus, (CHRNA2), within an 8p21 mapped locus, by sequence homology with rat DNA. *Somatic Cell & Molecular Genetics* 21:147-150.
- Xu W, Gelber S, Orr-Urtreger A, Armstrong D, Lewis RA, Ou CN, Patrick, Role L, De Biasi M, Beaudet AL. 1999. Megacystis, mydriasis, and ion channel defect in mice lacking the alpha 3 neuronal nicotinic acetylcholine receptor. *Proceedings of the National Academy of Sciences USA* 96:5746-5751.
- Yakel JL, Jackson MB. 1988. 5-HT₃ receptors mediate rapid responses in cultured hippocampus and a clonal cell line. *Neuron* 1:615-621.
- Yu CR, Role LW. 1998a. Functional contribution of the alpha5 subunit to neuronal nicotinic channels expressed by chick sympathetic ganglion neurons. *Journal of Physiology* 509:667-681.
- Yu CR, Role LW. 1998b. Functional contribution of the alpha7 subunit to multiple subtypes of nicotinic receptors in embryonic chick sympathetic neurons. *Journal of Physiology* 509:651-665.
- Zaninetti M, Tribollet E, Bertrand D, Raggenbass M. 1999. Presence of functional neuronal nicotinic acetylcholine receptors in brainstem motoneurons of the rat. *European Journal of Neuroscience* 11:2737-2748.
- Zevin S, Jacob P, Benowitz NL. 2000. Nicotine-mecamylamine interactions. *Clinical Pharmacology & Therapeutics* 68:58-66.
- Zhang Z-W, Vijayaraghavan S, Berg DK. 1994. Neuronal acetylcholine receptors that bind α -bungarotoxin with high affinity function as ligand-gated ion channels. *Neuron* 12:167-177.
- Zhou X, Galligan JJ. 1998. Non-additive interaction between nicotinic cholinergic and P2X purine receptors in guinea-pig enteric neurons in culture. *Journal of Physiology (London)* 513:685-697.
- Zia S, Ndoye A, Nguyen VT, Grando SA. 1997. Nicotine enhances expression of the alpha 3, alpha 4, alpha 5, and alpha 7 nicotinic receptors modulating calcium metabolism and regulating adhesion and motility of respiratory epithelial cells. *Research Communications In Molecular Pathology & Pharmacology* 97:243-262.
- Zoli M, Lena C, Picciotto MR, Changeux JP. 1998. Identification of four classes of brain nicotinic receptors using beta2 mutant mice. *Journal of Neuroscience* 18:4461-4472.
- Zwart R, Vijverberg HPM. 2000. Potentiation and inhibition of neuronal alpha 4 beta 4 nicotinic acetylcholine receptors by choline. *European Journal of Pharmacology* 393:209-214.