
From: Wagner, Karl A
Sent: Wednesday, January 02, 2008 4:03 PM
To: Dwyer, R.W. (Bill)
Subject: RE: MS markers
Attachments: Analytes.doc

Categories: Confidential
LHS:

Confidential

Hi Bill,

I attached a document with the current PI list. The analytes are broken out by method. The analytes in yellow are typically below LOQ while the analytes in green are sometimes below LOQ.

As I mentioned on the phone, I am interested in proposing a reduced testing list. A few of the analytes that 'are not for free' with the routine method are styrene and acetamide (which are determined with the VOCs). Similarly, the amount of work would be greatly reduced if we could simply look at NNN and NNK and not NPY or NDMA. There is the similar question for HCN and 2-nitropropane as those analytes are determined independently.

Please let me know when you have a few minutes to sit down and talk about this.

Thanks again,
Karl

From: Dwyer, R.W. (Bill)
Sent: Wednesday, January 02, 2008 9:09 AM
To: Wagner, Karl A
Subject: RE: MS markers

Karl,
Yes, I am delighted to talk to you about this topic.
Call me at 43552.
Happy new year.
Bill

From: Wagner, Karl A
Sent: Tuesday, December 18, 2007 2:55 PM
To: Dwyer, R.W. (Bill)
Subject: MS markers

Confidential

Hi Bill,

I was wondering if you could help me with a question: I am looking for information regarding MS analytes that may be used to predict the concentration (or change in concentration) of other analytes of the same chemical class. For example, could BaP be used as a representative marker to predict the concentration of other PAHs in smoke? Would you have a few minutes to talk and/or do you have some good references?

Thanks,
Karl

Karl A. Wagner, Ph.D.
Product Integrity

Philip Morris USA
(804) 274-2405